

## References

- Maurin M, Delbano JN, Mackaya L, Colomb H, Guier C, Mandjee A, et al. Human infection with *Schineria larvae*. Emerg Infect Dis. 2007;13:671–3. <http://dx.doi.org/10.3201/eid1304.061151>
- Roudiere L, Jean-Pierre H, Comte C, Zorogniotti I, Marchandin H, Jumas-Bilak E. Isolation of *Schineria* sp. from a man. Emerg Infect Dis. 2007;13:659–61. <http://dx.doi.org/10.3201/eid1304.061255>
- Barker HS, Snyder JW, Hicks AB, Yanoviak SP, Southern P, Dhakal BK, et al. First case reports of *Ignatzschineria* (*Schineria*) *indica* associated with myiasis. J Clin Microbiol. 2014;52:4432–4. <http://dx.doi.org/10.1128/JCM.02183-14>
- Rådström P, Bäckman A, Qian N, Kraggsbjerg P, Pählson C, Olcén P. Detection of bacterial DNA in cerebrospinal fluid by an assay for simultaneous detection of *Neisseria meningitidis*, *Haemophilus influenzae*, and streptococci using a seminested PCR Strategy. J Clin Microbiol. 1994;32:2738–44.
- Yamamoto S, Harayama S. PCR amplification and direct sequencing of *gyrB* genes with universal primers and their application to the detection and taxonomic analysis of *Pseudomonas putida* strains. Appl Environ Microbiol. 1995;61:1104–9.
- Tóth E, Kovács G, Schumann P, Kovács AL, Steiner U, Halbritter A, et al. *Schineria larvae* gen. nov., sp. nov., isolated from the 1st and 2nd larval stage of *Wohlfahrtia magnifica* (Diptera: Sarcophagidae). Int J Syst Evol Microbiol. 2001; 51:401–7.
- Tóth EM, Borsodi AK, Euzéby JP, Tindall BJ, Márialigeti K. Proposal to replace the illegitimate genus name *Schineria* Tóth et al. 2001 with the genus name *Ignatzschineria* gen. nov. and to replace the illegitimate combination *Schineria larvae* Tóth et al. 2001 with *Ignatzschineria larvae* comb. nov. Int J Syst Evol Microbiol. 2007;57:179–80. <http://dx.doi.org/10.1099/ijs.0.64686-0>
- Gupta AK, Dharme MS, Rangrez AY, Verma P, Ghate HV, Rohde M, et al. *Ignatzschineria indica* sp. nov. and *Ignatzschineria ureiclastica* sp. nov., isolated from adult flesh flies (Diptera: Sarcophagidae). Int J Syst Evol Microbiol. 2011;61:1360–9. <http://dx.doi.org/10.1099/ijs.0.018622-0>
- Tóth EM, Hell É, Kovács G, Borsodi AK, Márialigeti K. Bacteria isolated from the different developmental stages and larval organs of the obligate parasitic fly, *Wohlfahrtia magnifica* (Diptera: Sarcophagidae). Microb Ecol. 2006;51:13–21. <http://dx.doi.org/10.1007/s00248-005-0090-6>

Address for correspondence: Cécile Le Brun, CHU de Tours, Hôpital Bretonneau—Bactériologie, 2 Bd Tonnellé, Tours 37000, France; email: c.lebrun@chu-tours.fr

# etymologia

## *Ignatzschineria* [ig-nat"shi-ner'e-ə]

A genus of aerobic, gram-negative, nonmotile rods, *Ignatzschineria* was first isolated from flies of the family Sarcophagidae (from the Greek *sarco* ["flesh"] + *phage* ["eating"]) by Erika Tóth et al. in 2001. Tóth named the genus after Austrian entomologist Ignaz Rudolph Schiner (1813–1873), who first described the fly *Wohlfahrtia magnifica*. In 2007, Tóth discovered that *Schineria* had already been used for genus of tachina flies and proposed the replacement genus name *Ignatzschineria*.



Ignaz Rudolph Schiner  
Photographer unknown.  
Image via Wikimedia Commons.

## Sources

- Tóth EM, Borsodi AK, Euzéby JP, Tindall BJ, Márialigeti K. Proposal to replace the illegitimate genus name *Schineria* Tóth et al 2001 with the genus name *Ignatzschineria* gen. nov. and to replace the illegitimate combination *Schineria larvae* Tóth et al 2001 with *Ignatzschineria larvae* comb. nov. Int J Syst Evol Microbiol. 2007;57:179–80. <http://dx.doi.org/10.1099/ijs.0.64686-0>
- Tóth E, Kovács G, Schumann P, Kovács AL, Steiner U, Halbritter A, et al. *Schineria larvae* gen. nov., sp. nov., isolated from the 1st and 2nd larval stages of *Wohlfahrtia magnifica* (Diptera: Sarcophagidae). Int J Syst Evol Microbiol. 2001;51:401–7.

Address for correspondence: Ronnie Henry, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop E03, Atlanta, GA 30329-4027, USA; email: boq3@cdc.gov

DOI: <http://dx.doi.org/10.3201/eid2110.ET2110>