Article DOI: https://doi.org/10.3201/eid2911.230814

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

## Racial and Socioeconomic Equity of Tecovirimat Treatment during the 2022 Mpox Emergency, New York, New York, USA

## **Appendix**

**Appendix Table.** Cumulative percentages of persons with mpox treated with tecovirimat during 2-week intervals in study of racial and socioeconomic equity of tecovirimat treatment during 2022 mpox emergency, New York City, New York, USA, May 19, 2022–October 29, 2022\*

		Race/ethnicity					Neighborhood poverty level†			
		Asian/Pacific	Black/African	Hispanic/					•	
End dates	Overall	Islander/other	American	Latino	White	Unknown	Low	Medium	High	Very high
May 28	0	0	0	NI	0	0	0	0	0	NI
Jun 11	0	0	0	0	0	0	0	0	0	NI
Jun 25	11.3	0	0	15.0	13.2	0	15.0	9.7	11.8	0.0
Jul 9	11.0	0	7.9	12.6	13.4	9.7	13.3	11.1	10.6	7.3
Jul 23	18.3	19.7	17.0	15.7	22.9	14.2	18.6	17.6	18.7	20.2
Aug 6	27.3	23.2	24.9	25.3	34.4	22.1	24.8	28.0	28.0	27.2
Aug 20	29.3	28.1	25.8	27.4	37.7	25.3	27.7	29.7	30.3	29.0
Sep 3	30.5	28.2	28.1	28.3	38.3	27.1	29.1	30.9	30.9	30.7
Sep 17	31.0	27.5	29.9	28.8	38.5	26.5	29.0	31.4	31.9	31.1
Oct 1	31.5	28.5	30.2	29.7	38.6	26.8	29.6	31.8	32.7	31.1
Oct 15	31.7	28.9	30.5	30.0	38.5	27.2	29.7	31.7	33.4	31.6
Oct 29	31.8	29.1	30.6	30.2	38.6	26.9	30.1	31.6	33.4	32.0
Total±	32.4	30.1	31.3	31.0	38.8	27.7	30.6	32.0	33.9	33.4

<sup>\*</sup>End dates of 2-week intervals are indicated. NI, no identified persons with mpox.

<sup>†</sup>Poverty levels were defined as the percentage of residents in the patient's ZIP code living below the federal poverty level according to the American Community Survey (https://www.census.gov). Neighborhoods were categorized by using the NYC DOHMH EpiQuery tool (https://a816-health.nyc.gov/hdi/epiquery) into 4 groups: low poverty, <10% of population; medium, 10%−19.9%; high, 20%−29.9%; and very high, ≥30%. ‡Total percentages include treated persons who had no prescription date (n = 22) and, therefore, could not be assigned to a 2-week interval period.