

and death, possibly associated with multidrug-resistant *Shigella* spp., a review of the national policy for the management of bloody diarrhea is urgently needed.

Acknowledgments

We thank Darrel Cecil, Temas Ikanofi, Leomeldo Latorre, and Luisa Wanma for their diagnostic support; and Anthony Gomes, Irwin Law, Carmen Aramburu, and Eigil Sorensen for their technical support.

**Alexander Rosewell,
Rosheila Dagina,
Manoj Murhekar, Berry Ropa,
Enoch Posanai, Samir Dutta,
Ian Barr, Glen Mola,
Anthony Zwi,
and C. Raina MacIntyre**

Author affiliations: World Health Organization, Port Moresby, Papua New Guinea (A. Rosewell, M. Murhekar); University of New South Wales, Sydney, New South Wales, Australia (A. Rosewell, A.

Zwi, C.R. MacIntyre); National Department of Health, Port Moresby (R. Dagina, B. Ropa, E. Posanai); Port Moresby General Hospital, Port Moresby (S. Dutta); World Health Organization Collaborating Center for Reference and Research on Influenza, Melbourne, Victoria, Australia (I. Barr); and University of Papua New Guinea, Port Moresby (G. Mola)

DOI: 10.3201/eid1704.101021

References

1. Outbreak of influenza, Madagascar, July–August 2002. *Wkly Epidemiol Rec.* 2002;77:381–4.
2. Corwin AL, Simanjuntak CH, Ingkoku-sumo G, Sukri N, Larasati RP, Subianto B, et al. Impact of epidemic influenza A-like acute respiratory illness in a remote jungle highland population in Irian Jaya, Indonesia. *Clin Infect Dis.* 1998;26:880–8. DOI: 10.1086/513917
3. Sungu M, Sanders R. Influenza virus activity in Papua New Guinea. *P N G Med J.* 1991;34:199–203.
4. Clemens J, Kotloff K, Kay B. Generic protocol to estimate the burden of *Shigella* diarrhoea and dysenteric mortality. Geneva: World Health Organization; 1999.
5. Bennish ML, Wojtyniak BJ. Mortality due to shigellosis: community and hospital data. *Rev Infect Dis.* 1991;13(Suppl 4):S245–51.
6. Kotloff KL, Winickoff JP, Ivanoff B, Clemens JD, Swerdlow DL, Sansonetti PJ, et al. Global burden of *Shigella* infections: implications for vaccine development and implementation of control strategies. *Bull World Health Organ.* 1999;77:651–66.
7. Grais RF, Conlan AJK, Ferrari MJ, Djibo A, Le Menach A, Bjørnstad ON, et al. Time is of the essence: exploring a measles outbreak response vaccination in Niamey, Niger. *J R Soc Interface.* 2008;5:67–74. DOI: 10.1098/rsif.2007.1038
8. Situation report—diarrhoeal disease outbreaks in Papua New Guinea. Port Moresby (Papua New Guinea): Surveillance Unit, National Department of Health; 2009.
9. World Health Organization. Guidelines for the control of shigellosis, including epidemics due to *Shigella dysenteriae* type 1 [2010 May 20]. <http://www.who.int/topics/cholera/publications/shigellosis/en/index.html>

Address for correspondence: Alexander Rosewell, World Health Organization, 4th Floor, AOPI Centre, PO Box 5896, Port Moresby, Papua New Guinea; email: rosewella@wpro.who.int

Vol. 16, No. 12

An online Technical Appendix was omitted from the article *Mycobacterium tuberculosis* Infection of Domesticated Asian Elephants, Thailand (T. Angkawanish, et al.). The article has been corrected online (<http://www.cdc.gov/eid/content/16/12/1949.htm>).

Get the content you want delivered to your inbox.



**Table of Contents
Podcasts
Ahead of Print Articles
Medscape CME™
Specialized Content**

Online subscription: www.cdc.gov/ncidod/eid/subscribe.htm