

the genera *Eptesicus* and *Rhinolophus* in South Korea. However, nucleotide sequencing showed the presence of prototypical Hantaan virus indicating a spillover infection or laboratory contamination (7).

Further screening is necessary to confirm *N. hispidus* as a natural reservoir host of the virus. Although the presented bat-associated sequence is obviously distinct from other hantaviruses, which suggests association with a novel natural host, a spillover infection from another, yet unrecognized host cannot be ruled out. However, detection of the virus exclusively in 1 organ (lung but not in liver, kidney, and spleen; data not shown) suggests a persistent infection that is typically observed in natural hosts of hantaviruses (8).

To date, only a few reports exist on cases of hemorrhagic fever with renal syndrome in Africa (9,10). However, underreporting must be assumed because the symptoms resemble those of many other febrile infections. Moreover, in cases of infections by non-rodent-associated hantaviruses, cross-reactivity with routinely used rodent-borne virus antigens should be limited and may hamper human serodiagnostics (1). The results suggest that bats, which are hosts of many emerging pathogens (5), may act as natural reservoirs for hantavirus. The effect of this virus on public health remains to be determined.

This study was supported by Deutsche Forschungsgemeinschaft (grant KR1293/9-1). Work in Sierra Leone was supported by the Gola Forest Project. National and local authorities in Sierra Leone, Liberia, Mali, Senegal, and Republic of Congo kindly granted research and export permits. N.D.W. is supported by an award from the National Institutes of Health Director's Pioneer Award (grant DP1-OD000370).

**Sabrina Weiss,<sup>1</sup>  
Peter T. Witkowski,<sup>1</sup>  
Brita Auste, Kathrin Nowak,  
Natalie Weber, Jakob Fahr,  
Jean-Vivien Mombouli,  
Nathan D. Wolfe,  
Jan Felix Drexler,  
Christian Drosten,  
Boris Klempa,  
Fabian H. Leendertz,  
and Detlev H. Kruger**

Author affiliations: Robert Koch-Institute, Berlin, Germany (S. Weiss, K. Nowak, F.H. Leendertz); Charité School of Medicine, Berlin (P.T. Witkowski, B. Auste, B. Klempa, D.H. Kruger); Ulm University, Ulm, Germany (N. Weber); University of Braunschweig, Braunschweig, Germany (J. Fahr); Laboratoire National de Santé Publique, Brazzaville, Republic of the Congo (J.-V. Mombouli); Stanford University Program in Human Biology, Stanford, California, USA (N.D. Wolfe); University of Bonn Medical Centre, Bonn, Germany (J.F. Drexler, C. Drosten); and Slovak Academy of Sciences, Bratislava, Slovakia (B. Klempa)

DOI: <http://dx.doi.org/10.3201/eid1801.111026>

## References

1. Krüger DH, Schönrich G, Klempa B. Human pathogenic hantaviruses and prevention of infection. *Hum Vaccin*. 2011;7:685–93. doi:10.4161/hv.7.6.15197
2. Klempa B, Fichet-Calvet E, Lecompte E, Auste B, Aniskin V, Meisel H, et al. Hantavirus in African wood mouse, Guinea. *Emerg Infect Dis*. 2006;12:838–40.
3. Klempa B, Fichet-Calvet E, Lecompte E, Auste B, Aniskin V, Meisel H, et al. Novel hantavirus sequences in shrew, Guinea. *Emerg Infect Dis*. 2007;13:520–2. doi:10.3201/eid1303.061198
4. Kang HJ, Kadjo B, Dubey S, Jacquet F, Yanagihara R. Molecular evolution of Azagny virus, a newfound hantavirus harbored by the West African pygmy shrew (*Crocodyrus obscurior*) in Cote d'Ivoire. *Virology*. 2011;8:373. doi:10.1186/1743-422X-8-373
5. Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T. Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev*. 2006;19:531–45. doi:10.1128/CMR.00017-06
6. Murphy WJ, Eizirik E, Johnson WE, Zhang YP, Ryder OA, O'Brien SJ. Molecular phylogenetics and the origins of placental mammals. *Nature*. 2001;409:614–8. doi:10.1038/35054550
7. Jung YT, Kim GR. Genomic characterization of M and S RNA segments of hantaviruses isolated from bats. *Acta Virol*. 1995;39:231–3.
8. Schönrich G, Rang A, Lutteke N, Raftery MJ, Charbonnel N, Ulrich RG. Hantavirus-induced immunity in rodent reservoirs and humans. *Immunol Rev*. 2008;225:163–89. doi:10.1111/j.1600-065X.2008.00694.x
9. Coulaud X, Chouaib E, Georges AJ, Rollin P, Gonzalez JP. First human case of haemorrhagic fever with renal syndrome in the Central African Republic. *Trans R Soc Trop Med Hyg*. 1987;81:686. doi:10.1016/0035-9203(87)90455-X
10. Klempa B, Koivogui L, Sylla O, Koulemou K, Auste B, Kruger DH, et al. Serological evidence of human hantavirus infections in Guinea, West Africa. *J Infect Dis*. 2010;201:1031–4. doi:10.1086/651169

Address for correspondence: Detlev H. Kruger, Institute of Medical Virology, Helmut-Ruska-Haus, Charité Medical School, Charitéplatz 1, D-10117 Berlin, Germany; email: [detlev.kruger@charite.de](mailto:detlev.kruger@charite.de)

## Outbreak of Porcine Epidemic Diarrhea in Suckling Piglets, China

**To the Editor:** Beginning in October 2010, porcine epidemic diarrhea (PED), caused by a coronaviral infection affecting pigs, emerged in China in an outbreak characterized by high mortality rates among suckling piglets. The outbreak overwhelmed >10 provinces in southern China, and >1,000,000 piglets died. This outbreak was distinguished by ≈100% illness among piglets after birth (predominantly within 7 days and

<sup>1</sup>These authors contributed equally to this article.

sometimes within only a few hours) and death rates of 80%–100% (online Technical Appendix Table 1, [wwwnc.cdc.gov/EID/pdfs/11-1259-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-1259-Techapp.pdf)). Few sows or boars showed any clinical signs during the outbreak, which is not consistent with a recent report from Thailand (1). In that outbreak during late 2007, pigs of all ages were affected, exhibiting different degrees of diarrhea and no appetite. We characterized the genetic variation of the PED virus (PEDV) that caused a large-scale outbreak in China during 2010–2011 and compared it with viruses in other outbreaks. We also report a possible novel transmission pathway for PEDV.

A total of 177 samples (intestine, stool, and maternal milk) were collected from pigs from different farms who had diarrhea; 100% of farms had  $\geq 1$  porcine sample positive for PEDV. A total of 125/177 porcine samples were confirmed as positive for PEDV by reverse transcription PCR using primers as described (2). PEDV was detected in 105 (82.0%) of 128 fecal samples and 20 (40.8%) of 49 sow milk samples. Piglets infected with PEDV showed mild hemorrhage, undigested curdled milk in the stomach, and thin-walled intestines with severe mucosal atrophy and foamy fluid (data not shown).

The spike (S) gene of the family *Coronaviridae* has a high degree of variation and can induce neutralizing antibody (3). Reverse transcription PCR products of the 651-bp partial S gene of PEDV and the deduced amino acid sequences were aligned by using ClustalW ([www.genome.jp/tools/clustalw](http://www.genome.jp/tools/clustalw)), and a neighbor-joining tree with 1,000 bootstraps was constructed. Sequences of the S genes from this outbreak were 99.1%–100.0% homologous and had 88.7%–98.9% nt identity with all reference strains (online Technical Appendix Table 2), 98.5%–98.9% with Thailand strains, and 94.5%–95.1% with vaccine strain CV777. The partial S

gene deduced amino acid sequences were compared and also showed a high degree of homology (98.0%–100.0%); they had 85.3%–98.7% identity with all reference strains listed in online Technical Appendix Table 2, 98.0%–98.7% with Thailand strains, and 93.3%–94.7% with vaccine strain CV777 (data not shown).

Phylogenetic analysis indicated that the PEDV in the China outbreak was different from foreign and other domestic strains on the basis of the reported partial S gene sequences. All new strains were clustered in the same branch, close to the cluster of Thailand strains, and far from the cluster of vaccine strain CV777 (Figure).

In the China outbreak, PEDV caused severe diarrheal disease in piglets; heavy economic losses in many provinces resulted, despite use of commercial vaccines (inactivated

transmissible gastroenteritis [TGEV H] and porcine epidemic diarrhea [CV777]). To determine why the vaccines showed poor efficacy, we investigated evolution of the virus. Comparison of amino acid sequences from isolates from the outbreak and from the CV777 vaccine strain showed 9 amino acid mutations of fragments containing major hydrophilic regions: 16 (L→H), 18 (S→G), 22 (V→I), 44 (T→S), 89 (G→S), 100 (A→E), 107 (L→F), 130 (I→V) and 160 (I→F) (online Technical Appendix Figure, panel A). Three of these 9 mutations were at positions 16, 18, and 22 in the isolates from China; they influenced the hydrophobicity of the S protein as compared with that for CV777 (online Technical Appendix Figure, panel B).

Phylogenetic analysis showed that strain CV777 did not cluster with current common strains and showed




Figure. Phylogenetic tree constructed by using the neighbor-joining method based on the 9 porcine epidemic diarrhea virus (PEDV) sequences identified in a study of porcine epidemic diarrhea in China. Partially amplified spike genes of the PEDV isolates plus 18 PEDV sequences downloaded from GenBank were compared. Sequences included in each cluster are listed in online Technical Appendix Table 3 ([wwwnc.cdc.gov/EID/pdfs/11-1259-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-1259-Techapp.pdf)). Strains from Thailand and China and the CV777 vaccine strain are indicated. Scale bar indicates nucleotide substitutions per site.

considerable genetic distance from them. Isolates in the outbreak in China had only a minor nucleotide sequence variation from the Thailand isolates, indicating that the virus has a high genetic relatedness to the Southeast Asia strain. However, previous studies showed that isolates from Europe, South Korea, and China were serologically identical to the prototype CV777 strain (1,4).

To our knowledge, fecal–oral transmission is probably the main or only route of PEDV transmission (5–7). In our study, if a fecal sample from a sick piglet was found to be positive for PEDV, we also collected and studied milk from its mother. These results showed that PEDV was present in sow milk (online Technical Appendix Table 3), but the detection rate was lower for these samples (40.8%) than for the fecal samples (82.0%).

On the basis of these results, we hypothesize that sow milk could represent a possible (and potentially major) route for the vertical transmission of PEDV from sow to suckling piglet. This hypothesis could be indirectly verified by our field observation that piglet death rates decreased as a result of fostering (data not shown). Our findings show that PEDV was identified not only in fecal samples from sick piglets, as expected, but also in the milk of the sow, which suggests vertical transmission of the virus.

#### Acknowledgments

We thank the technicians from the pig farms for assistance in sample collection and Bin Wu for assistance in figure preparation. We also thank the Guangdong Province Pig Industry Innovation Projects for their support.

**Rui-Qin Sun, Ru-Jian Cai,  
Ya-Qiang Chen,  
Peng-Shuai Liang,  
De-Kun Chen,  
and Chang-Xu Song**

Author affiliations: Guangdong Academy of Agricultural Sciences Veterinary Medical Institute, Guangzhou, China (R.-Q. Sun, R.-J. Cai, C.-X. Song); Northwest A & F University, Xi'an, Shanxi, China (R.-Q. Sun, D.-K. Chen); Inner Mongolia Agriculture University, Huhhot, Inner Mongolia, China (Y.-Q. Chen); and Henan Agriculture University, Zhengzhou, Henan, China (P.-S. Liang)

DOI: <http://dx.doi.org/10.3201/eid1801.111259>

#### References

1. Puranaveja S, Poolperm P, Lertwatcharasarakul P, Kesdaengsakonwut S, Boonsoongnern A, Urairong K, et al. Chinese-like strain of porcine epidemic diarrhea virus, Thailand. *Emerg Infect Dis*. 2009;15:1112–5. doi:10.3201/eid1507.081256
2. Kim SY, Song DS, Park BK. Differential detection of transmissible gastroenteritis virus and porcine epidemic diarrhea virus by duplex RT-PCR. *J Vet Diagn Invest*. 2001;13:516–20. doi:10.1177/104063870101300611
3. Jiménez G, Correa I, Melgosa MP, Buldido MJ, Enjuanes L. Critical epitopes in transmissible gastroenteritis virus neutralization. *J Virol*. 1986;60:131–9.
4. Pospischil A, Hess RG, Bachmann PA. Light microscopy and ultrahistology of intestinal changes in pigs infected with epizootic virus diarrhoea (EVD): comparison with transmissible gastroenteritis (TGE) virus and porcine rotavirus infections. *Zentralbl Veterinarmed B*. 1981;28:564–77. doi:10.1111/j.1439-0450.1981.tb01774.x
5. Riley S. Large-scale spatial-transmission models of infectious disease. *Science*. 2007;316:1298–301. doi:10.1126/science.1134695
6. Utiger A, Tobler K, Bridgen A, Ackermann M. Identification of the membrane protein of porcine epidemic diarrhea virus. *Virus Genes*. 1995;10:137–48. doi:10.1007/BF01702594
7. Turgeon DC, Morin M, Jollette J, Higgins R, Marsolais G, DiFranco E. Coronavirus-like particles associated with diarrhea in baby pigs in Quebec. *Can Vet J*. 1980;21:100–xxiii.

Address for correspondence: Chang-Xu Song, Veterinary Medicine Institute, Guangdong Academy of Agricultural Sciences, Guangzhou, People's Republic of China; email: [cxsong2004@163.com](mailto:cxsong2004@163.com)

## *Bartonella quintana* Transmission from Mite to Family with High Socioeconomic Status

**To the Editor:** Urban trench fever caused by *Bartonella quintana* has been reported in persons who abuse alcohol and in homeless persons in large cities worldwide. Symptoms vary from asymptomatic intermittent bacteremia to serious complications (1). *Pediculus humanus* mites, the known vector of the infection, are not always identified, which raises the possibility that other vectors might also be involved (2). We report on an outbreak of *B. quintana* infection among a young family of high socioeconomic status and their visiting relatives.

The family resides in a regional city (population 104,000) in northern Czech Republic in an old, renovated apartment located on the top floor, just under the roof. In the summer of 2007, hundreds of ectoparasitic mites migrated from a whole in the roof and settled on the inner side of a permanently open window before infesting family members. Two weeks later (day 1 of symptom onset), a papular rash and pruritic vesicular lesions were noted by the parents on the body and legs of their 2 children, a 1-year-old girl and a 3-year-old boy. On day 3, the girl's body temperature rose to 38.0°C, and the boy's temperature rose to 39.5°C. The rash resolved in ≈10 days in both children. Vesicular lesions on the girl's buccal mucosal membrane resolved in 5 days. Excoriated areas resulting from spontaneous rupture of lesions or scratching were still visible on day 14.

On day 4, a fever (temperature, 38.5°C) and intense tibialgia, which persisted for 5 days, developed in the 33-year-old father of the infected children. On day 5, a vesicular rash, which resolved in 10 days, developed in the 33-year-old mother. The children's