


Seroincidence of Enteric Fever, Juba, South Sudan

Appendix



Appendix Figure. Age-dependent seroprevalence curves using negative control cutoff. Each point is an individual classified into positive or negative based on a cutoff derived from a North American control population for hemolysin E immunoglobulin A (HlyE IgA) (left) and hemolysin E immunoglobulin G (HlyE IgG) (right). The purple line is the age-dependent seroprevalence curve fit using generalized additive models.

Additional References

16. Teunis PFM, van Eijkeren JCH. Estimation of seroconversion rates for infectious diseases: effects of age and noise. *Stat Med*. 2020;39:2799–814. [PubMed](https://doi.org/10.1002/sim.8578) <https://doi.org/10.1002/sim.8578>
17. Leonard MM, Camhi S, Huedo-Medina TB, Fasano A. Celiac Disease Genomic, Environmental, Microbiome, and Metabolomic (CDGEMM) study design: approach to the future of personalized prevention of celiac disease. *Nutrients*. 2015;7:9325–36. [PubMed](https://doi.org/10.3390/nu7115470) <https://doi.org/10.3390/nu7115470>

18. Wood SN. Generalized additive models: an introduction with R. London: Chapman and Hall/CRC; 2006.
19. Marra G, Wood SN. Coverage properties of confidence intervals for generalized additive model components. *Scand J Stat*. 2012;39:53–74. <https://doi.org/10.1111/j.1467-9469.2011.00760.x>
20. Meiring JE, Shakya M, Khanam F, Voysey M, Phillips MT, Tonks S, et al.; STRATAA Study Group. Burden of enteric fever at three urban sites in Africa and Asia: a multicentre population-based study. *Lancet Glob Health*. 2021;9:e1688–96. [PubMed https://doi.org/10.1016/S2214-109X\(21\)00370-3](https://doi.org/10.1016/S2214-109X(21)00370-3)
21. Crump JA, Luby SP, Mintz ED. The global burden of typhoid fever. *Bull World Health Organ*. 2004;82:346–53. [PubMed](#)