

Dynamics of SARS-CoV-2 Antibody Response to CoronaVac followed by Booster Dose of BNT162b2 Vaccine

Marcela Helena Gambim Fonseca, Ana Carolina Matias Dinelly Pinto, Maria Francilene Souza Silva, Amanda Campelo Lima de Melo, Germana Silva Vasconcelos, Eduardo Ruback dos Santos, Fernanda Montenegro de Carvalho Araújo, Luiz Odorico Monteiro de Andrade

We evaluated the longitudinal dynamics of antibody response to the SARS-CoV-2 vaccine CoronaVac and the effect of a booster dose of BNT162b2 vaccine. We found a robust antibody response after the second dose of CoronaVac that wanes over time. The response was recovered by BNT162b2, which boosted anti-spike antibody titers.

Long-term protection against SARS-CoV-2 requires the persistence of vaccine antibodies above protective thresholds, the maintenance of immune memory cells capable of reactivation after subsequent viral exposure, or both (1). A decay of circulating SARS-CoV-2 antibodies over time in persons who received CoronaVac (Sinovac, <http://www.sinovac.com>) have been reported, suggesting the necessity of a third shot of vaccine (2). In Brazil, the third dose has been administered, preferably, with the BNT162b2 vaccine (Pfizer-BioNTech, <https://www.pfizer.com>) (3,4). Limited information is available about antibody dynamics after CoronaVac vaccine and the recent supplementing with the BNT162b2 booster. Therefore, we evaluated the longitudinal dynamics of the antibody response to CoronaVac up to 230 days after the second dose in a cohort of healthcare workers (HCWs) and evaluated the effect of a booster dose of BNT162b2 on antibody levels. The study was approved by the Ethics Committee of the Hospital Geral Dr. César Cals (Fortaleza, Brazil; approval no. CAAE 39691420.7.0000.5049). We obtained informed consent from all participants.

The Study

We included in this study 99 HCWs of both sexes, ≥ 18 years of age, who had received 2 doses of the CoronaVac vaccine, with an interval of 28 days between doses, and then a booster shot of BNT162b2 vaccine

8 months after the second CoronaVac dose. Blood collections and serologic tests were performed at Fundação Oswaldo Cruz (Fiocruz; Ceará, Brazil) and analyzed at 7 different timepoints: before vaccination (P1); 28 days after the first CoronaVac dose (P2); 30 (P3) 90 (P4), 180 (P5), and 230 (P6) days after the second CoronaVac dose; and 15 days after the BNT162b2 dose (P7). We monitored the HCWs for SARS-CoV-2 infection by PCR over time.

We tested all serum samples for IgG against nucleocapsid (N) and spike (S) proteins of SARS-CoV-2 by using chemiluminescent microparticle immunoassays on the ARCHITECT i2000SR equipment (Abbott, <https://www.abbott.com>). The cutoff value was 50 AU/mL for S antibodies and 1.4 index (S/CO) for N antibodies.

We used GraphPad Prism version 9 (<https://www.graphpad.com>) for statistical analyses. We describe data as median and interquartile range (IQR) or percentage. In group comparisons, we used χ^2 test to analyze the seropositivity data and Kruskal-Wallis test with subsequent Dunn's test to analyze the IgG values. We considered differences with $p < 0.05$ to be statistically significant.

The cohort was 70.71% women and 29.29% men. Average age was 32.31 years (95% CI 30.3–34.3 years). The age distribution of persons was as follows: 18–30 years, 42 (42.4%); 31–45 years, 51 (51.5%); and >45 years, 6 (6.1%). Median age for each age group was as follows: 18–30 years, 23.0 years (IQR 20–27.3 years); 31–45 years, 36 years (IQR 32–39 years), >45 years, 56.50 years (IQR 52–63.3 years).

Although all HCWs completed the vaccination schedule, some HCWs were unable to give a blood sample in subsequent phases of the study. Therefore, serum samples were obtained from 99 participants in P1 and P2, 95 in P3, 94 in P4, 89 in P5, 84 in P6, and 74 in P7.

We evaluated the seropositivity and IgG levels for S and N proteins at the different timepoints

Author affiliations: Fundação Oswaldo Cruz, Eusébio, Brazil

DOI: <https://doi.org/10.3201/eid2806.220061>

(Figure 1). At baseline (P1), S IgG were detectable in 25.3% of HCWs, increasing to 84.9% in P2 and reaching 100% in P3. We then observed a decline in seropositivity was observed to 98.9% in P4, 94.4% in P5, and 89.3% in P6. In P7, seropositivity had recovered to 100%. N IgG was detectable in 8.1% of HCWs in P1, 19.2% in P2, and 52.6% in P3, then reduced to 29.8% in P4, 13.5% in P5, 10.7% in P6, and 12.2% in P7 (Figure 1, panel A). The seroconversion rate was 79.7% for S antibodies and 11.9% for N antibodies after the first CoronaVac dose, increasing to 100% for S antibodies and 43.6% N antibodies after the second dose.

After the first CoronaVac dose (P2), S IgG levels were significantly elevated compared with baseline values ($p < 0.0001$) (Figure 1, panel B; Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/28/6/22-0061-App1.pdf>). Those S IgG levels increased significantly after the second dose (P3) ($p < 0.0001$). However, antibodies levels waned over time. The third BNT162b2 dose again significantly increased S IgG levels ($p < 0.0001$). We observed a similar change in N IgG (Figure 1, panel C; Appendix Table 1). Median values of N IgG were significantly higher after the second CoronaVac dose ($p < 0.0001$) and declined significantly after vaccination ($p = 0.0002$). In contrast, the third dose with BNT162b2 did not increase N IgG levels.

We evaluated the antibody response to the vaccine in relation to a previous SARS-CoV-2 infection. Twenty-five volunteers were seropositive in P1 and were included in the COVID-19-positive group. Eight volunteers had a positive PCR result during the study (4 in P3, 4 in P4); they were moved to the COVID-19-positive group.

The HCWs who had COVID-19 maintained the anti-S seropositivity at 100% over time. In relation to HCWs who did not have COVID-19, 79.7% of persons became seropositive to S protein in P2. The seropositivity increased to 100% in P3 but decreased in the next timepoints, recovering in P7 (Figure 2, panel A). The differences in seropositivity between the groups were statistically significant in P2 ($p = 0.0145$). Anti-N seropositivity was 6 (95% CI 2.7–15.1) times higher in the COVID-19-positive group in P2 compared with the COVID-19-negative group. This difference was reduced in P3, increasing in P4 and stabilizing in P5, reaching levels of 5 (95% CI 1.4–18.4) times higher in the COVID-19-positive group.

In the antibody levels analysis, antibody titers for S protein were higher in the COVID-19-positive group than for the COVID-19-negative group at all timepoints except P3 and P7 (Figure 2, panel B; Appendix Table 2). N IgG levels of COVID-19-positive persons were statistically higher than for COVID-19-negative persons at all timepoints (Figure 2, panel C; Appendix Table 2).

Conclusions

We found an antibody response to N and S protein after 2 doses of CoronaVac vaccine. However, the antibodies declined over time. After immunization, the decline of antibodies is expected because not all vaccine-induced plasmablasts commit or are maintained as long-lived memory plasma cells (5). Thus, the success of vaccines depends on the generation and maintenance of immunologic memory (6). Administration of BNT162b2 as the third vaccine dose boosted S IgG but not N IgG. The substantial increase of S IgG after




Figure 1. Antibody response over time in a cohort of healthcare workers vaccinated with 2 doses of CoronaVac vaccine (<https://www.sinovac.com>) followed by a BNT162b2 vaccine (Pfizer-BioNTech, <https://www.pfizer.com>) booster dose. A) S and N IgG seropositivity. B) S IgG levels. C) N IgG levels. Antibody responses were evaluated before vaccination (timepoint P1); 28 days after the first dose of CoronaVac vaccine (P2); 30 (P3) 90 (P4), 180 (P5), and 230 (P6) days after the second dose of CoronaVac vaccine; and 15 days after the booster dose with BNT162b2 vaccine (P7). For panels B and C, black lines indicate median levels values and error bars interquartile ranges; horizontal dotted lines indicate cutoff values. Statistical analysis performed using the Kruskal–Wallis test with subsequent Dunn’s multiple testing correction. N, nucleocapsid protein; S, spike protein; S/CO, signal-to-cutoff ratio.




Figure 2. Comparison of antibody response over time among healthcare workers vaccinated with 2 doses of CoronaVac vaccine (https://www.sinovac.com) followed by a BNT162b2 vaccine (Pfizer-BioNTech, https://www.pfizer.com) booster dose, by COVID-19 positivity status. A) S and N IgG seropositivity. B) S IgG levels. C) N IgG levels. Antibody responses were evaluated before vaccination (timepoint P1); 28 days after the first dose of CoronaVac vaccine (P2); 30 (P3) 90 (P4), 180 (P5), and 230 (P6) days after the second dose of CoronaVac vaccine; and 15 days after the booster dose with BNT162b2 vaccine (P7). For panels B and C, black lines indicate median levels values and error bars interquartile ranges; the horizontal dotted line indicates the cutoff value of the assays. Numbers below p values indicate numbers of COVID-19–positive and COVID-19–negative persons in each timepoint. Statistical analysis performed using the Kruskal–Wallis test with subsequent Dunn’s multiple testing correction. N, nucleocapsid protein; S, spike protein; S/CO, signal-to-cutoff ratio; (–), COVID-19 negative; (+), COVID-19 positive.

the booster dose suggests that CoronaVac vaccine induced immune memory. The third BNT162b2 dose did not increase the N IgG because mRNA vaccines do not induce a response to the N protein (7,8).

Previously infected participants had a significantly higher antibody level than previously uninfected participants in almost all phases of the study. In addition, we found that those without previous infection showed a faster waning of antibodies over time, a result also reported in previous studies (9). The antibody-making B cells multiply after each exposure, whether attributable to the infection or vaccination; therefore, antibody levels in the previously infected HCWs can reflect the sum of the antibodies produced after infection and vaccine (10).

In summary, a booster dose of BNT162b2 vaccine in HCWs administered 8 months after the second dose with CoronaVac vaccine recalled a specific immune response to SARS-CoV-2. That response had declined substantially 230 days after the second dose of CoronaVac vaccine, resulting in an increase of S IgG after BNT162b2 vaccination and indicating that the 2-dose CoronaVac vaccine schedule generates immune memory.

Acknowledgments

We thank the healthcare workers of COVID-19 Diagnosis Support Unit of Fiocruz, Ceará, Brazil, for participating in this study. We also thank the Paulo Goberlânio de Barros Silva for the statistical analysis.

The project is funded by Fiocruz and Ministério da Saúde, Brazil.

About the Author

Dr. Gambim Fonseca is a researcher at and coordinator of the Serology Laboratory of the COVID-19 Diagnosis Support Unit of Fiocruz Ceará, Brazil. Her primary research interests include antibodies and coronavirus disease.

References

- Siegrist C-A. Vaccine immunology. 2018 Jan 1 [cited 2022 Mar 22]. https://www.who.int/immunization/documents/Elsevier_Vaccine_immunology.pdf
- Sauré D, O’Ryan M, Torres JP, Zuniga M, Santelices E, Basso LJ. Dynamic IgG seropositivity after rollout of CoronaVac and BNT162b2 COVID-19 vaccines in Chile: a sentinel surveillance study. *Lancet Infect Dis.* 2022;22:56–63. [https://doi.org/10.1016/S1473-3099\(21\)00479-5](https://doi.org/10.1016/S1473-3099(21)00479-5)
- Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al.; C4591001 Clinical Trial Group. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med.* 2020;383:2603–15. <https://doi.org/10.1056/NEJMoa2034577>
- Kanno AI, Barbosa MMF, Moraes L, Leite LCC. SARS-CoV-2 vaccine development and how Brazil is contributing. *Genet Mol Biol.* 2021;44(Suppl 1):e20200320. <https://doi.org/10.1590/1678-4685-gmb-2020-0320>
- Naaber P, Tserel L, Kangro K, Sepp E, Jürjenson V, Adamson A, et al. Dynamics of antibody response to BNT162b2 vaccine after six months: a longitudinal prospective study. *Lancet Reg Health Eur.* 2021;10:100208. <https://doi.org/10.1016/j.lanepe.2021.100208>
- Palm AE, Henry C. Remembrance of things past: long-term B cell memory after infection and vaccination. *Front Immunol.* 2019;10:1787. <https://doi.org/10.3389/fimmu.2019.01787>
- Mueller T. Antibodies against severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) in individuals with and without COVID-19 vaccination: a method comparison of two different commercially available serological assays from the same manufacturer. *Clin Chim*


- Acta. 2021;518:9–16. <https://doi.org/10.1016/j.cca.2021.03.007>
8. Keskin AU, Bolukcu S, Ciragil P, Topkaya AE. SARS-CoV-2 specific antibody responses after third CoronaVac or BNT162b2 vaccine following two-dose CoronaVac vaccine regimen. *J Med Virol.* 2022;94:39–41. <https://doi.org/10.1002/jmv.27350>
9. Vicenti I, Basso M, Gatti F, Scaggiante R, Boccutto A, Zago D, et al. Faster decay of neutralizing antibodies in never infected than previously infected healthcare workers three months after the second BNT162b2 mRNA COVID-19 vaccine dose. *Int J Infect Dis.* 2021;112:40–4. <https://doi.org/10.1016/j.ijid.2021.08.052>
10. Ali H, Alahmad B, Al-Shammari AA, Alterki A, Hammad M, Cherian P, et al. Previous COVID-19 infection and antibody levels after vaccination. *Front Public Health.* 2021;9:778243. <https://doi.org/10.3389/fpubh.2021.778243>

Address for correspondence: Marcela Helena Gambim Fonseca, Fundação Oswaldo Cruz, São Jose s/n, Eusebio, Ceara, Brazil; email: marcela.gambim@fiocruz.br

April 2022

Zoonotic Infections

- Citywide Integrated *Aedes aegypti* Mosquito Surveillance as Early Warning System for Arbovirus Transmission, Brazil
- *Shewanella* spp. Bloodstream Infections in Queensland, Australia
- Increasing Antimicrobial Resistance in World Health Organization Eastern Mediterranean Region, 2017–2019
- Phylogenetic Analysis of Spread of Hepatitis C Virus Identified during HIV Outbreak Investigation, Unnao, India
- SARS-CoV-2 IgG Seroprevalence among Blood Donors as a Monitor of the COVID-19 Epidemic, Brazil
- Diminishing Immune Responses against Variants of Concern in Dialysis Patients 4 Months after SARS-CoV-2 mRNA Vaccination
- Genomic Epidemiology of Early SARS-CoV-2 Transmission Dynamics, Gujarat, India
- Reassessing Reported Deaths and Estimated Infection Attack Rate during the First 6 Months of the COVID-19 Epidemic, Delhi, India
- Mapping the Risk for West Nile Virus Transmission, Africa
- Isolation of Heartland Virus from Lone Star Ticks, Georgia, USA, 2019
- *Bordetella hinzii* Pneumonia in Patient with SARS-CoV-2 Infection



- Increased Attack Rates and Decreased Incubation Periods in Raccoons with Chronic Wasting Disease Passed through Meadow Voles
- Fatal Human Alphaherpesvirus 1 Infection in Free-Ranging Black-Tufted Marmosets in Anthropized Environments, Brazil, 2012–2019
- Molecular Surveillance for Imported Antimicrobial Resistant *Plasmodium falciparum*, Ontario, Canada
- Unique Clinical, Immune, and Genetic Signature in Patients with Borrelial Meningoradiculoneuritis
- Durability of Antibody Response and Frequency of SARS-CoV-2 Infection 6 Months after COVID-19 Vaccination in Healthcare Workers

- SARS-CoV-2 Outbreak among Malayan Tigers and Humans, Tennessee, USA, 2020
- Zika Virus after the Public Health Emergency of International Concern Period, Brazil
- Vehicle Windshield Wiper Fluid as Potential Source of Sporadic Legionnaires' Disease in Commercial Truck Drivers
- Coccidioidomycosis Cases at a Regional Referral Center, West Texas, USA, 2013–2019
- In Vitro Confirmation of Artemisinin Resistance in *Plasmodium falciparum* from Patient Isolates, Southern Rwanda, 2019
- *Rigidoporus corticola* Colonization and Invasive Fungal Disease in Immunocompromised Patients, United States
- Zoonotic Pathogens in Wildlife Traded in Markets for Human Consumption, Laos
- Infectious Toscana Virus in Seminal Fluid of Young Man Returning from Elba Island, Italy
- Multisystem Inflammatory Syndrome in Adult after First Dose of mRNA Vaccine
- Hantavirus Pulmonary Syndrome in a COVID-19 Patient, Argentina, 2020

**EMERGING
INFECTIOUS DISEASES**

To revisit the April 2022 issue, go to:
<https://wwwnc.cdc.gov/eid/articles/issue/28/4/table-of-contents>