

RESEARCH

18. National Committee for Clinical Laboratory Standards. Method for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Approved standard. NCCLS document M7-A3. Villanova (PA): The Committee; 1993.
19. Hooper DC. New uses for new and old quinolones and the challenge of resistance. *Clin Infect Dis* 2000;30:243-54.
20. Tee W, Mijch A, Wright E, Yung A. Emergence of multidrug resistance in *Campylobacter jejuni* isolates from three patients infected with human immunodeficiency virus. *Clin Infect Dis* 1995;21:634-8.
21. Burnens AP, Heitz M, Brodard I, Nicolet J. Sequential development of resistance to fluoroquinolones and erythromycin in an isolate of *Campylobacter jejuni*. *Zentralbl Bakteriol* 1996;283:314-21.
22. Lu PL, Hsueh PR, Hung CC, Chang SC, Luh KT, Lee CY. Bacteremia due to campylobacter species: high rate of resistance to macrolide and quinolone antibiotics. *J Formos Med Assoc* 2000;99:612-7.
23. Steinbigel NH. Erythromycin, lincomycin, and clindamycin. In: Mandell GL, Douglas RG, Bennett JE, editors. Principles and practice of infectious diseases, 3rd ed. New York: Churchill Livingstone; 1990. p. 309.
24. Van Duong D, Binns CW, Van Le T. Availability of antibiotics as over-the-counter drugs in pharmacies: a threat to public health in Vietnam. *Trop Med Int Health* 1997;2:1133-9.
25. Thamlikitkul V. Antibiotic dispensing by drug store personnel in Bangkok, Thailand. *J Antimicrob Chemother* 1988;21:125-31.
26. Visitsunthorn N, Komolpis P. Antimicrobial therapy in *Plesiomonas shigelloides*-associated diarrhea in Thai children. *Southeast Asian J Trop Med Public Health* 1995;26:86-90.

Address for correspondence: Daniel W. Isenbarger, 7543 Spring Lake Dr., #D-1, Bethesda, MD 20817, USA; fax: 202-782-3813; e-mail: isenbarg-erdw@hotmail.com

Erratum Vol. 8, No. 1

In the article, "A Large Outbreak of Legionnaires' Disease at a Flower Show, the Netherlands," by Jeroen W. Den Boer et al., errors appear in two figures and their legends.

In Figure 3, part 3a should be identified as IgM and part 3b as IgG. The correct legends are as follows:

Figure 3a. Smoothed mean geometric immunoglobulin (Ig) M antibody titers to *Legionella pneumophila* of nearest 35 exhibitors in Halls 3 and 4 per 63 cm² of exhibition area; confirmed and probable cases among exhibitors in halls 3 and 4.

● = confirmed case in exhibitor; ○ = probable case in exhibitor;
Bu = bubblemat; W = whirlpool spa.

Figure 3b. Smoothed mean geometric IgG antibody titers to *L. pneumophila* of 35 exhibitors nearest to whirlpool in halls 3 and 4 per 63 cm² of exhibition area; exhibitors ill with confirmed and probable cases in halls 3 and 4.

● = confirmed case in exhibitor; ○ = probable case in exhibitor;
Bu = bubblemat; W = whirlpool spa.

In Figure 4, parts b and c were inadvertently omitted. The complete figure appears below. The correct legends are as follows:

Figure 4. Exhibition hall, West Frisian Flower Show, Bovenkarspel, the Netherlands, 1999.

4a. Circles indicate locations in water-supply system where water samples were taken. PE = polyethylene.

4b. Assessment of risk for *Legionella pneumophila* infection, by distance from water-using devices.

4c. Water samples taken and culture status, by distance from water-using devices.

We regret any confusion these errors may have caused. The article is corrected online, available at <http://www.cdc.gov/ncidod/eid/vol8no1/01-0176.htm>

