

# Sexual Health and Sexually Transmitted Infections in the North American Arctic

Dionne Gesink Law,\* Elizabeth Rink,† Gert Mulvad,‡ and Anders Koch§

Our objective was to describe the basic epidemiology of sexually transmitted infections for Arctic and sub-Arctic regions of North America. We summarized published and unpublished rates of chlamydial infection and gonorrhea reported from 2003 through 2006 for Alaska, Canada, and Greenland. In 2006, Alaska reported high rates of chlamydial infection (715 cases/100,000 population) compared with the United States as a whole; northern Canada reported high rates of chlamydial infection (1,693 cases/100,000) and gonorrhea (247 cases/100,000) compared with southern Canada; and Greenland consistently reported the highest rates of chlamydial infection (5,543 cases/100,000) and gonorrhea (1,738 cases/100,000) in the Arctic. Rates were high for both men and women, although the highest incidence of infection was predominantly reported for young women in their early twenties. We propose that community-based participatory research is an appropriate approach to improve sexual health in Arctic communities.

Four million people live in the Arctic (1), yet little is known about sexual health and sexually transmitted infections (STIs) in the circumpolar North. Arctic communities in North America comprise a large proportion of Native American, First Nation, Metis, Inuit, and other aboriginal peoples living in harsh climates, diverse landscapes, and a variety of community structures including urban, metropolitan, reserves or reservations, towns, villages, settlements, and remote fly-in communities. Access to healthcare varies by community and country and patient concerns about the preservation of confidentiality remain a barrier to accessing healthcare.

\*University of Toronto, Toronto, Ontario, Canada; †Montana State University, Bozeman, Montana, USA; ‡Centre for Primary Health Care, Nuuk, Greenland; and §Statens Serum Institut, Copenhagen, Denmark

STI intervention and prevention strategies have been developed primarily for urban and suburban environments (2,3), the rural South (4–7), Latino communities (8), and developing countries, primarily in Africa (9,10). Cultural differences alone will affect their generalizability to communities in the Arctic. This is further emphasized by Bjerregaard et al. (11) who stated: “Intervention models developed under quite different circumstances cannot be expected to work in Greenland and intervention studies are highly needed.” However, combining the global knowledge gained from previous interventions involving other populations with the local knowledge and infrastructure of Arctic communities is important to develop innovative, culturally appropriate, and sustainable STI intervention strategies.

Our objective was to describe STI trends in the circumpolar Arctic, focusing on the North American continent (United States, Canada, and Greenland). We also propose a community-based participatory research approach to conducting research and planning interventions involving Arctic communities.

## Methods

Data on chlamydial infection and gonorrhea in the United States, Canada, and Greenland were collected from a variety of sources. Rates for the United States and Alaska were obtained from federal (12) and state (13) reports. Rates for Canada were obtained in collaboration with the Public Health Agency of Canada STI Surveillance and Epidemiology Section (Public Health Agency of Canada, unpub. data; see also [14] for published summaries). Data reported for Canada’s northern territories (Yukon, Northwest Territories, and Nunavut) were combined into 1 statistic and compared to data reported for Canada’s southern provinces, which were also combined into 1 statistic. Data for Greenland were obtained from the Office of the Chief

Medical Officer in Greenland (15,16) and compared with data reported for Denmark by the Statens Serum Institut ([www.ssi.dk](http://www.ssi.dk)). In situations where STI rates were not already available (primarily Greenland and Denmark), rates were calculated by dividing the number of cases by the total population and multiplying by 100,000. Population estimates were obtained from the US Census Bureau, Statistics Canada, Statistics Greenland, and Statistics Denmark (StatBank).

Chlamydial infection and gonorrhea rates reported for the year 2006 were standardized by age and sex to the year 2000 US population so that rates could be compared across countries after correcting for age and sex differences between the different populations. Rates were standardized to the year 2000 US population by using age- and sex-stratified counts available from the US Census Bureau ([www.census.gov](http://www.census.gov)). Additionally, chlamydial infection and gonorrhea rates were stratified by the basic demographic characteristics of age, sex, and race (when available) to gain insights into target populations for community interventions. Rates by race were only available for Alaska.

## Results

As expected, chlamydial infection was the most highly reported STI for the United States, Canada, and Greenland (Table 1). Compared with other states in the United States, Alaska reported the highest rates of chlamydial infection in 2003 and second highest in 2004 and 2005 (12). Canada's northern territories consistently reported the highest rates of chlamydial infection in Canada, which is consistent with the 1987–1994 rates measured by Orr and Brown (17) for the Keewatin District of the Canadian Central Arctic. Greenland reported chlamydial infection rates higher than Denmark and higher than any other country in the North American Arctic.

Co-infection with chlamydial infection and gonorrhea is common so we expected gonorrhea rates to be high for the Arctic regions. However, Alaska reported some of the lowest gonorrhea rates in the United States (12). As expected, however, the Canadian Northern Territories reported higher gonorrhea rates than their southern counterparts, and again, Greenland reported gonorrhea rates higher than those in Denmark and in any other country in the North American Arctic (Table 1).

Chlamydial infection rates reported for women were much higher than rates reported for men in Alaska (Table 2), Canada (Table 3), and Greenland and Denmark (Table 4). Compared to gonorrhea rates reported for men, however, gonorrhea rates were higher for women  $\leq 30$  years of age in Alaska (Table 2),  $\leq 20$  years of age in Canada (until 2006, when rates remained higher for women  $\leq 24$  years of age; Table 3), and  $\leq 20$  years of age or  $\leq 30$  years of age for women in Greenland (Table 4). Gonorrhea rates reported for men in Denmark were consistently higher than rates reported for women (Table 4). Reported rates of chlamydial infection and gonorrhea were consistently high for both men and women 15–30 years of age, particularly for those 20–24 years of age, regardless of country.

## Discussion

Chlamydial infection rates were higher for Arctic and sub-Arctic areas in North America than for their southern counterparts. Gonorrhea rates reported for northern Canada and Greenland were also much higher than for their southern counterparts, although rates reported for Alaska were not very high. In 1741, Hans Egede, the first missionary to Greenland noted that “It is strange ... that even though [Greenlanders] have free intercourse with other people, these are not infected” (11). However, for the past several years Greenland has reported chlamydial infection rates

Table 1. Comparison of sexually transmitted infection rates reported for North America's Arctic countries, 2003–2006

Yearly rate*	United States	Alaska, USA	Canada	Northern territories,†	Southern provinces,†	Denmark	Greenland
Chlamydial infection							
2003	301.7	601.1	189.4	1,433	185	342	3,255
2004	316.7	609.4	197.1	1,805	195	401	3,208
2005	332.5	664.4	200.4	1,952	195	441	4,762
2006	347.8	682	202.2	1,922	197	458	4,527
2006 standardized‡	470.9	715	205	1,693	200	681	5,543
Gonorrhea							
2003	115.2	88.3	26.0	264	25	3.5	1,162
2004	113.5	87.4	28.9	215	29	7.7	1,148
2005	115.6	91.5	27.8	212	28	8.2	1,350
2006	120.9	95	33.1	281	32	7.5	1,418
2006 standardized	164.4	101	33	247	32	6.5	1,738

\*Per 100,000 population. Data from Centers for Disease Control and Prevention, 2006 (12); Public Health Agency of Canada, 2007 (14); Office of the Chief Medical Officer in Greenland (15,16); Statens Serum Institute surveillance Epi-data online ([www.ssi.dk](http://www.ssi.dk)).

†Canadian Northern territories: Yukon Territory, Northwest Territories, and Nunavut; Southern provinces: British Columbia, Alberta, Saskatchewan, Manitoba, Ontario, Quebec, Newfoundland, Nova Scotia, New Brunswick, and Prince Edward Island.

‡2006 standardized estimates are directly standardized to the year 2000 US population distributed by age and sex.

Table 2. Chlamydial infection and gonorrhea rates per 100,000 population by age, sex, and race reported for Alaska, 2006\*

Characteristic	Chlamydial infection rates		Gonorrhea rates	
	M	F	M	F
Age, y				
15–19	966	4,158	78	346
20–24	2,673	4,990	344	496
25–29	1,250	2,253	185	309
30–34	607	854	162	162
35–39	269	420	134	81
Race				
White	235	389	28	41
Alaska Native/ American Indian	927	3,012	153	344

\*Source: (13).

≈10× higher, and gonorrhea rates ≈100× higher, than rates reported for Denmark and the highest rates of both infections in the North American Arctic (Table 1).

Chlamydial infection and gonorrhea rates reported for the Arctic and sub-Arctic are very high for both men and women, although the highest incidence of infection is predominantly reported for young women in their early 20s (Tables 2–4). True rates could be higher than reported for a variety of reasons. As in other settings, asymptomatic infection is high for both men and women and can result in missed cases. How much knowledge exists in remote communities about STIs, their symptoms, and what to do if one suspects he or she has an infection is unclear. Even if a person suspects that he or she has an infection, accessing healthcare can be a challenge since many of the Arctic communities are remote fly-in communities with limited

healthcare resources. Additionally, many Arctic residents spend their summers away hunting or whaling, usually at great distances from their communities, and certainly far away from a healthcare provider. Another barrier to care in small communities is the issue of confidentiality and the common perception that it can be breached easily. This results in delayed healthcare seeking or missed infections. Partner notification can also be hindered by cultural norms and taboos. For instance, in some communities, talking about something can be regarded as the same as wishing it upon the people. Therefore there can be a reluctance or even movement against talking about STIs or naming sexual contacts. Finally, reporting infections can become a challenge in an already overtaxed healthcare system with limited infrastructure.

STI rates are quite variable across the North American Arctic and sub-Arctic (Tables 1–4). Access to healthcare and reporting differences could explain some of the difference in rates. For instance, Greenland has universal healthcare. Canada has universal healthcare, but it differs for on-reserve and off-reserve aboriginal people. Alaska only has universal healthcare for indigenous people. These different healthcare coverage strategies could affect the healthcare-seeking behavior of the populations that live with them. Another nuance of northern rates is the small underlying populations from which cases arise. The addition of 1 new case can result in a large change in the rate of infection. Additionally, because no international surveillance system is in place to monitor STIs, the information collected is not standardized between the countries. For instance, in the United States, the only country that collects racial informa-

Table 3. Chlamydial infection and gonorrhea rates per 100,000 population by age and sex reported for northern territories (NT) and southern provinces (SP) in Canada, 2004–2006\*

Characteristic	2004				2005				2006			
	NT		SP		NT		SP		NT		SP	
	M	F	M	F	M	F	M	F	M	F	M	F
Chlamydia, age, y												
<14	22	319	1	19	22	361	0.8	18	8	296	1	16
15–19	3,050	10,014	276	1,428	3,193	11,866	270	1,367	3,374	10,771	278	1,329
20–24	4,778	9,408	695	1,478	5,255	8,893	701	1,470	4,982	9,431	703	1,475
25–29	3,154	4,492	405	552	3,623	4,435	423	562	3,192	5,024	419	592
30–39	1,292	1,913	141	158	1,461	1,856	157	158	1,697	1,812	164	170
40–59	338	359	31	22	486	450	34	21	399	432	36	24
≥60	120	142	4	1	90	215	4	1	84	126	5	2
Total	1,190	2,451	128	260	1,339	2,595	132	256	1,312	2,556	134	258
Gonorrhea, age, y												
<14	0	23	0	2	7	38	0	3	0	23	0	4
15–19	376	761	57	124	437	737	53	112	671	1,473	63	132
20–24	820	763	126	99	968	650	118	102	1,038	1,246	132	118
25–29	738	332	91	43	689	433	93	41	557	526	104	57
30–39	306	169	65	14	311	86	61	15	267	234	67	21
40–59	137	15	23	3	71	45	23	2	86	25	27	0
≥60	24	0	4	0	22	0	4	0	42	25	5	0
Total	240	189	37	21	239	184	35	20	251	312	40	25

\*See (14).

Table 4. Chlamydial infection and gonorrhea rates per 100,000 population by age and sex reported for Greenland (GLD) and Denmark (DK), 2004–2006

Characteristic	2004				2005				2006			
	GLD		DK		GLD		DK		GLD		DK	
	M	F	M	F	M	F	M	F	M	F	M	F
Chlamydia, age, y												
15–19	9,378	20,332	944	3,361	7,986	31,383	1,243	3,891	12,462	27,125	1,345	4,095
20–24	13,229	16,890	1,966	3,526	9,003	20,594	2,264	3,720	17,154	21,854	2,391	3,768
25–29	6,444	8,590	978	1,284	5,776	11,006	1,089	1,322	10,837	10,445	1,114	1,335
≥30	916	1,229	86	41	805	1,005	85	87	1,630	1,507	93	88
Total	2,481	4,158	287	511	3,852	5,597	324	554	3,704	5,468	343	571
Gonorrhea, age, y												
15–19	3,714	7,346	13.9	7.7	2,141	7,801	11.5	4.0	5,360	8,763	11.1	6.5
20–24	4,663	4,450	46.4	7.4	2,993	5,648	39.9	13.1	5,994	5,858	32.5	16.0
25–29	3,056	1,795	51.3	4.5	1,984	3,184	44.9	4.7	3,755	2,374	39.4	10.8
≥30	493	434	10.7	1.2	224	312	14.1	1.1	363	498	12.4	1.2
Total	1,047	1,301	13.6	1.9	1,174	1,550	14.6	2.0	1,252	1,609	12.5	2.6

tion as part of its surveillance program, chlamydial infection, gonorrhea, and syphilis rates reported for American Indians and Alaska Natives were recently reported to be 2–6× higher than rates reported for non-Hispanic whites (18). In Canada, STI rates are suspected of being higher for aboriginal people, but no data exist to confirm this hypothesis. In Greenland, 89% of the population is Kalaallit Inuit, making it of arguable importance to collect racial information for Greenland.

There is a dearth of research pertaining to the factors that contribute to sexual health and STIs among aboriginal people (18,19). Chlamydial infection and gonorrhea are major causes of ectopic pregnancy in the Canadian Arctic (17), and STIs are highest for Canadian aboriginal people 15–24 years of age (20). Westernization, culture, and identity have been suggested as possible factors influencing STI transmission among Inuit youth in northern Canada (20); however, research is still needed to provide evidence for this hypothesis. Most other studies have focused on HIV and AIDS (19,21).

In Greenland, much of the STI research has also focused on HIV/AIDS. HIV/AIDS came late to Greenland compared to the rest of Europe and has remained limited to a heterosexual, alcohol-abusing group of persons of low socioeconomic status living in 2 communities in western Greenland. One reason that HIV, unlike other STIs, has not become a widespread epidemic across Greenland is because the prevalence of needle sharing and men who have sex with men is limited, considerably affecting the modes of transmission (11). However, research on HIV transmission among heterosexual persons, as well as the increased risk for co-infection with STIs and HIV, suggests that chlamydial and gonorrhea rates in Greenland are a public health concern that warrant further investigation (22–24).

As suggested by Steenbeek et al. (20), colonization and westernization in the Arctic may be responsible for increased rates of STIs for Arctic communities. We fur-

ther hypothesize that these factors are contributing to separately high STI rates in the Arctic through individual, familial, social, cultural, and environmental domains. We also hypothesize that high STI rates may only be a marker of greater underlying public health concerns such as substance abuse, poor mental health, and the legacy of historic trauma.

### Implications for Future Research

We propose that community-based participatory research (CBPR) is an appropriate approach to address sexual health and STIs in the Arctic. Sexual and reproductive health data for aboriginal populations are often not reported in national surveillance and survey reports (25). Also, indigenous communities have historically been reluctant to participate in research projects because traditional research methods, which emphasize the researcher as “the expert,” have not engaged indigenous communities in designing and implementing research projects (25). CBPR has been identified as an effective and essential strategy for conducting research with indigenous peoples because of its emphasis on community participation to build ownership of research projects and community-based interventions as well as empowering the community to address its health disparities (26,27).

Several components of CBPR support its use as a methodologic framework for conducting research in aboriginal communities. First, CBPR engages aboriginal or indigenous people in full and equal partnership with those communities in efforts to observe and respect tribal sovereignty and the right to self-determination (28). Second, the growing interest in addressing the interrelatedness of historic trauma and health disparities in indigenous populations and the inherent complexities of unraveling the interconnected components and concepts related to historic trauma and health can best be understood by discussions and conversations with indigenous communities (29,30). Third, a legacy of harm



from past research, as well as mistrust of researchers, warrants the use of CBPR as a means to ensure that all phases of a research project, from the development of research questions to research design and data collection methods to dissemination of results, have community input and approval (26,31). Fourth, CBPR provides a forum to ensure timely communication of research results to the community by using information dissemination mechanisms that best meet the community's needs. Finally, the limited research on sexual health among indigenous populations primarily focuses on problem theory that provides insights into the predisposing, enabling, and reinforcing factors related to engagement in high-risk sexual behavior among aboriginal communities. However, emerging evidence in the field of aboriginal sexual health suggests that a risk-based approach to understanding sexual behavior in these communities not only has a narrow and negative focus, with scant opportunities for indigenous groups to capitalize on their strengths, but also is not congruent with indigenous cultural and social beliefs and historical experiences (32). CBPR, because of its collaborative nature, empowers community members to capitalize on the strengths and resources available in their community.

## Conclusion

The use of CBPR as a framework in which to conduct sexual health research with and among indigenous populations is a promising approach that joins the strengths and skills of researchers with local knowledge, wisdom, traditions, and resources. The CBPR approach is much like taking a Bayesian approach to study design, data collection, analysis, interpretation, dissemination, and follow-up. Researchers provide global (prior) knowledge that is then integrated and updated with local (likelihood) knowledge provided by the community to produce a more holistic model of health. This approach means that study designs can be more effective, data collection can be more accurate and complete, interpretation of the results can be more insightful and relevant, dissemination of the study results can be more efficient and translated at the appropriate level for the community by community members, and interventions can be more effective, culturally appropriate, and sustainable. Community involvement in the project can also help facilitate translation of the research findings into clinical and political practice.

## Acknowledgments

We thank Gaya Jayaraman for providing us with access to unpublished surveillance data for Canada, Flemming Stenz and Jytte Hey for providing us with access to published and unpublished data for Greenland, Susan Cowan and Steen Hoffman for access to unpublished data for Denmark, and Jessica Leston for her assistance accessing data for Alaska.

This research was supported in part by the University of Toronto and Montana State University.

Dr Gesink Law is an epidemiologist and assistant professor in the Department of Public Health Sciences at the University of Toronto. She has research interests in sexual, reproductive, and aboriginal health and, with Dr Rink, has been building a research program in sexual health and STIs with northern frontier and aboriginal communities in North America.

## References

1. Arctic Climate Impact Assessment. Impacts of a warming Arctic: Arctic climate impact assessment. New York: Cambridge University Press; 2004.
2. Gaydos CA, Kent CK, Rietmeijer CA, Willard NJ, Mrazek JM, Chapin JB, et al. Prevalence of *Neisseria gonorrhoeae* among men screened for *Chlamydia trachomatis* in four United States cities, 1999–2003. *Sex Transm Dis*. 2006;33:314–9.
3. Levine SB, Coupey SM. Adolescent substance use, sexual behavior, and metropolitan status: is “urban” a risk factor? *J Adolesc Health*. 2003;32:350–5.
4. Adimora AA, Schoenbach VJ, Doherty IA. HIV and African Americans in the southern United States: sexual networks and social context. *Sex Transm Dis*. 2006;33(Suppl):S39–45.
5. Thomas JC. From slavery to incarceration: social forces affecting the epidemiology of sexually transmitted diseases in the rural South. *Sex Transm Dis*. 2006;33(Suppl):S6–10.
6. Aral SO, O’Leary A, Baker C. Sexually transmitted infections and HIV in the southern United States: an overview. *Sex Transm Dis*. 2006;33(Suppl):S1–5.
7. Farley TA. Sexually transmitted diseases in the Southeastern United States: location, race, and social context. *Sex Transm Dis*. 2006;33(Suppl):S58–64.
8. Rhodes SD, Eng E, Hergenrather KC, Remnitz IM, Arceo R, Montañó J, et al. Exploring Latino men’s HIV risk using community-based participatory research. *Am J Health Behav*. 2007;31:146–58.
9. Johnson LF, Coetzee DJ, Dorrington RE. Sentinel surveillance of sexually transmitted infections in South Africa: a review. *Sex Transm Infect*. 2005;81:287–93.
10. Sangani P, Rutherford G, Wilkinson D. Population-based interventions for reducing sexually transmitted infections, including HIV infection. *Cochrane Database Syst Rev*. 2004;2:CD001220.
11. Bjerregaard P, Mulvad G, Olsen J. Studying health in Greenland: obligations and challenges. *Int J Circumpolar Health*. 2003;62:5–16.
12. Centers for Disease Control and Prevention. Sexually transmitted disease surveillance, 2006. Atlanta: US Department of Health and Human Services; 2007.
13. Gessner BD, McLaughlin J, eds. *Chlamydia trachomatis*—Alaska, 2006. State of Alaska Epidemiology Bulletin. 2007 [cited 2007 Oct 25]. Available from <http://www.epi.alaska.gov>
14. Public Health Agency of Canada. STI data tables. Surveillance and Epidemiology Section, Community Acquired Infections Division, Centre for Infectious Disease Prevention and Control. 2007 [cited 2007 Oct 25]. Available from [http://www.phac-aspc.gc.ca/std-mts/stddata\\_pre06\\_04/index\\_tab\\_e.htm](http://www.phac-aspc.gc.ca/std-mts/stddata_pre06_04/index_tab_e.htm)
15. Office of the Chief Medical Officer in Greenland. Ukiumoortumik Nalunaarut Årsberetning 2004. Nuuk, Greenland: Embedslægeinstitutionen i Grønland; 2005.
16. Office of the Chief Medical Officer in Greenland. Ukiumoortumik Nalunaarut Årsberetning 2003. Nuuk, Greenland: Embedslægeinstitutionen i Grønland; 2004.
17. Orr PH, Brown R. Incidence of ectopic pregnancy and sexually transmitted disease in the Canadian central Arctic. *Int J Circumpolar Health*. 1998;57(Suppl 1):127–34.

18. Wong D, Swint E, Paisano EL, Cheek JE. Indian Health surveillance report sexually transmitted diseases 2004. Atlanta: Centers for Disease Control and Prevention and Indian Health Services; 2006.
19. Kaufman CE, Shelby L, Mosure DJ, Marrazzo J, Wong D, de Ravello L, et al. Within the hidden epidemic: sexually transmitted diseases and HIV/AIDS among American Indians and Alaska Natives. *Sex Transm Dis.* 2007;34:767-77.
20. Steenbeek A, Tyndall M, Rothenberg R, Sheps S. Determinants of sexually transmitted infections among Canadian Inuit adolescent populations. *Public Health Nurs.* 2006;23:531-4.
21. Larkin J, Flicker S, Koleszar-Green R, Mintz S, Dagnino M, Mitchell C. HIV risk, systemic inequities, and Aboriginal youth: widening the circle for HIV prevention programming. *Can J Public Health.* 2007;98:179-82.
22. Elwy AR, Hart GJ, Hawkes S, Petticrew M. Effectiveness of interventions to prevent sexually transmitted infections and human immunodeficiency virus in heterosexual men: a systematic review. *Arch Intern Med.* 2002;162:1818-30.
23. Nusbaum MR, Wallace RR, Slatt LM, Kondrad EC. Sexually transmitted infections and increased risk of co-infection with human immunodeficiency virus. *J Am Osteopath Assoc.* 2004;104:527-35.
24. Lapidus JA, Bertolli J, McGowan K, Sullivan P. HIV-related risk behaviors, perceptions of risk, HIV testing, and exposure to prevention messages and methods among urban American Indians and Alaska Natives. *AIDS Educ Prev.* 2006;18:546-59.
25. Hellerstedt WL, Peterson-Hickey M, Rhodes KL, Garwick A. Environmental, social, and personal correlates of having ever had sexual intercourse among American Indian youths. *Am J Public Health.* 2006;96:2228-34.
26. Holkup PA, Tripp-Reimer T, Salois EM, Weinert C. Community-based participatory research: an approach to intervention research with a Native American community. *ANS Adv Nurs Sci.* 2004;27:162-75.
27. Smith A, Christopher S, McCormick AK. Development and implementation of a culturally sensitive cervical health survey: a community-based participatory approach. *Women Health.* 2004;40:67-86.
28. Mail PD, Conner J, Conner CN. New collaborations with native Americans in the conduct of community research. *Health Educ Behav.* 2006;33:148-53.
29. Whitbeck LB, Adams G, Hoyt D, Chen X. Conceptualizing and measuring historical trauma among American Indian people. *Am J Community Psychol.* 2004;33:119-30.
30. Jervis L, Beals J, Croy C, Klein S, Manson S. Historical consciousness among two American Indian tribes. *Am Behav Sci.* 2006;50:526-49.
31. Quigley D. A review of improved ethical practices in environmental and public health research: case examples for native communities. *Health Educ Behav.* 2006;33:130-47.
32. Whitbeck L. Some guiding assumptions and a theoretical model for developing culturally specific preventions with Native American people. *J Community Psychol.* 2006;34:183-92.

Address for correspondence: Dionne Gesink Law, Department of Public Health Sciences, Health Sciences Bldg, University of Toronto, 155 College St, 6th Floor, Toronto, Ontario M5T 3M7, Canada; email: dionne.gesinklaw@utoronto.ca

# EMERGING INFECTIOUS DISEASES

Full text free online at  
[www.cdc.gov/eid](http://www.cdc.gov/eid)

The print journal is available at no charge to public health professionals

YES, I would like to receive Emerging Infectious Diseases.

Please print your name and business address in the box and return by fax to 404-639-1954 or mail to

EID Editor  
CDC/NCID/MS D61  
1600 Clifton Road, NE  
Atlanta, GA 30333

Moving? Please give us your new address (in the box) and print the number of your old mailing label here \_\_\_\_\_

**EID**  
*Online*  
[www.cdc.gov/eid](http://www.cdc.gov/eid)