

results for Japanese encephalitis virus-specific IgM. Of the 11 patients, 10 had no history of travel to India or other dengue-endemic countries. DF or DHF was initially diagnosed for 7 patients, and viral encephalitis, typhoid fever, or viral fever was diagnosed for others without serologic tests. Reverse transcription-PCR and virus isolation were performed at Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan, but the dengue virus genome was not detected, and no virus was isolated, likely because sample collection was delayed and the sample was transported to Japan in a deteriorated condition.

DF/DHF have been considered to be a possible public health threat to Nepal because DF/DHF epidemics have occurred recently in India and Pakistan, which reported several thousand cases and >100 deaths (6). The first DF case in Nepal was reported in 2004 (7). Further, the first DENV-2 strain of Nepal origin was isolated from a Japanese traveler who visited Nepal and in which DF developed after the patient returned to Japan. The isolated DENV-2 (GenBank accession no. AB194882) was 98% homologous with DENV-2 isolated in India (8). The prevalence of dengue virus antibody was reported to be 10.4% in the southwestern region of Nepal (9). These reports suggest that dengue virus has been circulating in Nepal for several years. Thus, DF/DHF has likely been misdiagnosed and illness caused by dengue virus underestimated in Nepal. In contrast, Japanese encephalitis has been a public health problem in southwestern region of Nepal, and large epidemics have occurred almost every year since 1978 (10). Nepal has no dengue surveillance programs, and health professionals do not usually consider dengue as a differential diagnosis.

The emergence occurred in the lowland Terai belt region, which borders the state of Bihar, India. The *Aedes* mosquito is known to persist in

this region. The emerging DENV-2 is likely to have been introduced into Nepal from India.

**Basu Dev Pandey,*
Kouichi Morita,†
Santa Raj Khanal,‡
Tomohiko Takasaki,§
Isao Miyazaki,¶
Tetsuro Ogawa,¶
Shingo Inoue,†
and Ichiro Kurane§**

*Tropical and Infectious Disease Hospital, Kathmandu, Nepal; †Nagasaki University, Nagasaki, Japan; ‡Tribhuban University, Kathmandu, Nepal; §National Institute of Infectious Diseases, Tokyo, Japan; and ¶Pentax Company Limited, Tokyo, Japan

References

1. World Health Organization. Dengue and dengue haemorrhagic fever. Fact sheet no.117; revised 2002 Apr [cited 2006 Oct 31]. Available from <http://www.who.int/mediacentre/factsheets/fs117/en>
2. World Health Organization. Dengue outbreak in Bhutan. Communicable Disease Newsletter. 2007;4(1).
3. Islam MA, Ahmed M, Begum N, Choudhury N, Khan A, Parquet C, et al. Molecular characterization and clinical evaluation of dengue outbreak in 2002 in Bangladesh. *Jpn J Infect Dis.* 2006;59:85–91.
4. Jamil B, Hasan R, Zafar A, Bewley K, Chamberlain J, Mioulet V, et al. Dengue virus serotype, Karachi, Pakistan. *Emerg Infect Dis.* 2007;13:182–3.
5. Pandey B, Yamamoto A, Morita K, Kurosawa Y, Rai S, Adhikari S, et al. Serodiagnosis of Japanese encephalitis among Nepalese patients by the particle agglutination assay. *Epidemiol Infect.* 2003;131:881–5.
6. Gupta E, Dar L, Kapoor G, Broor S. The changing epidemiology of dengue in Delhi. *Virology.* 2006;3:92.
7. Pandey BD, Rai SK, Morita K, Kurane I. First case of dengue in Nepal. *Nepal Med Coll J.* 2004;6:157–9.
8. Takasaki T, Kotaki A, Nishimura K, Sato Y, Tokuda A, Lim C, et al. Dengue virus type 2 from an imported dengue patient in Japan. First isolation of dengue virus from Nepal. *J Travel Med.* 2007;14:445–8.
9. Sherchand JB, Pandey BD, Haruki K, Jimba M. Sero-diagnosis of Japanese encephalitis and dengue virus infection from clinically suspected patients of Nepal. *J Inst Med.* 2001;23:25–31.
10. Epidemiology and Disease Control Division, Ministry of Health Nepal. The annual report of malaria, Kala-azar and Japanese encephalitis in Nepal, 2005. Kathmandu, Nepal: Ministry of Health; 2006.

Address for correspondence: Basu Dev Pandey, Tropical and Infectious Disease Hospital, Teku Road, GPO Box 9045, Kathmandu, Nepal; email: basupandey@wlink.com.np

Human Tuberculosis Caused by *Mycobacterium bovis*, Taiwan

To the Editor: *Mycobacterium bovis* is one of the causative agents of tuberculosis (TB) in humans and animals. Drinking unpasteurized milk, eating undercooked meat, and close contact with infected animals are the main sources of infection for humans. Currently, 119 *M. bovis* spoligotypes are contained in the fourth international spoligotyping database (SpolDB 4) and are categorized into 3 main sublineages corresponding to ST prototypes 482, 683, and 479 (1).

Although an *M. bovis* surveillance program for farm animals has been implemented by the Taiwan Council of Agriculture, no surveillance system exists for human TB cases caused by *M. bovis*. To monitor the epidemiology of *M. bovis* in domestic animals, a regular tuberculin skin test (TST) is compulsory for cattle and sheep and optional for deer in Taiwan (2). In 2005, screening of *Mycobacterium* spp. infections by TST was performed for 111,412 cattle and 73,396 caprine and ovine herds, of which 188 (0.17%) and 148 (0.2%), respectively, were positive (2). We used spacer oligonucleotide typing (spoligotyping) and

mycobacterial interspersed repetitive units–variable number tandem repeat (MIRU-VNTR) methods to investigate human TB caused by *M. bovis* in Taiwan.

During 2004–2005, a total of 3,321 mycobacteria isolates from individual patients were sent to the reference laboratory for strain typing. Of the 3,321 patients, 2,427 (73.1%) were male, 903 (27.2%) were from eastern Taiwan, and 513 (15.4%) were aboriginal persons. The mean age of the patients was 58.7 years: 224 (6.7%) were <25 years of age, 667 (20.1%) were 25–44 years of age, 906 (27.3%) were 45–64 years of age, and 1,524 (45.9%) were ≥65 years of age.

Isolates were screened by using a GenoType kit (3) and multiplex PCR (4). To differentiate between *M. bovis* and *M. bovis* BCG strains, presence or absence of region of difference 1 was analyzed (5). Spoligotyping was performed with a commercial kit (Isogen Bioscience BV, Maarssen, the Netherlands) following the manufacturer's instructions (6). Spoligotyping profiles were analyzed by using Bionumerics software, version 4.51 (Applied Maths, Kortrijk, Belgium). The resolved spoligotype was designated by comparing it to SpoIDB4 (1). The MIRU-VNTR assay was performed by using a modified, high-throughput, 15-loci MIRU typing system that we developed (7).

Of the 3,321 patient isolates, 3,306 (99.5%) were *M. tuberculosis* and only 15 (0.5%) were *M. bovis*. Mean age of the 15 *M. bovis*-infected patients was 62.2 years (Table). Twelve (80%) patients were male, and 3 (20%) were female. Of these 15 patients, 10 (66.7%) had newly diagnosed TB and 5 (33.3%) had been treated with anti-TB drugs. Most (11/15, 73%) of the patients were from eastern Taiwan, where the reporting rate for TB was the highest among the 4 regions of Taiwan; 60% (9/15) were aboriginal persons. Of the 15 patients, 13 (86.7%) had pulmonary TB, 1 had both pulmonary and extrapulmonary TB, and 1 had extrapulmonary TB. Only 2 patients (cases 2 and 3) had known contact with farm animals. Univariate analysis showed that region (eastern region 1.2% vs. other region 0.2%; odds ratio [OR] 7.4, 95% confidence interval [CI] (2.4–23.4) and ethnicity (aboriginal 1.8% vs. nonaboriginal 0.2%; OR 8.3, 95% CI 3.0–23.5) were associated with *M. bovis* infection but not age and sex. The association between region and *M. bovis* disappeared after controlling for age and ethnicity. Aboriginal ethnicity was the only factor significantly associated with TB caused by *M. bovis* after controlling for age (adjusted OR 12.7, 95% CI 4.2–38.9).

Spoligotyping profiles of the 15 *M. bovis* isolates were typical of *M.*

bovis with the absence of spacers 3, 6, 9, 16, 21, and 39–43 (8). Although 15 cases were reported separately from different regions of Taiwan, only 1 spoligotyping profile was identical to spoligotype ST 684 of the *bovis* sublineage. In addition, we identified 2 similar MIRU-VNTR profiles: 523–23232–42533–22 (13/14, 92.9%, cases 1–9, 11–13, and 15) and 523–22232–42523–22 (1/14, 7.1%, case 14). We were unable to obtain sufficient DNA from 1 strain (case 10) for MIRU-VNTR typing.

Aboriginal persons in Taiwan were more likely to have TB caused by *M. bovis* and had a 5-fold higher reporting rate for TB (9) than nonaboriginal persons. Environmental and genetic factors may be associated with a higher reporting rate for TB among aboriginal populations (10), but the contribution of *M. bovis* infection needs to be investigated. Because only 1 major spoligotype and 2 similar MIRU patterns were found in this case series, spread of a predominant clone in Taiwan is likely. In addition, because of insufficient epidemiologic data, we were unable to determine the proportion of cases caused by reactivation of latent infection and those caused by recent transmission. In our study population, *M. bovis* infection in humans appeared to be predominately indigenous in Taiwan because no imported case was noted. We are now

Table. Characteristics of 15 patients with tuberculosis caused by *Mycobacterium bovis*, Taiwan, 2004–2005

Patient no.	Age at diagnosis, y	Sex	Type of tuberculosis	Ethnicity	Region
1	53	M	Pulmonary	Aboriginal	Northern
2	71	M	Pulmonary	Han	Central
3	76	M	Pulmonary	Han	Central
4	84	M	Pulmonary	Han	Southern
5	71	F	Pulmonary	Aboriginal	Eastern
6	55	M	Pulmonary	Aboriginal	Eastern
7	51	M	Pulmonary	Aboriginal	Eastern
8	65	F	Pulmonary	Aboriginal	Eastern
9	51	M	Pulmonary	Han	Eastern
10	90	M	Pulmonary	Han	Eastern
11	34	M	Pulmonary	Aboriginal	Eastern
12	81	M	Pulmonary	Han	Eastern
13	64	F	Pulmonary	Aboriginal	Eastern
14	44	M	Bladder/urinary	Aboriginal	Eastern
15	43	M	Pulmonary/miliary	Aboriginal	Eastern

genotyping *M. bovis* strains isolated from farm animals to help elucidate the source of infection and transmission of *M. bovis* in Taiwan.

Acknowledgments

We thank Chen-Che Chiu and Pei-Ju Chin for excellent technical assistance.

This work was supported by grant DOH95-DC-2011 from the Taiwan Centers for Disease Control, Department of Health, and joint grant 95-0324-19-F-01-00-00-00-35 from National Science Council and Department of Health, Taiwan, Republic of China.

Ruwen Jou,* Wei-Lun Huang,* and Chen-Yuan Chiang†

*Centers for Disease Control, Taipei, Taiwan, Republic of China; and †International Union Against Tuberculosis and Lung Disease, Paris, France

References

1. Brudey K, Driscoll JR, Rigouts L, Prodinger WM, Gori A, Al-Hajj SA, et al. *Mycobacterium tuberculosis* complex genetic diversity: mining the fourth international spoligotyping database (SpolDB4) for classification, population genetics and epidemiology. *BMC Microbiol.* 2006;6:23–39.
2. Bureau of Animal and Plant Health Inspection and Quarantine, Council of Agriculture, Executive Yuan, Taiwan. Annual report. Taipei: The Bureau; 2005.
3. Richter E, Weizenegger S, Rusch-Gerdes S, Niemann S. Evaluation of genotype MTBC assay for differentiation of clinical *Mycobacterium tuberculosis* complex isolates. *J Clin Microbiol.* 2003;41:2672–5.
4. Yeboah-Manu D, Yates MD, Wilson SM. Application of a simple multiplex PCR to aid in routine work of the mycobacterium reference laboratory. *J Clin Microbiol.* 2001;39:4166–8.
5. Behr MA, Wilson MA, Gill WP, Salamon H, Schoolnik GK, Rane S, et al. Comparative genomics of BCG vaccines by whole-genome DNA microarray. *Science.* 1999;284:1520–3.
6. Kamerbeek J, Schouls L, Kolk A, van Agterveld M, van Soolingen D, Kuijper S, et al. Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J Clin Microbiol.* 1997;35:907–14.
7. Chin PJ, Jou R. A modified automated high-throughput mycobacterial interspersed repetitive unit method for genotyping *Mycobacterium tuberculosis*. *Diagn Microbiol Infect Dis.* 2005;53:325–7.
8. Aranaz A, Lie'bana E, Mateos A, Dominguez L, Vidal D, Domingo M, et al. Spacer oligonucleotide typing of *Mycobacterium bovis* strains from cattle and other animals: a tool for studying epidemiology of tuberculosis. *J Clin Microbiol.* 1996;34:2734–40.
9. Hsu YH, Chen CW, Sun HS, Jou R, Lee JJ, Su IJ. Association of NRAMP 1 gene polymorphism with susceptibility to tuberculosis in Taiwanese aborigines. *J Formos Med Assoc.* 2006;105:363–9.
10. Wang WH, Chang SJ, Wang TN, Cheng LS, Feng YP, Chen CJ, et al. Complex segregation and linkage analysis of familial gout in Taiwanese aborigines. *Arthritis Rheum.* 2004;50:242–6.

Address for correspondence: Ruwen Jou, Reference Laboratory of Mycobacteriology, Research and Diagnostic Center, Department of Health, Centers for Disease Control, 161 Kun-Yang St, Nan-Kang, Taipei 115, Taiwan, Republic of China; email: rwj@cdc.gov.tw

Marine Mammal *Brucella* Genotype Associated with Zoonotic Infection

To the Editor: Brucellosis is a zoonotic disease that remains endemic to many parts of the world. There are 6 classic *Brucella* species described with different preferred hosts. Human disease is most commonly associated with consumption of unpasteurized dairy products or with occupational exposure for veterinarians, agricultural workers, laboratory workers, meat industry workers, and hunters.

In recent years, it has become clear that novel members of the genus, yet to be formally named, are associated with a variety of marine mammal species, particularly dolphins, por-

poises, and seals (1). To date there are 3 reports in the literature of naturally acquired infection of humans with *Brucella* species originating from marine mammals (2,3) One other case, representing infection of a laboratory worker, has also been reported (4). Two of the naturally acquired cases were reported in Peru (2). One person had consumed raw shellfish and swam in the Pacific Ocean but did not report any direct contact with marine mammals; the second person reported infrequent visits to the coast and no contact with marine mammals but had consumed raw shellfish. An additional naturally acquired case was recently reported from New Zealand, where extensive molecular testing characterized the strain involved as a marine mammal type (3). This patient again reported no exposure to marine mammals but did report that he fished regularly, had contact with uncooked fish bait, and consumed raw snapper. The cases in Peru were notable for severe, atypical symptoms; both patients had symptoms of neurobrucellosis. The New Zealand case was associated with spinal osteomyelitis (3). In contrast, the laboratory-acquired infection was mild and uncomplicated (4).

We have characterized these isolates by a variety of molecular approaches in conjunction with ongoing studies, which examine genetic diversity within *Brucella* species isolated from marine mammals. Multilocus sequence analysis (5) showed that all 3 isolates from naturally acquired human infection with *Brucella* species from marine mammals shared an identical genotype (ST27). In previous characterization of 56 *Brucella* isolates from marine mammals, ST27 was found only once. Strain F5/99, originally isolated from an aborted bottlenose dolphin fetus off the western coast of the United States (6), shares this genotype. Use of an alternative typing approach, based on restriction fragment length polymorphism analysis of outer membrane protein-encoding genes