
Seasonal Cholera from Multiple Small Outbreaks, Rural Bangladesh

O. Colin Stine,* Munirul Alam,† Li Tang,*
G. Balakrish Nair,† A. Kasem Siddique,†
Shah M. Faruque,† Anwar Huq,‡ Rita Colwell,*§
R. Bradley Sack,§ and J. Glenn Morris, Jr*¶

Clinical and environmental *Vibrio cholerae* organisms collected from February 2004 through April 2005 were systematically isolated from 2 rural Bangladeshi locales. Their genetic relatedness was evaluated at 5 loci that contained a variable number of tandem repeats (VNTR). The observed minimal overlap in VNTR patterns between the 2 communities was consistent with sequential, small outbreaks from local sources.

Cholera is a major cause of illness in the developing world. The World Health Organization reported in 2006 that 236,896 cases of cholera occurred in 52 countries, a 79% increase over 2005 (1). Although major advances in the understanding of the molecular basis of *Vibrio cholerae* pathogenicity have been made, including defining the environmental reservoirs for the microorganism (2–4), we do not fully understand the cause of seasonal epidemics in cholera-endemic areas nor the factors that drive epidemics. Specifically, whether these seasonal epidemics arise from a single clonal strain or reflect superimposition of multiple small outbreaks is not clear.

The Study

From February 2004 through April 2005, we systematically collected clinical and environmental *V. cholerae* from Bakerganj and Mathbaria, 2 small communities 50 miles apart in the southern part of coastal Bangladesh. Samples were collected on 3 consecutive days every 2 weeks throughout the year. Clinical isolates were collected from ≈20% of all patients who had symptoms of cholera when seen at the local clinics. Environmental isolates were cultured from water, sediment, and plankton samples taken at 6 sites (ponds or river sites) in each of the 2 communities. The same sites were used throughout the 15-month study,

*University of Maryland School of Medicine, Baltimore, Maryland, USA; †International Center for Diarrheal Disease Research, Dhaka, Bangladesh; ‡University of Maryland, College Park, Maryland, USA; §Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA; and ¶University of Florida, Gainesville, Florida, USA

and the same method was applied at all sites and across all time points. Isolation was performed by standard culture methods, and *V. cholerae* was identified by a combination of biochemical (5), molecular, and serologic techniques (6). All samples were collected according to protocols approved by Institutional Review Boards at Johns Hopkins University, University of Maryland, and the International Centre for Diarrheal Disease Research, Bangladesh.

For multilocus sequence typing (MLST) and variable number of tandem repeat (VNTR) determinations, each locus was PCR amplified by using standard conditions and appropriate primers from the literature (7) (see online Technical Appendix, available from www.cdc.gov/EID/content/14/5/831-Techapp.pdf). The resulting fragments were sequenced by using Big Dye Kit (Applied Biosystems, Foster City, CA, USA). Trace files were generated by using an ABI 3730xl automatic sequencer and read using either 1) the Phred (8,9), Phrap (www.washington.edu), or Consed (10) package or 2) Sequencher (AGCT, Gene Codes Corporation, Ann Arbor, MI, USA).

A total of 391 environmental and clinical isolates of *V. cholerae* were collected and identified from February 2004 through April 2005. Of these, 267 environmental isolates were identified as belonging to non-O1 and non-O139 serogroups and did not carry the gene for cholera toxin (*ctx*). Analysis of these 267 by MLST (using the 7 loci identified previously [7]) yielded a genetic background that was distinct from that of the clinical/epidemic strains. The other 68 (20%) of 335 environmental *V. cholerae* isolates shared a genetic background identical or nearly identical to clinical/epidemic *V. cholerae*. These 68 and all 56 clinical isolates collected (all of which were related by MLST) were further analyzed by examining 5 VNTR loci.

Sequence typing was based on 5 polymorphic VNTR loci. These loci were identified with the program Tandem Repeat Finder (11). Four of the 5 loci had hexameric repeats in coding regions. The loci were identified by those genes in which they occur: VC0147vntr, VC0436–7vntr (intergenic), VC1650vntr, VC0171vntr, and VCA0283vntr. Alleles were distinguished by the number of tandem repeats as determined by Tandem Repeat Finder (11) (online Technical Appendix). Sequences from 1 locus with identical numbers of repeats were assigned to the identical allele. The alleles at the 5 loci were ordered to generate a sequence type (ST), for example, 3,5,2,2,8. Each locus was polymorphic with 7, 6, 6, 20, and 16 alleles, respectively. Thirty-six STs were observed. The various STs were defined as related if they were identical at 4 of the 5 loci. When we defined a VNTR genetic group as differing by a single locus variant from another member of the group, 3 large VNTR genetic groups were identified and 5 VNTR genetic groups composed of only 2 isolates and 7 unrelated strains. These 7 singletons differed from all other STs at 2 or more loci.

There was statistically significant agreement between serogroup and VNTR genetic group. For *V. cholerae* O139, all STs were 4,1,1,x,x (online Technical Appendix). Thus, the isolates were considered to be related because x,x = 1,1; 2,1; or 2,8, i.e., a change in a single locus serially connected all isolates. Summing the number of isolates of a sequence type, we found that the 23 ctx⁺ O139 strains formed a VNTR genetic group. A second group comprised 75 ctx⁺ O1 Inaba isolates. Finally, 18 ctx⁺ O1 Ogawa clustered into 3 additional VNTR genetic groups. There were 10 exceptions, i.e., 3 non-O1, non-O139 ctx⁺ isolates were in groups; 3 ctx⁻ O139, 2 ctx⁻ O1 Inaba, 1 ctx⁻, and 1 ctx⁺ O1 Ogawa were not.

We found that Bakerganj and Mathbaria yielded distinct *V. cholerae* populations; only 2 (ST 3,5,2,2,7 and 1,1,3,9,8) of 36 STs identified were found at both locations (Table 1; online Technical Appendix). There was substantial divergence in STs among strains isolated from patients, compared with strains from the environment in Mathbaria; only 1 (ST 3,5,2,2,7) of 16 STs were found in both patient and environmental isolates. Similarly, in Bakerganj, only 2 (ST 3,5,2,2,6 and 3,5,2,1,5) of 24 STs were found in both clinical and environmental isolates.

Clinical or environmental isolates from a given period were more likely to have a common ST (online Technical Appendix). For example, at Mathbaria, 49 of the 53 isolates with an ST identical to that of another isolate were found in the same or neighboring month. Similarly, at Bakerganj, 33 of 36 isolates with identical STs were found in the same or neighboring month.

Variation in the VNTR loci appeared to be greater among clinical isolates than among environmental isolates. A total of 29 STs occurred in clinical isolates, whereas only 12 occurred in environmental isolates (Table 1). When we controlled for location and month of collection (Table 2), the total number of STs among environmental isolates (7 ST/35 isolates) was less than that among clinical isolates (16 ST/32 isolates) ($\chi^2 = 4.4$, df 1, $p = 0.036$). Common STs were found among environmental isolates, despite the isolates coming from samples from different ponds and distinct subsamples (e.g., water, phytoplankton, zooplankton).

Table 1. Number of *Vibrio cholerae* sequence types in distinct serotypes and sample types in Bakerganj and Mathbaria, Bangladesh, 2004–2005

Serotype	Source	Bakerganj	Mathbaria
O1 Inaba	Clinic	10	7
	Environment	7	1
O1 Ogawa	Clinic	6	6
	Environment	1	0
O139	Environment	0	3

Conclusions

Our data do not support the concept of seasonal cholera epidemics occurring by movement of a single clonal wave across the countryside. They are consistent, instead, with the natural occurrence of *V. cholerae* year-round in the aquatic environment of each site, with each site having its own, distinct grouping of strains (12,13). The limited overlap between STs in environmental and clinical isolates is an enigma that remains to be resolved. However, the extensive variation in VNTR STs in this short time frame and small geographic area suggests that VNTR STs can be useful in assessing genetic relatedness of isolates during outbreaks/epidemics. The strong temporal clustering of the variation arising in the VNTR STs of clinical isolates is consistent with the hypothesis that clinical cases reflect the occurrence of multiple small outbreaks.

Our data are drawn from rural Bangladesh; however, cholera is a global disease. Its epidemiology may well differ in sub-Saharan Africa, the Americas, or other parts of Asia, or in the mega-cities that are increasingly the hallmark of the developing world. These variations emphasize the need for application of similar techniques in these diverse settings.

Our work was supported by a National Institutes of Health award to R.B.S.

Dr Stine is an associate professor of epidemiology and preventive medicine at the University of Maryland School of Medicine. He is actively using genetic variation in bacteria and humans to elucidate medical problems.

Table 2. Sequence type (ST) variations among *Vibrio cholerae* O1 Inaba isolates from environmental and clinical sources by month of collection, Bangladesh, 2004–2005


Location	Date	Source	No. ponds	No. isolates	No. STs	Variation*
Mathbaria	2004 Dec	Environment	4	12	1	0.08
Bakerganj	2004 Sep	Environment	5	16	4	0.25
Bakerganj	2005 Apr	Environment	4	7	2	0.29
Bakerganj	2004 Oct	Clinic		9	4	0.44
Mathbaria	2004 May	Clinic		11	5	0.45
Mathbaria	2004 Apr	Clinic		8	4	0.50
Bakerganj	2004 Dec	Clinic		4	3	0.75

*Variation, no. STs/no. isolates.

References

1. World Health Organization. Cholera 2005. *Wkly Epidemiol Rec*. 2006;81:297–308.
2. Huq A, Small EB, West PA, Huq MI, Rahman R, Colwell RR. Ecological relationships between *Vibrio cholerae* and planktonic crustacean copepods. *Appl Environ Microbiol*. 1983;45:275–83.
3. Brayton PR, Tamplin ML, Huq A, Colwell RR. Enumeration of *Vibrio cholerae* O1 in Bangladesh waters by fluorescent-antibody direct viable count. *Appl Environ Microbiol*. 1987;53:2862–5.
4. Franco AA, Fix AD, Prada A, Paredes E, Palomino JC, Wright AC, et al. Cholera in Lima, Peru, correlates with prior isolation of *Vibrio cholerae* from the environment. *Am J Epidemiol*. 1997;146:1067–75.
5. Tison DL, Kelly MT. *Vibrio* species of medical importance. *Diagn Microbiol Infect Dis*. 1984;2:263–76.
6. Alam M, Sultana M, Nair GB, Sack RB, Sack DA, Siddique AK, et al. Toxigenic *Vibrio cholerae* in the aquatic environment of Mathbaria, Bangladesh. *Appl Environ Microbiol*. 2006;72:2849–55.
7. Garg P, Aydanian A, Smith D, J Glenn M Jr, Nair GB, Stine OC. Molecular epidemiology of O139 *Vibrio cholerae*: mutation, lateral gene transfer, and founder flush. *Emerg Infect Dis*. 2003;9:810–4.
8. Ewing B, Green P. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res*. 1998;8:186–94.
9. Ewing B, Hillier L, Wendl MC, Green P. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res*. 1998;8:175–85.
10. Gordon D. Viewing and editing assembled sequences using Consed. In: Baxevanis A, Davison D, editors. *Current protocols in bioinformatics*. New York: John Wiley & Co; 2004. p. 11.2.1–11.2.43.
11. Benson G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res*. 1999;27:573–80.
12. Colwell RR. Global climate and infectious disease: the cholera paradigm. *Science*. 1996;274:2025–31.
13. Huq A, Sack RB, Nizam A, Longini IM, Nair GB, Ali A, et al. Critical factors influencing the occurrence of *Vibrio cholerae* in the environment of Bangladesh. *Appl Environ Microbiol*. 2005;71:4645–54.

Address for correspondence: O. Colin Stine, University of Maryland School of Medicine, 596 Howard Hall, 660 W Redwood St, Baltimore, MD 21201, USA; email: ostin001@umaryland.edu









Measurable Indicators


Search
past Issues

EID
Online
www.cdc.gov/eid

Distribution of sequence types by serotype and source in Bakerganj and Mathbaria. For each sequence type, the source is indicated by a C for clinical or an E for environmental, the number of isolates follows after n=, and the month(s) when it was isolated. The arrows indicate sequence types found at both locales, while the barbells indicate sequence types found in both clinical and environmental samples within a locale.

	Bakerganj	Mathbaria
O1 Inaba	 C 3,5,2,20,7 n=1 Apr-04	C 3,5,2,2,14 n=3 Apr 04
	C 3,5,2,1,7 n=1 Apr-04	C 2,5,2,2,6 n=6 Apr May 04
	C 3,5,2,2,6 n=2 Jun-04	C 5,5,2,2,6 n=3 Apr May 04
	E 3,4,2,1,12 n=13 Sep 04	C 3,5,2,2,16 n=4 Apr May 04
	E 3,5,2,15,12 n=1 Sep 04	C 3,5,2,2,7 n=1 May 04 
	E 3,5,2,20,12 n=1 Sep 04	C 2,5,2,18,6 n=1 May 04
	E 2,5,6,15,12 n=1 Sep 04	C 2,5,2,17,6 n=1 May 04
	C 3,5,2,1,6 n=5 Sep Oct Nov 04	E 3,5,2,2,7 n=12 Dec 04 
	C 3,5,2,1,8 n=1 Oct 04	
	C 3,5,2,1,5 n=5 Oct 04	
	C 5,5,2,16,5 n=1 Oct 04	
	E 3,5,2,1,5 n=2 Nov 04	
	C 4,5,2,3,6 n=2 Dec 04	
	C 3,5,2,3,6 n=1 Dec 04	
	C 3,5,2,3,7 n=1 Dec 04	
E 3,5,2,2,7 n=7 Feb-04 Apr 05 		
E 3,5,2,2,6 n=1 Apr 05		
O1 Ogawa	C 1,1,3,8,8 n=1 Jul 04	C 4,2,2,7,6 n=1 Apr 04
	C 1,1,3,9,8 n=4 Jul Aug Oct 04 	C 5,2,3,10,8 n=1 Apr 04
	C 6,1,3,8,8 n=2 Dec 04	C 1,1,3,9,8 n=1 Apr 04 
	C 2,1,3,7,8 n=1 Dec 04	C 2,1,4,10,4 n=1 Apr 04
	C 2,1,3,8,7 n=1 Dec 04	C 5,1,4,10,4 n=1 Apr 04
	C 5,1,3,8,8 n=1 Dec 04	C 1,1,3,6,7 n=2 Nov 04
	E 2,1,3,8,8 n=1 Dec 04 Apr 05	
O139	None	E 4,1,1,1,1 n=5 Mar Jul Sep 04
		E 4,1,1,2,1 n=17 Sep 04 Apr 05
		E 4,1,1,2,8 n=1 Apr 05

C: clinical isolate; **E**: environmental isolate

 : indicates genotypes are found in both clinical and environmental isolates

 : indicates genotypes found in both Bakerganj and Mathbaria