

3. Baseman JB, Tully JG. *Mycoplasma*: sophisticated, reemerging, and burdened by their notoriety. *Emerg Infect Dis*. 1997;3:21–32.
4. Baker AS, Ruoff KL, Madoff S. Isolation of *Mycoplasma* species from a patient with seal finger. *Clin Infect Dis*. 1998;27:1168–70. DOI: 10.1086/514980
5. Lierz M, Hagen N, Harcourt-Brown N, Hernandez-Divers SJ, Lueschow D, Hafez HM. Prevalence of mycoplasmas in eggs from birds of prey using culture and a genus-specific mycoplasma polymerase chain reaction. *Avian Pathol*. 2007;36:145–50. DOI: 10.1080/03079450701213347
6. Lierz M, Deppenmeier S, Gruber AD, Brokat S, Hafez HM. Pathogenicity of *Mycoplasma lipofaciens* strain ML64 for turkey embryos. *Avian Pathol*. 2007;36:389–93. DOI: 10.1080/03079450701589126
7. Bradbury JM, Forrest M, Williams A. *Mycoplasma lipofaciens*, a new species of avian origin. *Int J Syst Bacteriol*. 1983;33:329–35.
8. Bencina D, Dorrer D, Tadina T. *Mycoplasma* species isolated from six avian species. *Avian Pathol*. 1987;16:653–64. DOI: 10.1080/03079458708436413
9. Lierz M, Stark R, Brokat S, Hafez HM. Pathogenicity of *M. lipofaciens* strain ML64, isolated from an egg of a Northern Goshawk (*Accipiter gentilis*), for chicken embryos. *Avian Pathol*. 2007;36:151–3. DOI: 10.1080/03079450701213354
10. Christensen NH, Yavari CA, McBai AJ, Bradbury JM. Investigations into the survival of *Mycoplasma gallisepticum*, *Mycoplasma synoviae* and *Mycoplasma iowae* on materials found in the poultry house environment. *Avian Pathol*. 1994;23:127–43. DOI: 10.1080/03079459408418980

Address for correspondence: Michael Lierz, Institute for Poultry Diseases, Free University of Berlin, Königsweg 63, 14163 Berlin, Germany; email: lierz.michael@vetmed.fu-berlin.de

Urinary Tract Infection Caused by Capnophilic *Escherichia coli*

To the Editor: Increased atmospheric CO₂ concentrations promote the growth of fastidious microorganisms. However, the possibility that a strain of *Escherichia coli* can be CO₂ dependent is exceptional (1).

An isolate of capnophilic *E. coli* was responsible for a urinary tract infection (UTI) in a 77-year-old woman at the University Hospital of Guadalajara (Spain) in November 2002. Urine was cultured on a cystine-lactose-electrolyte-deficient agar plate and incubated at 37°C in an atmosphere containing 6% CO₂ for 1 day. After 24 hours, the culture yielded gram-negative rods (>10⁵ CFU/mL) in pure culture. The organism was motile, catalase positive, and oxidase negative. The strain could not be identified by using the MicroScan WalkAway-40 system (DadeBehring, Inc., West Sacramento, CA, USA). A subculture was performed, and the organism did not grow on sheep blood agar and MacConkey agar plates at 37°C in ambient air. However, a subculture incubated at 37°C for 24 hours in an atmosphere of 6% CO₂ produced smooth colonies 2–3 mm in diameter on sheep blood agar and MacConkey agar plates. The organism fermented lactose, and the indole reaction (BBL DrySlidet, Becton Dickinson Co., Sparks, MD, USA) performed on sheep blood agar was negative. The strain grew well on Schaedler agar plates after anaerobic incubation for 48 hours. The isolate remained capnophilic after 5 subcultures. The strain was identified as *E. coli* by using the Biolog GN2 panel (Biolog, Inc., Hayward, CA, USA) (100%, T = 0.534), after incubation of the panel in an atmosphere containing 6% CO₂ for 1 day. The API 20E system (bioMérieux, Marcy-l’Etoile, France) according to the manufac-

turer’s instructions without CO₂ incubation also identified *E. coli* (profile 5004512). The identification was confirmed by means of 16S rDNA sequence analysis (1,472 bp obtained by PCR amplification by a previously reported method [2]), which showed 99% similarity with *E. coli* sequence (GenBank accession no. CP000802). The 16S sequence showed similarity with *Shigella* species; however, this identification was not considered because the strain fermented lactose on MacConkey agar and agglutinations with *Shigella* antiserum were negative. The original 16S rDNA sequence was deposited in GenBank (accession no. EU555536).

The antimicrobial drug susceptibility profile was determined by incubating Mueller-Hinton agar plates at 37°C in an atmosphere containing 6% CO₂ by the disk diffusion method, according to National Committee for Clinical Laboratory Standards recommendations (3). The isolate was susceptible to ampicillin, amoxicillin/clavulanic acid, piperacillin, cefazolin, cefuroxime, cefotaxime, nitrofurantoin, fosfomicin, trimethoprim-sulfamethoxazole, gentamicin, tobramycin, amikacin, norfloxacin, and ciprofloxacin. MICs were obtained for the following antimicrobial agents with the E-test method (AB Biodisk, Solna, Sweden), performed on Mueller-Hinton agar plates incubated in a 6% CO₂ atmosphere: ampicillin (1.5 µg/mL), amoxicillin (3 µg/mL), cefotaxime (0.064 µg/mL), imipenem (0.094 µg/mL), piperacillin (2 µg/mL), and ciprofloxacin (0.008 µg/mL).

E. coli is the most common pathogen among patients with uncomplicated UTIs (4). Two cases of UTIs due to carbon dioxide-dependent strains of *E. coli* have been reported (1). The mechanisms for development of CO₂ dependence are unknown (5). CO₂ can play a role in the growth of *E. coli* as a substrate for carboxylation reactions (6). Other members of the family *En-*

**Search
past issues**

EID
Online
www.cdc.gov/eid

terobacteriaceae (such as some strains of *Klebsiella* spp.) and other organisms (such as *Staphylococcus aureus*), can have similar requirements (7,8).

There is not 1 best way of performing urine cultures. Guidelines for the diagnosis of UTI includes the use of sheep blood agar and either MacConkey agar or a similar selective medium for routine urine culture. The plates should be incubated overnight (at least 16 hours) at 37°C in ambient air; alternatively, the blood agar plate can be incubated in elevated (3%–8%) CO₂ (9). For fastidious microorganisms, chocolate agar can be added to the MacConkey agar and the plates incubated in 5% CO₂ for 2 days (9).

The real incidence of these infections is unknown, but the rarity of these strains suggests that the incidence is low. However, the real incidence of UTI caused by capnophilic *E. coli* may be underestimated because urine cultures are not usually incubated in CO₂. In addition, urine cultures are not performed for many women with uncomplicated cystitis. Other fastidious uropathogens such as *Haemophilus influenzae* and *H. parainfluenzae*, also require special media and incubation in an atmosphere of CO₂ (9). The low frequency of these strains suggests that incubation of routine urine cultures in an atmosphere containing CO₂ is not necessary. Incubation in CO₂ should be ordered only if the patient has pyuria and a previous negative urine culture after incubation in ambient air or if the patient is unresponsive to empiric therapy and routine urine culture is negative. Good clinician–laboratory communication is vital. Further studies should be performed to ascertain the real incidence of UTIs caused by capnophilic strains of *E. coli*.

Because no breakpoints are available for antimicrobial agents against capnophilic strains of *E. coli*, we used published interpretative criteria or *Enterobacteriaceae* (3). The strain was susceptible to all antimicrobial agents that we tested. The impact of CO₂ on

the susceptibility of capnophilic strains of *E. coli* is unknown. Susceptibility of some antimicrobial agents such as quinolones can be influenced by the pH change and enhanced growth that occur during CO₂ incubation when testing capnophilic organisms (10).

Daniel Tena,*
Alejandro González-Praetorius,*
Juan Antonio Sáez-Nieto,†
Sylvia Valdezate,†
and Julia Bisquert*

*University Hospital of Guadalajara, Guadalajara, Spain; and †Instituto de Salud Carlos III, Majadahonda, Madrid, Spain

DOI: 10.3201/eid1407.071053

References

1. Eykyn S, Phillips I. Carbon dioxide-dependent *Escherichia coli*. *BMJ*. 1978;1:576.
2. Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral JP, Raoult D. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacteria isolates. *J Clin Microbiol*. 2000;38:3623–30.
3. National Committee for Clinical Laboratory Standards. Performance standards for antimicrobial disk susceptibility tests. Approved standard M2–A8. Wayne (PA): The Committee; 2003.
4. Kahlmeter G. An international survey of the antimicrobial susceptibility of pathogens from uncomplicated urinary tract infection: the ECO.SENS Project. *J Antimicrob Chemother*. 2003;51:69–76. DOI: 10.1093/jac/dkg028
5. Repaske R, Clayton MA. Control of *Escherichia coli* growth by CO₂. *J Bacteriol*. 1978;135:1162–4.
6. Kozliak EI, Fuchs JA, Guilloton MB, Anderson PM. Role of bicarbonate/CO₂ in the inhibition of *Escherichia coli* growth by cyanate. *J Bacteriol*. 1995;177:3213–9.
7. Barker J, Brookes G, Johnson T. Carbon dioxide-dependent *Klebsiellae*. *BMJ*. 1978;1:300.
8. Rahman M. Carbon dioxide-dependent *Staphylococcus aureus* from abscess. *BMJ*. 1977;2:319.
9. Clarridge JE, Johnson JR, Pezzlo MT. Laboratory diagnosis of urinary tract infections. In: Weissfeld AS, editor. *Cumitech 2B*. Washington: ASM Press; 1988. p. 2–19.
10. Bolmstrom A, Karlsson A. Influence of CO₂ incubation on quinolone activity against *Streptococcus pneumoniae* and *Haemophilus influenzae*. *Diagn Microbiol*

Infect Dis. 2002;42:65–9. DOI: 10.1016/S0732-8893(01)00338-8

Address for correspondence: Daniel Tena, Section of Microbiology, University Hospital of Guadalajara, Calle Donantes de Sangre s/n 19002 Guadalajara, Spain; email: daniel@seccam.jccm.es

Duck Migration and Past Influenza A (H5N1) Outbreak Areas

To the Editor: In 2005 and 2006, the highly pathogenic avian influenza (HPAI) virus subtype H5N1 rapidly spread from Asia through Europe, the Middle East, and Africa. Waterbirds are considered the natural reservoir of low pathogenic avian influenza viruses (1), but their potential role in the spread of HPAI (H5N1), along with legal and illegal poultry and wildlife trade (2), is yet to be clarified.

The garganey (*Anas querquedula*) is the most numerous duck migrating between Eurasia and Africa: ≈2 million gather in the wetlands of Western Africa every northern winter (3). We report on a spatial correlation between the 2007 migration path of a garganey monitored through satellite telemetry and areas that had major HPAI (H5N1) outbreaks from 2005 through 2007.

Seven garganeys were captured, sampled, and fitted with a 12-g satellite transmitter in northern Nigeria (Hadejia-Nguru Wetlands; 12°48'N; 10°44'E) in the period February 7–15, 2007. All cloacal and tracheal swabs tested negative for avian influenza virus by real-time reverse transcription–PCR analysis of the matrix gene. One second-year (>9-month-