

Nine patients originated from Somalia, Ethiopia, or Yemen. Few cases have previously been reported from this area (3), although Ethiopian migrants have been shown to have an egg positivity of 0.4% on routine screening (4). Patients 5 and 6 had not returned to Africa for >20 years, suggesting that they acquired their infection in Europe. Therefore, a risk factor may exist that is specific to this ethnic group within the United Kingdom.

Six cases were diagnosed at 1 hospital. All 6 patients reported current or past use of locally bought *khat*, a leaf chewed for its stimulant properties. It is imported fresh to the United Kingdom from Africa and is an ideal environment for the survival of *Fasciola cercariae*. It is used most commonly by migrants from the Horn of Africa and Yemen and has been reported in association with acute fascioliasis in the United Kingdom (5). Use of imported *khat* may explain the apparently higher incidence of fascioliasis in this ethnic group residing in the United Kingdom.

Despite the described parallel rise in human and veterinary fascioliasis, none of these cases provide clear evidence that recent human cases resulted from zoonotic transmission within the United Kingdom. Most cases occurred in migrants from the Horn of Africa and Yemen, some of whom may have acquired *Fasciola* spp. in their country of origin; other cases appear likely to have been acquired in the United Kingdom, possibly due to use of imported *khat*. Physicians need a heightened awareness of fascioliasis when investigating impaired liver function or abnormal abdominal imaging in migrants or travelers from high-risk areas.

**Meera A. Chand,
Joanna S. Herman,
David G. Partridge,
Kirsten Hewitt,
and Peter L. Chiodini**


Author affiliations: Hospital for Tropical Diseases, London, UK (M.A. Chand, J.S. Herman, P.L. Chiodini); Royal Hallamshire Hospital, Sheffield, UK (D.G. Partridge); Health Protection Agency Centre for Infections, London (K. Hewitt); and London School of Hygiene and Tropical Medicine, London (P.L. Chiodini)

DOI: 10.3201/eid1511.090511

References

1. Marcos LA, Terashima A, Gotuzzo E. Update on hepatobiliary flukes: fascioliasis, opisthorchiasis and clonorchiasis. *Curr Opin Infect Dis*. 2008;21:523–30.
2. Marcos LA, Tagle M, Terashima A, Busalieu A, Ramirez C, Carrasco C, et al. Natural history, clinoradiologic correlates and response to triclabendazole in acute massive fascioliasis. *Am J Trop Med Hyg*. 2008;78:222–7.
3. Control of foodborne trematode infections. Report of a WHO study group. *World Health Organ Tech Rep Ser*. 1995;849:1–157.
4. Nahmias J, Greenberg Z, Djerrasi L, Giladi L. Mass treatment of intestinal parasites among Ethiopian immigrants. *Isr J Med Sci*. 1991;27:278–83.
5. Doherty JF, Price N, Moody AH, Wright SG, Glynn MJ. Fascioliasis due to imported *khat*. *Lancet*. 1995;345:462. DOI: 10.1016/S0140-676(95)90450-6

Address for correspondence: Meera A. Chand, Hospital for Tropical Diseases, Mortimer Market, Capper Street, London WC1E 6JB, UK; email: meera.chand@nhs.net



Order your own copy of CDC's trusted travel medicine desktop reference:
CDC Health Information for International Travel 2010
www.us.elsevierhealth.com

Gastroenteritis Outbreaks in 2 Tourist Resorts, Dominican Republic

To the Editor: Noroviruses are an important cause of acute gastroenteritis, and outbreaks caused by these viruses have emerged as a major challenge to the healthcare, leisure, and tourism industries. The primary reason is their highly efficient transmission among persons in semiclosed populations such as those in healthcare facilities, hotels, and cruise ships. During an outbreak, primary cases result from exposure to a fecally contaminated vehicle (e.g., food or water), whereas secondary and tertiary cases among contacts of primary case-patients result from person-to-person transmission (1). Airborne and fomite transmission also play a role in the virus spread during outbreaks (2). Transmission through recreational water has also been described (3).

We investigated 2 outbreaks of norovirus gastroenteritis in tourist resorts in the Dominican Republic in January 2005. A total of 402 persons and 371 persons at 2 resorts, 1 located in Punta Cana (attack rate 6.8%) and another in Puerto Plata (attack rate 6.2%), respectively, reported symptoms of diarrhea, vomiting, headache, and fatigue. A total of 35 stools samples, 28 from Punta Cana and 7 from Puerto Plata, were negative for bacterial or parasitic pathogens. However, norovirus was confirmed by the IDEIA norovirus immunoassay (DakoCytomation, Ely, UK) in 11 samples from Punta Cana and 7 samples from Puerto Plata.

Active measures to reduce norovirus transmission were adopted by the 2 resorts, including an increase in cleaning frequency and increase in concentration of chlorine used for surface disinfection of public areas (1,000 mg/L), kitchenware (200 mg/L for 15 min), and fruits and vegetables (150 mg/L for 15 min). Personnel involved

in food preparation were invited to fill out an epidemiologic questionnaire and provide stool samples.

Because these preventive measures were not effective, water was suspected as a possible route of transmission, and diverse approaches were used to isolate viral RNA from different types of water. For sewage water, 100 mL were concentrated by a Molecular Weight Cutoff (MWCO) 100,000 Da Vivaspin20 centrifugal concentrator (Sartorius, Madrid, Spain) to a 5-mL concentrated sample. Five milliliters of chloroform:isoamyl alcohol (24:1) was added to the sample, mixed in a vortex, and centrifuged at $1,000 \times g$ for 10 min (4). A second concentration step to 1 mL was performed with the resulting aqueous phase using a MWCO 100,000 Da Vivaspin6 centrifugal concentrator (Sartorius). For tap and recreational water, a sample of 2 L was processed through a cellulose nitrate filter, as described (5). The filter was first activated by 5 mL of 0.25 mol/L AlCl_3 . After a washing step with 200 mL 0.5 mmol/L H_2SO_4 , viruses were eluted from the filter with 10 mL of 1 mmol/L NaOH and neutralized with 50 μL of 0.1 mol/L H_2SO_4 and 1.1 mL $10\times$ pH 8.0 Tris-EDTA buffer. Finally, samples were concentrated to a final volume of 1 mL with a Vivaspin6, centrifugal concentrator, as described above.

Viral RNA was isolated by the QIAamp Viral RNA Mini Kit (QIAGEN, Hilden, Germany), treated with Dnase I (Invitrogen, Carlsbad, CA, USA), and reverse transcribed to cDNA with Superscript II reverse transcriptase (Invitrogen). Finally, cDNA was amplified by multiplex-PCR with Taq Gold polymerase (Applied Biosystems, Foster City, CA, USA) by using forward primers Nor31N (5'-CAGATTAYACWGCWTGGGA-3') and Nor32M (5'-CAGATTAYTCWCGWTGGGA-3'); and reverse primers Nor41M (5'-CCARTGATTTATGCTG TTCAC-3') and Nor42N (5'-CCAGT

GGCGATGGAGTTC-3'), specific for the RNA polymerase gene (provided by J.A. Boga and M. Oña) (6). PCR was performed with the following conditions: 12 min at 94°C, (1 min at 94°C, 1 min at 55°C, 1 min at 72°C) for 40 cycles, and 10 min at 72°C. PCR products were analyzed by agarose gel electrophoresis. A band of 221 bp was considered a positive result for norovirus.

Norovirus was detected in the 4 samples of sewage water analyzed (2 from each location) collected after intervention, indicating that viral carriers still remained in the resorts. Moreover, norovirus particles were detected in the 2 deputed water samples (1 from each area), then indicating insufficient treatment conditions. Further investigation showed that deputed water was being used to water plants and grass by sprinkling, and becoming an important secondary source for infection. Norovirus-contaminated sewage was treated with different concentrations of chlorine and reverse transcription-PCR demonstrated that 15 mg/L for 1 h did not give a positive signal. Garden sprinkling watering was replaced by inundation watering. Although the virus was not detected either in tap or recreational water, hyperchlorination was carried out to prevent possible dissemination of norovirus. Preventive measures described above for surfaces, kitchenware and food were maintained for 2 additional weeks, and no additional cases were detected after 1 month had passed since the first case. To summarize, only interventions at multiple points, including eradication of secondary sources, and preventive measures to avoid person-to-person transmission enabled the outbreaks to be controlled.

**Antonio Doménech-Sánchez,
Carlos Juan, Antonio J. Rullán,
José L. Pérez,
and Clara I. Berrocal**

Author affiliations: Saniconsult Ibérica SL, Palma de Mallorca, Spain (A. Doménech-Sánchez, C.I. Berrocal); Instituto Universitario de Investigación en Ciencias de la Salud, Palma de Mallorca (A. Doménech-Sánchez, C. Juan, J.L. Pérez); Hospital Son Dureta. Palma de Mallorca (C. Juan, J.L. Pérez); and Servicio Navarro de Salud, Pamplona, Spain (A.J. Rullán)

DOI: 10.3201/eid1511.090350

References

1. Parashar U, Quiroz ES, Mounts AW, Monroe SS, Fankhauser RL, Ando T, et al. Norwalk-like viruses. Public health consequences and outbreak management. *MMWR Recomm Rep*. 2001;50(RR-9):1-17.
2. Marks PJ, Vipond IB, Carlisle D, Deakin D, Fey RE, Caul EO. Evidence for airborne transmission of Norwalk-like virus (NLV) in a hotel restaurant. *Epidemiol Infect*. 2000;124:481-7. DOI: 10.1017/S0950268899003805
3. Doménech-Sánchez A, Olea F and Berrocal CI. Infections related to recreational waters [in Spanish]. *Enferm Infecc Microbiol Clin* 2008;26 (Suppl 4):33-8
4. Schwab KJ, De Leon R, Sobsey MD. Concentration and purification of beef extract mock eluates from water samples for the detection of enteroviruses, hepatitis A virus, and Norwalk virus by reverse transcription-PCR. *Appl Environ Microbiol*. 1995;61:531-7.
5. Katayama H, Shimasaki A, Ohgaki S. Development of a virus concentration method and its application to detection of enterovirus and Norwalk virus from coastal seawater. *Appl Environ Microbiol*. 2002;68:1033-9. DOI: 10.1128/AEM.68.3.1033-1039.2002
6. Boga JA, Ordás J, Melón S, Villar M, González D, Temprano MA et al. Detection, genotyping and temporal distribution of norovirus producing sporadic cases and epidemic outbreaks of gastroenteritis in Asturias [in Spanish]. *Enferm Infecc Microbiol Clin* 2004;22(Suppl 1): 186.

Address for correspondence: Antonio Doménech-Sánchez, Saniconsult Ibérica SL, Can Foradí 37, Bajos Son Cladera Nou, E 07009 Palma de Mallorca, Spain; email: adomenech@saniconsult.net

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the U.S. Department of Health and Human Services.