
Evidence-based Tool for Triggering School Closures during Influenza Outbreaks, Japan

Asami Sasaki, Anne Gatewood Hoen, Al Ozonoff, Hiroshi Suzuki, Naohito Tanabe, Nao Seki, Reiko Saito, and John S. Brownstein

Guidelines available to school administrators to support school closure decisions during influenza outbreaks are usually not evidence based. Using empirical data on absentee rates of elementary school students in Japan, we developed a simple and practical algorithm for determining the optimal timing of school closures for control of influenza outbreaks.

Influenza pandemic preparedness and seasonal influenza control programs have focused on vaccine development and antiviral drugs, which are only partially effective and not always available to all persons at risk (1–3). Nonpharmaceutical interventions, such as social distancing, represent additional key tools for mitigating the impact of outbreaks. Because children are a major factor in the transmission of influenza within communities and among households, school closure may be a valuable social distancing method (4,5).

Japan has a unique system of monitoring school absenteeism and of instituting school closures during influenza outbreaks. Individual classes, specific grade levels, or the entire school may be closed; final decision-making authority is given to school principals. However, as in the United States and other countries, there are no regulations to support these decisions (6). Our study suggests a simple system to help determine when schools should be closed; daily influenza-related absentee thresholds are measured to predict outbreaks.

Author affiliations: Harvard School of Public Health, Boston, Massachusetts, USA (A. Sasaki); University of Niigata Prefecture, Niigata, Japan (A. Sasaki); Children's Hospital, Boston (A. Gatewood Hoen, J.S. Brownstein); Harvard Medical School, Boston (A. Gatewood Hoen, J.S. Brownstein); Boston University School of Public Health, Boston (A. Ozonoff); and Niigata University Graduate School of Medical and Dental Sciences, Niigata (H. Suzuki, N. Tanabe, N. Seki, R. Saito)

DOI: 10.3201/eid1511.090798

The Study

We used data on absenteeism caused by influenza from the 54 elementary schools in Joetsu City, Niigata Prefecture, Japan during the 4 influenza seasons during 2005–2008. Data was obtained between the second week of January to the third week of March for each influenza season. Average school size was 221 students. Current public health policy prevents influenza-infected children from attending school until 2 days after fever has disappeared. An illness requires 2 physician visits: 1 for the initial diagnosis and 1 to obtain written permission from the treating physician to return to school. Diagnoses are usually made by using a rapid antigen test and patients are treated with the antiviral drugs, oseltamivir or zanamivir.

Based on elementary school daily influenza-related absentee surveillance, the most intense influenza seasons were 2005 and 2007 (Figure 1). The number of schools reporting outbreaks during the 4 influenza seasons was 34 (63%, 2005), 13 (24%, 2006), 35 (65%, 2007) and 18 (33%, 2008), respectively. Rates of absenteeism caused by confirmed influenza infection in the 54 elementary schools in Joetsu City were well correlated with national reports of influenza-like illness by 5,000 sentinel physicians, who reported 322, 205, 226, and 142 cumulative cases of infection per sentinel in each season (online Technical Appendix, available from www.cdc.gov/EID/content/15/11/1841-Techapp.pdf).

We evaluated the optimal influenza-related absentee rate for predicting outbreaks of influenza. For this study, we defined an influenza outbreak in a school as a daily influenza-related absentee rate of >10%, on the basis of the 95th percentile of daily absentee rates (10.7%) in 54 elementary schools during 4 influenza seasons (online Technical Appendix).

Next, we considered 9 different daily influenza-related absentee threshold levels for initiating early school closures: 1%, 2%, 3%, ..., 9%. In addition, for each threshold level, we considered 3 scenarios: 1) a single-day scenario, in which daily influenza-related absentee rates are observed for the first time above a given threshold for 1 day; 2) a double-day scenario, in which rates reached a given threshold for the first time for 2 consecutive days; the rate for the second day was the same or higher than for the first day; and 3) a triple-day scenario, in which rates reached a given threshold for the first time for 3 consecutive days; rates for the second and third days were the same or higher than the rate for the first day. The double-day and triple-day scenarios did not include weekends. To evaluate the performance of prediction for each threshold, we determined the school's outbreak status in the 7-day period starting on the first day of each scenario (online Technical Appendix) JMP7.0.1 (SAS Institute, Inc., Cary, NC, USA) was used for statistical analysis.




Figure 1. Four-year surveillance of influenza-related absentee rates in 54 elementary schools in Joetsu City and national surveillance of influenza-like illness (ILI) reported by sentinel physicians in Japan. Data were collected from the second week of January (after the winter holiday) to the third week of March (before the spring holiday). The average of the daily absentee rates for 54 elementary schools during 4 influenza seasons (2005–2008) were 3.29%, 1.77%, 2.97%, and 1.92%, respectively.

We calculated the sensitivity and specificity of each scenario at all 9 threshold levels, and presented these data as a plot in Figure 2. The area under the curve for the single-, double-, and triple-day scenarios was 0.80 (95% confidence interval [CI] 0.77–0.83), 0.85 (95% CI 0.82–0.89) and 0.87 (95% CI 0.83–0.91), respectively.

We used the Youden index for calculating optimal thresholds (7). The Youden index = (sensitivity) + (specificity) – 1. A perfect test result would have a Youden index of 1. For the single-day scenario, the optimal threshold was 5%, with a sensitivity of 0.77 and specificity of 0.73. For the double-day scenario, the optimal threshold was 4%, with a sensitivity of 0.84 and specificity of 0.77. For the triple-day scenario, the optimal threshold was 3%, with a sensitivity of 0.90 and specificity of 0.72.

Conclusions

We have demonstrated the predictive value of a simple and practical detection method for triggering school closures early after influenza outbreaks. Our analysis suggests that a single-day at a threshold influenza-related absentee rate of 5%, double-days $\geq 4\%$, or triple-days $\geq 3\%$ are optimal levels for alerting school administrators to consider school closure. The double- and triple-day scenarios performed similarly, and gave better results than the single-day. Thus, the double-day scenario might be the preferred early warning trigger.

Our study had the advantage of reliable empirical data on influenza-related absenteeism in schools. Data were based on physician and laboratory diagnosis and a strong absentee surveillance program. However, there are limitations to our approach. We did not have available vaccina-

tion or medication histories of patients. Also, our results are based on data from only 1 city's school district; validation in a broader area will be required. Although separate analyses may be required for other geographic regions, we present a simple approach that can be easily reapplied.

Influenza outbreak detection from surveillance data typically relies on relatively complex time series analysis or smoothing (8,9). The noisiness of school surveillance data makes detection of outbreaks difficult (10). However, complex statistical analyses are not practical to use in the context of daily decision-making in schools. Despite the limitations of our study, we have presented a method that provides a basis for empirical data-supported decision-making by school administrators that is intuitive and practical.

School closure could be an effective method of social distancing, although evidence supporting its effectiveness is incomplete. Some studies suggest that though child-to-child transmission might decrease, transmission might increase in other age groups (11,12). During school closures, children may need to forgo participation in external activities that could increase contact rates. Additionally, working parents staying home to care for their children (13) could result in a decrease in household income, causing loss of productivity and economic losses (14). Decision-makers will need to consider these factors when considering school closures.




Figure 2. The receiver operating characteristic (ROC) curve for detection of influenza outbreak by 1%–9% thresholds under single-day, double-day, triple-day scenarios. ROC space is defined on the x axis as specificity and on the y axis as sensitivity. The area under the curve (AUC) is an indicator of the quality of a model; larger AUC values corresponded to better performance. Optimal thresholds for the 3 scenarios are *single-day, 5%; †double-day, 4%; and ‡triple-day, 3%.

During the early days of the outbreak of influenza A pandemic (H1N1) 2009 virus, the US Centers for Disease Control and Prevention (Atlanta, GA, USA) released 2 different recommendations for school dismissal after the appearance of the first suspected case: dismiss for 7 days (as of April 26) and then for 14 days (as of May 1). Later, to reflect new knowledge about the extent of community spread and disease severity, the recommendation was revised to advise against school closure unless absentee rates interfered with school function (15). The pandemic (H1N1) 2009 influenza outbreak highlights the need for a flexible national policy that can be quickly adapted to reflect current situations. The evidence-based strategy for predicting outbreaks based on influenza-related absentee rates that we present here provides local administrators, who may need to consider school closure, with a simple and practical tool to aid in their decisions.

Acknowledgments

We thank the Education Board of Joetsu City, Niigata Prefecture, Japan for providing school surveillance data.

This work was supported by Takemi Program, which is funded by the Japan Foundation for the Promotion of International Medical Research Cooperation, the National Institute of Allergy and Infectious Disease, the National Institutes of Health Research, and the Canadian Institutes of Health Research. A.S. is a recipient of a fellowship grant in the 2008–09 Takemi Program in the Department of Global Health and Population at the Harvard School of Public Health.

Dr Sasaki is an associate professor at the Department of Health and Nutrition, University of Niigata Prefecture. Her research focuses on the epidemiology of infectious diseases and health surveillance systems.

References

- Davey VJ, Glass RJ, Min HJ, Beyeler WE, Glass LM. Effective, robust design of community mitigation for pandemic influenza: a systematic examination of proposed US guidance. *PLoS One*. 2008;3:e2606. DOI: 10.1371/journal.pone.0002606
- Glezen WP. Clinical practice. Prevention and treatment of seasonal influenza. *N Engl J Med*. 2008;359:2579–85. DOI: 10.1056/NEJMc0807498
- Lipsitch M, Cohen T, Murray M, Levin BR. Antiviral resistance and the control of pandemic influenza. *PLoS Med*. 2007;4:e15. DOI: 10.1371/journal.pmed.0040015
- Cauchemez S, Valleron AJ, Boelle PY, Flahault A, Ferguson NM. Estimating the impact of school closure on influenza transmission from Sentinel data. *Nature*. 2008;452:750–4. DOI: 10.1038/nature06732
- Heymann A, Chodick G, Reichman B, Kokia E, Laufer J. Influence of school closure on the incidence of viral respiratory diseases among children and on health care utilization. *Pediatr Infect Dis J*. 2004;23:675–7. DOI: 10.1097/01.inf.0000128778.54105.06
- Kahn LH. Pandemic influenza school closure policies. *Emerg Infect Dis*. 2007;13:344–5. DOI: 10.3201/eid1302.061109
- Fluss R, Faraggi D, Reiser B. Estimation of the Youden index and its associated cutoff point. *Biom J*. 2005;47:458–72. DOI: 10.1002/bimj.200410135
- Cowling BJ, Wong IO, Ho LM, Riley S, Leung GM. Methods for monitoring influenza surveillance data. *Int J Epidemiol*. 2006;35:1314–21. DOI: 10.1093/ije/dyl162
- Gault G, Larrieu S, Durand C, Josseran L, Jouvès B, Filleul L. Performance of a syndromic system for influenza based on the activity of general practitioners, France. *J Public Health (Oxf)*. 2009;31:286–92. DOI: 10.1093/pubmed/udp020
- Besculides M, Heffernan R, Mostashari F, Weiss D. Evaluation of school absenteeism data for early outbreak detection, New York City. *BMC Public Health*. 2005;5:105. DOI: 10.1186/1471-2458-5-105
- Mikolajczyk RT, Akmatov MK, Rastin S, Kretzschmar M. Social contacts of school children and the transmission of respiratory-spread pathogens. *Epidemiol Infect*. 2008;136:813–22. DOI: 10.1017/S0950268807009181
- Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med*. 2008;5:e74. DOI: 10.1371/journal.pmed.0050074
- Johnson AJ, Moore ZS, Edelson PJ, Kinnane L, Davies M, Shay DK, et al. Household responses to school closure resulting from outbreak of influenza B, North Carolina. *Emerg Infect Dis*. 2008;14:1024–30. DOI: 10.3201/eid1407.080096
- Sadique MZ, Adams EJ, Edmunds WJ. Estimating the costs of school closure for mitigating an influenza pandemic. *BMC Public Health*. 2008;8:135. DOI: 10.1186/1471-2458-8-135
- Centers for Disease Control and Prevention. Update on school (K–12) and child care programs: interim CDC guidance in response to human infections with the novel influenza A (H1N1) virus. 2009 May 09 [cited 2009 May 27]. Available from http://www.cdc.gov/h1n1flu/K12_dismissal.htm

Address for correspondence: Asami Sasaki, Division of Health and Nutrition, 471 Ebigase, Higashi-Ku, Nigata City, Nigata 950-8680, Japan; email: asammy@unii.ac.jp




Bonus article content available on our podcast page.
<http://www.cdc.gov/ncidod/eid/podcast/index.htm>

New bonus content available for select articles each month.
ONLINE ONLY

Evidence-based Tool for Triggering School Closures during Influenza Outbreaks, Japan

Technical Appendix



Technical Appendix Figure 1. Plot of weekly absentee rates of influenza cases from 54 elementary schools vs. weekly national influenza-like-illness (ILI) cases reported by Sentinel physicians, 2005–2008. * $p < 0.01$.

Relation between national surveillance with +1 week lag and school surveillance

Correlations				
			sentinel	schoolaverag erate
Spearman's rho	sentinel	Correlation Coefficient	1.000	.753**
		Sig. (2-tailed)		.000
		N	40	40
	schoolaverag erate	Correlation Coefficient	.753**	1.000
		Sig. (2-tailed)	.000	
		N	40	40

** . Correlation is significant at the 0.01 level (2-tailed).

Relation between school surveillance with+ 1 week lag and national surveillance

Correlations				
			sentinel	schoolaverag erate
Spearman's rho	sentinel	Correlation Coefficient	1.000	.806**
		Sig. (2-tailed)		.000
		N	39	39
	schoolaverag erate	Correlation Coefficient	.806**	1.000
		Sig. (2-tailed)	.000	
		N	39	39

** . Correlation is significant at the 0.01 level (2-tailed).

Relation between national surveillance with+2 week lag and school surveillance

Correlations				
			sentinel	schoolaverag erate
Spearman's rho	sentinel	Correlation Coefficient	1.000	.587**
		Sig. (2-tailed)		.000
		N	35	35
	schoolaverag erate	Correlation Coefficient	.587**	1.000
		Sig. (2-tailed)	.000	
		N	35	35


** . Correlation is significant at the 0.01 level (2-tailed).

Relation between school surveillance with+2 week lag and national surveillance

Correlations				
			sentinel	schoolaverag erate
Spearman's rho	sentinel	Correlation Coefficient	1.000	.452**
		Sig. (2-tailed)		.003
		N	41	41
	schoolaverag erate	Correlation Coefficient	.452**	1.000
		Sig. (2-tailed)	.003	
		N	41	41

** . Correlation is significant at the 0.01 level (2-tailed).


Technical Appendix Figure 2. Correlation between lagged weekly rates of absenteeism due to confirmed influenza cases from 54 elementary schools and weekly national influenza-like-illness cases reported by Sentinel physicians, 2005–2008. None of the lagged comparisons resulted in an improved correlation over the unlagged relationship shown in Technical Appendix Figure 1.



Technical Appendix Figure 3. Histogram of daily rate of absenteeism related to confirmed influenza cases in 54 elementary schools. We defined an influenza outbreak in a school as a daily influenza-related absentee rate of >10%, on the basis of the 95th percentile of daily absentee rates (10.7%) for 4 influenza seasons.


Single, Double and Triple-day scenario and outbreak(10%) status

Ex) School 1



(A) **Single-day scenario 2%** (First time reaches 2% for only one day)


Is there 10% in 7 days? **NO**



(B) **Double-day scenario 3%**

(First time reaches 3% with second day same or higher rate)


Is there 10% in 7 days? **YES**



(C) **Triple-day scenario 2%**

(First time reaches 2% with second and third day same or higher rate)

Is there 10% in 7 days? **YES**



Technical Appendix Figure 4. Schematic illustration of our method for evaluating and optimizing our algorithm. For each of the 54 elementary schools, and for each influenza season of the study, we considered 3 scenarios: a single-day scenario, in which daily influenza-related absentee rates are observed for the first time above a given threshold for 1 day; a double-day scenario, in which rates reached a given threshold for the first time for 2 consecutive days, with the second day at the same rate or higher than the first; and a triple-day scenario, in which rates reached a given threshold for the first time for 3 consecutive days, with the second and third days at the same rate or higher than the first. Each scenario was evaluated at 9 different absentee threshold points: 1%, 2% ... 9%. The example illustrated above shows how we evaluated the algorithm at 1 school during 1 influenza season under 3 arbitrarily chosen scenario-threshold combinations. A) For the single-day scenario evaluated at the 2% threshold, we calculated the date that absenteeism due to confirmed influenza reached at least 2% and noted whether the outbreak threshold of 10% was reached in the following 7 days. B) For the double-day scenario evaluated at the 3% threshold level, we calculated the date that absenteeism due to confirmed influenza reached at least 3% and was sustained at $\geq 3\%$ for at least 2 consecutive days (excluding weekends), and then noted whether the outbreak threshold of 10% was reached within the 7 days after the first day. C) For the triple-day scenario evaluated at the 2% threshold level, we calculated the date that absenteeism due to confirmed influenza reached at least 2% and was sustained at $\geq 2\%$ for at least 3 consecutive days (excluding weekends), and then noted whether the outbreak threshold of 10% was reached in the 7 days after the first day.