

Appendix Table. *Mycobacterium* spp. implicated in lymphadenopathy in children, France*

<i>Mycobacterium</i> spp.	Clinical features			Microbiologic tests			Histopathology				Reference
	No. patients	Treatment	Outcome	Direct examination†	Culture	Molecular identification	Ring of epithelial cells	Giant cells	Caseous necrosis	Altered leukocytes	
<i>M. tuberculosis</i>				±	+	IS6110	+	+	+	–	(6)
<i>M. avium</i>	183	NR	NR	NR	+	NR	+	+	NR	NR	(7)
<i>M. avium</i> subsp. <i>hominissuis</i>	34	NR	NR	NR	+	IS1245	NR	NR	NR	NR	(5)
<i>M. haemophilum</i>	39	NR	NR	+	+	16S–23S rDNA	+	+	+	–	(8)
<i>M. lentiflavum</i>	3	SE	Cured	+	+	16S rRNA	+	+	NR	+	(9)
<i>M. bohemicum</i>	4	SE/C, R	Cured	±	+	16S rRNA	+	+	+	–	(10)
<i>M. simiae</i>	1	SE	Cured	+	+	HPLC	+	+	+	–	(3)
<i>M. colombiense</i>	1	SE	Cured	+	+	16S rRNA	+	+	+	–	This study
	1	SE	Cured	–	+	<i>rpoB</i>	+	+	+	–	

*IS, insertion sequence; NR, not reported; SE, surgical excision; C, clarithromycin; R, rifampin; NR, not reported; HPLC, high-performance liquid chromatography.

†Ziehl-Nelsen staining.