

4. Butler WR, Floyd MM, Brown J, Toney SR, Daneshvar M, Cooksey RC, et al. Novel mycolic acid-containing bacteria in the family *Segniliparaceae* fam. nov., including the genus *Segniliparus* gen. nov., with descriptions of *Segniliparus rotundus* sp. nov., and *Segniliparus rugosus* sp. nov. *Int J Syst Evol Microbiol*. 2005;55:1615–24.
5. Brown-Elliott B, Beierle K, McGlasson M, Wallace R. MICs of non-tuberculous mycobacteria using the TREK Vizion system compared to manual readings. In: 18th European Congress of Clinical Microbiology and Infectious Diseases; Barcelona, Spain; 2008 April 19–22. Abstract R2410.
6. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial testing; sixteenth informational supplement. CLSI14 document M100–S16. Wayne (PA): The Institute; 2006.
7. National Committee on Clinical and Laboratory Standards (NCCLS). Susceptibility testing of mycobacteria, nocardiae, and other aerobic actinomycetes; approved standard. NCCLS document M24–A. Wayne (PA): The Committee; 2003.
8. Butler WR, Guthertz LS. Mycolic acid analysis by high performance liquid chromatography for identification of mycobacterium species. *Clin Microbiol Rev*. 2001;14:704–26. DOI: 10.1128/CMR.14.4.704-726.2001

Address for correspondence: Tarrant Hansen, Pathology Queensland, Mycobacterium Reference Laboratory, Brisbane, Queensland, Australia; email: tarrant_hansen@health.qld.gov.au

Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article's publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have 1 Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.

Multigenotype Q Fever Outbreak, the Netherlands

To the Editor: Q fever is a zoonosis caused by *Coxiella burnetii* (*I*). An ongoing Q fever outbreak has occurred in the Netherlands since 2007; incidence rates have increased >50-fold compared with the baseline rate (2). The source of this outbreak is unknown. Identifying the source of an infection is complicated because of difficulties in obtaining sufficient clinical and/or environmental samples for testing. Molecular diagnosis of Q fever has focused on the use of serum samples. Up-to-date genotyping of *C. burnetii* has depended on cultivation and enrichment of the isolate before analysis (3). We report multiple-locus variable-number tandem repeat analysis (MLVA) typing of *C. burnetii* for a variety of human and animal clinical samples obtained from different locations in the Netherlands (Table).

Severe pneumonia developed in patient 1 after close contact with sheep (ewes) and intimate cuddling with a newborn lamb. Patients 2 and 3 (a dairy goat farmer and his wife from another village) tested positive for Q fever after a large part of their goat herd aborted offspring. The farmer had no clinical symptoms; his wife had mild symptoms that disappeared spontaneously within 2 days. No samples from any of the goats were available. Two additional patients were tested, 1 of which lived in the same village as patients 2 and 3.

Swab specimens from all sheep and lambs tested in the first case yielded identical MLVA genotypes. The same genotype was also found in patient 1 but not in the other examined samples, implicating sheep as the origin of patient 1's infection. Although patients 2 and 3 lived together, the genotype found in patient 2 differed from the (partial) genotype

found in patient 3. Yet another genotype was found in a patient from the same village (patient 4). However, an identical genotype found in patient 2 was found in a patient from a distant village (patient 5). The village had only 1 goat farm, and if this herd of goats was the source of infection for the farmer, his wife, and patient 4, it would have contained >1 genotype. At least 1 of the obtained genotypes has spread over a wider surface area in the Netherlands.

Our results show that the unprecedented, ongoing Q fever outbreak in the Netherlands involves multiple genotypes of *C. burnetii*. Because most of the genotypes differ only by a single repeat difference, they might represent microvariants of a hypervirulent strain that has been introduced in the Dutch animal population. MLVA schemes with up to 17 markers have been previously reported (3). In this “proof of concept” (applying direct genotyping of *C. burnetii* on clinical samples), we focused on the 3 shortest repeat units because we believed that these units might have the highest a priori chance of successful amplification in clinical samples (especially in serum/plasma). Similar genotypes as those reported here were found in the MLVA database (<http://mlva.u-psud.fr>), but these similarities need confirmation by using more markers. Although using only 3 markers may lead to poor discriminatory power, we were still able to distinguish 4 different genotypes in a relatively small collection of serum samples. We are currently exploring the use of additional MLVA markers.

Our results also show a poor correlation between DNA load and clinical symptoms. Multiple human and animal clinical samples, including serum and plasma, throat or genital swabs, or sputum and urine, may be useful for direct genotyping and outbreak source tracking.

Table. Genotyping results for human and animal clinical samples, Q fever outbreak, the Netherlands

Patient/animal no.	Sample	Ms27*	Ms28*	Ms34*	Symptoms	Ct value†	Location
Patient 1	Plasma	3	3	8	Severe	34.4	1
Ewe 1	Vaginal swab	3	3	8	None	25.7	1
Ewe 2	Vaginal swab	3	3	8	None	16.3	1
Ewe 3	Vaginal swab	3	3	8	None	18.8	1
Lamb 1	Throat swab	3	3	8	None	27.9	1
Lamb 2	Throat swab	3	3	8	None	29.9	1
Lamb 3	Throat swab	3	3	8	None	28.9	1
Patient 2	Urine	3	3	7	None	31.7	2
	Throat swab	3	3	7		31.8	
Patient 3	Urine	NR‡	3	4	Mild	36.7	2
Patient 4	Sputum	4	3	7	Severe	34.2	2
Patient 5	Sputum	3	3	7	Severe	31.9	3
Nine Mile	Reference strain	4*	6*	5*			

*The allele-calling convention used was as published (3), resulting in a 4, 6, 5 code assigned respectively to the 6-bp repeat unit loci Ms-27, Ms-28, and Ms-34 for the genome sequence of the Nine Mile RSA-493 strain (GenBank accession no. NC002971.1). Primers for these markers were redesigned to amplify significantly shorter PCR products and were combined into 1 multicolor multiplex PCR. Primer sequences for Ms-27 were 5'-HEX-TCTTTATTTTCAGGCCGGAGT-3' and 5'-GAACGACTCATTGAACACACG-3; for Ms-28, 5'-TMR-AGCAAAGAAATGTGAGGATCG-3' and 5'-GCCAAAGGGATATTTTGTCTTC-3; for Ms-34, 5'-FAM-TTCTTCGGTGAGTTGCTGTG-3' and 5'-GCAATGACTATCAGCGACTCGAA-3'.

†Cycle threshold (Ct) value obtained by using real-time PCR targeting the IS1111a element.

‡NR, no result obtained. A full genotype was obtained only in samples with the highest DNA loads (Ct value ≤35).

**Corné H.W. Klaassen,
Marrigje H. Nabuurs-Franssen,
Jeroen J.H.C. Tilburg,
Maurice A.W.M. Hamans,
and Alphons M. Horrevorts**

Author affiliations: Canisius Wilhelmina Hospital, Nijmegen, the Netherlands (C.H.W. Klaassen, M.H. Nabuurs-Franssen, J.J.H.C. Tilburg, A.M. Horrevorts); Radboud University Medical Centre, Nijmegen (M.H. Nabuurs-Franssen); and Food and Consumer Product Safety Authority, Eindhoven, the Netherlands (M.A.W.M. Hamans)

DOI: 10.3201/eid1504.081612

References

1. Raoult D, Marrie TJ, Mege JL. Natural history and pathophysiology of Q fever. *Lancet Infect Dis*. 2005;5:219–26. DOI: 10.1016/S1473-3099(05)70052-9
2. Schimmer B, Morray G, Dijkstra F, Schneeberger PM, Weers-Pothoff G, Timer A, et al. Large ongoing Q fever outbreak in the south of the Netherlands, 2008. *Euro Surveill*. 2008;13:pii=18939 [cited 2009 Feb 18]. Available from <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=18939>
3. Arricau-bouvery N, Hauck Y, Bejaoui A, Frangoulidis D, Bodier CC, Souriau A, et al. Molecular characterization of *Coxiella burnetii* isolates by infrequent restriction site-PCR and MLVA typing. *BMC Microbiol*. 2006;6:38. DOI: 10.1186/1471-2180-6-38

Address for correspondence: Corné H.W. Klaassen, Department of Medical Microbiology and Infectious Diseases, Canisius Wilhelmina Hospital, Weg door Jonkerbos 100, 6532 SZ Nijmegen, the Netherlands; email: c.klaassen@cwz.nl

Correlation between Buruli Ulcer and Vector-borne Notifiable Diseases, Victoria, Australia

To the Editor: Buruli ulcer (BU) is a destructive skin disease caused by the toxin-producing environmental pathogen *Mycobacterium ulcerans*. Since the 1980s, BU has emerged as a major public health problem in rural West and Central Africa (1), where some researchers have suggested a role for aquatic insects as either reservoirs or vectors of *M. ulcerans* (2,3). However, this hypothesis remains unproven (4).

In contrast to the emerging BU–endemic areas in tropical rural West Africa, the climate of the Australian state of Victoria is temperate, yet locally acquired BU also has increased there in recent years (5). In addition, notifications have varied markedly from year to year for reasons not yet explained.

During the investigation of a new outbreak of BU in Victoria, we demonstrated that *M. ulcerans* is detectable by PCR in mosquitoes and that being bitten by mosquitoes increases the odds of being diagnosed with BU (6,7). However, *M. ulcerans*–positive mosquitoes might reflect only the presence of *M. ulcerans* in the local environment and play no role in transmission. To further investigate links between BU and mosquitoes, we compared patterns of notifications of BU with other notifiable diseases in Victoria. In particular, we were interested in any association between BU and the locally transmitted vector-borne alphaviruses Ross River virus (RRV) and Barmah Forest virus (BFV). Areas of BU and RRV/BFV endemicity overlap geographically, but areas with RRV and BFV are more extensive and include inland river systems where BU has not so far been reported.