

# White-Nose Syndrome Fungus (*Geomyces destructans*) in Bat, France

Sébastien J. Puechmaille, Pascal Verdeyroux, Hubert Fuller, Meriadeg Ar Gouilh, Michaël Bekaert, and Emma C. Teeling

White-nose syndrome is caused by the fungus *Geomyces destructans* and is responsible for the deaths of >1,000,000 bats since 2006. This disease and fungus had been restricted to the northeastern United States. We detected this fungus in a bat in France and assessed the implications of this finding.

Biologists are struggling to understand a recent emerging infectious disease, white-nose syndrome (WNS) (1), which potentially threatens >20% of all mammalian diversity (bats) (2). WNS is a deadly epidemic that has swept through the northeastern United States over the past 3 years and caused the death of >1,000,000 bats, with decreases of ≈100% in some populations (3).

This disease is associated with hibernating, cave-roosting bats. A visually conspicuous white fungus grows on the face, ears, or wings of stricken bats; infiltration of the hyphae into membranes and tissues leads to severe damage (4). Bats that exhibit WNS have little or no fat reserves, which are essential for their survival throughout and after hibernation (5). The fungus associated with WNS is a newly described, psychrophilic (cold-loving) species (*Geomyces destructans*) (6). It is closely related to *G. pannorum*, which causes skin infections in humans (7).

Although it is not known whether the fungus is primarily responsible for deaths of bats or is a secondary infection, it is directly associated with deaths of bats (5). Bacteriologic, virologic, parasitologic, and postmortem evaluations for the cause of death did not identify any other agents, which reinforces the suspicion that this fungus is the causative agent (4,5). To date, WNS has been found only in the northeastern United States. However, researchers have suggested its presence in Europe. We investigated whether

*G. destructans* is present in bats in Europe and assessed the implications of its presence.

## The Study

During intensive monitoring of bat hibernation in France, 1 bat (*Myotis myotis*) found on March 12, 2009, near Périgueux (45°8'N, 0°44'E), showed a powdery, white fungal growth on its nose (Figure 1, panel A), which is characteristic of WNS. Sterile dry cotton swabs were used to collect fungus material from the nose of the bat. The bat was then weighed, measured, and released. Swabs were moistened with 50 μL of sterile water and streaked onto plates containing potato dextrose agar supplemented with 0.1% mycologic peptone. Plates (9 cm in diameter) were sealed with parafilm and incubated inverted at 10°C. A dense fungus growth developed within 14 days (Figure 1, panel B).

Cultures were established by transferring inoculum to other mycologic media, including malt extract agar and Sabouraud agar. Colonies on malt extract agar were initially white but after spore production and aging they quickly darkened from the center to a dull gray, often showing a faint green hue. Spores were hyaline, irregularly curved, broadly crescent-shaped (typically 6–8 μm long and 3–4 μm wide), and narrowed at each end, one of which was broadly truncate, often showing an annular frill (Figure 1, panel C). Fungal cultures have been deposited in the culture collection of the Industrial Microbiology Department of University College Dublin (Reference IMD Z2053).

Microscopic examination of the original swab samples showed numerous spores with the above-mentioned features. The psychrophilic nature of the fungus and its species-specific morphologic features (online Technical Appendix, [www.cdc.gov/EID/content/16/2/290-Techapp.pdf](http://www.cdc.gov/EID/content/16/2/290-Techapp.pdf)) led to the conclusion that this fungus was *G. destructans*, which was recently isolated from WNS-positive bats in the northeastern United States (6).

Two molecular markers were sequenced from 6 randomly chosen fungus cultures to confirm species identity. DNA was extracted by using a Blood and Tissue Kit (QIAGEN, Hilden, Germany) following the manufacturer's instructions with slight modifications (after step 3, we added an incubation time of 10 min at 70°C). The internal transcribed spacer (ITS) regions (ITS1, 5.8S, and ITS2) and the small subunit (SSU) of the rRNA gene were amplified separately.

PCRs were performed in 25-μL volumes containing 1 μL of DNA (10–75 ng/μL), 1.5 mmol/L MgCl<sub>2</sub>, 0.4 μmol/L of each primer, 0.2 mmol/L dNTP, 1× PCR buffer, and 1 U of Platinum Taq DNA Polymerase High Fidelity (Invitrogen, Carlsbad, CA, USA). Identical PCR cycling conditions were used for both fragments: an initial step at 95°C for 15 s; 10 cycles at 95°C for 30 s, 60°C (reduction of 2°C

Author affiliations: University College Dublin, Dublin, Ireland (S.J. Puechmaille, H. Fuller, M. Bekaert, E.C. Teeling); Groupe Chiroptères Aquitaine, Erdoia, France (P. Verdeyroux); and Institut Pasteur, Paris, France (M. Ar Gouilh)

DOI: 10.3201/eid1602.091391




Figure 1. A) *Myotis myotis* bat found in a cave on March 12, 2009, in France, showing white fungal growth on its nose (arrow). B) Fungus colony on malt extract medium after incubation for 3 weeks at 10°C. Scale bar = 1 cm. C) Clusters of unstained spores of *Geomyces destructans*. Spores in the inset were stained with lactophenol cotton blue, which shows the truncate spore base (arrows) and surface granulation. Scale bars = 10 µm.

every 2 cycles) for 1 min and 45 s, and 72°C for 1 min; 30 cycles at 95°C for 30 s, 50°C for 1 min and 45 s, and 72°C for 1 min; and a final step at 72°C for 10 min. PCR products were purified and sequenced in both directions by using primers listed in the Table. Complementary sequences were assembled and edited for accuracy by using Codon-Code Aligner version 3.0.3 ([www.codoncode.com/aligner/download.htm](http://www.codoncode.com/aligner/download.htm)).

The ITS and SSU sequences from the 6 WNS fungus cultures were identical. They were deposited in GenBank as single sequences: ITS (GQ489024) and SSU (GQ489025). Sequences obtained for the 2 genetic markers showed a 100% sequence identity with the described *G. destructans* fungus (Figure 2, panels A, B). Thus, morphologic and genetic data support the presence of *G. destructans* infection in a bat in France.

## Conclusions

Our results show that the WNS fungus was present in a bat in France and has implications for WNS research, bat conservation, and emerging infectious disease control. We suggest 3 possible scenarios for our findings. The first scenario is that the fungus has only recently arrived in Europe and all bats in Europe are now at risk for infection. Thus, conservation steps must be taken to minimize the spread of this disease, especially because this disease is specific for hibernating bats. After the hibernation period, *M. myotis* bats may migrate up to 436 km to reach their summer roosts (10), a behavior that could quickly increase the chance of fungus transmission. A second scenario is that the fungus has been present in Europe for a long time. Because mass deaths have not been observed in bats in Europe, these bats may be immune to WNS. Therefore, identification of mechanisms of this immunity will advance understanding of this disease and fungus resistance in mammals. The third scenario is that the *G. destructans* fungus is not the primary

cause of death but acts as an opportunistic pathogen in bats already immunocompromised by other pathogens such as viruses or bacteria (1). Comparison of pathogens in bats in Europe and the United States infected with *G. destructans* should identify the primary causative agent.

The bat in our study showing fungal growth was not underweight (Figure 2, panel C), as is typical of bats in the United States with WNS (4). This finding favors the second or third scenarios. Also, a 6-year (2004–2009) annual monitoring program of wintering bat populations at the site and 5 sites within a 2-km radius did not show any cases of WNS or deaths and showed stable bat populations. The 3 scenarios indicate that studying *G. destructans* in bats in Europe and the United States is necessary to understand and control this disease.

Another fungus, *Batrachochytrium dendrobatidis*, is the etiologic agent responsible for chytridiomycosis (11), which currently threatens >50% of all amphibian species and is primarily responsible for the global decrease and extinction of >200 amphibian species in the past decade (12). Because bats account for >20% of mammalian diversity (2) and play major roles in ecosystem functions, we need to understand, monitor, and control the progression of WNS. Otherwise, we may be faced with similar unprec-

Table. Primers used for PCR amplification and sequencing of fungus in bats, France\*

Gene	Primer sequence (5' → 3')	PCR
ITS	TCCTCCGCTTATTGATATGC	Forward
	GGAAGTAAAAGTCGTAACAAGG	Reverse
SSU rRNA	CTGGTTGATTCTGCCAGT	Forward
	AAACCTTGTTACGACTTTTA	Reverse
	CCGGAGAAGGAGCCTGAGAAAC	†
	AACTTAAAGGAATTGACGGAAG	†
	CTCATTCCAATTACAAGACC	†
GAGTTTCCCCGTGTTGAGTC	†	

\*ITS, internal transcribed spacer; SSU, small subunit.

†Used for sequencing only.

edented and irreversible losses of mammalian biodiversity and entire ecosystems. Because bats control insect populations throughout the world (13–15), a large decrease in bat

populations would result in insect proliferations that would damage agricultural crops and spread many insect-borne diseases.




Figure 2. Genetic distance between fungus A) internal transcribed spacer (ITS) (474 nt) and B) small subunit (SSU) rRNA (1,865 nt) gene sequences and other closely related fungus species present in GenBank. Results are based on pairwise sequence comparisons with gaps and missing data removed. Error bars in panels A and B indicate mean  $\pm$  SD. C) Estimation of weight of *Myotis myotis* bats after hibernation as a function of the range of percentage of weight loss reported. Posthibernation of a bat's weight was estimated from prehibernation measured weights ( $n = 155$  bats) minus winter fat loss. A strong positive relationship exists between body mass and fat mass during prehibernation (8). Fat reserves between 15% and 30% of body mass at the onset of hibernation have been reported to be necessary for *Myotis* species to survive winter (9). The posthibernation weight ( $W_{\text{post}}$ ) was thus estimated as  $W_{\text{post}} = W_{\text{pre}} - (W_{\text{pre}} \times W_{\text{loss}} / 100)$ , where  $W_{\text{pre}}$  is prehibernation weight and  $W_{\text{loss}}$  is percentage of body mass lost during hibernation. Mean  $\pm$  SD prehibernation weight of 155 bats captured in France during August–October 2009 ( $27.42 \pm 2.87$  g) was used for the estimate. Black line represents the mean, gray area represents the mean  $\pm$  SD, and red dashed line represents the 24-g weight of the bat caught in France with white-nose syndrome posthibernation. The bat was in good condition (24 g) because it weighed more than the expected average for a posthibernating bat despite having *Geomyces destructans* growth on its snout.

## Acknowledgments

We thank Arnaud Le Houédec, Etienne Ouvrard, Les Naturalistes Vendéens/LPO-85, Christophe Hervé, Nicolas Galand, and Nicolas Harter for providing morphometric data.

This study was supported by a grant from The President of Ireland Young Researcher Award Science Foundation Ireland to E.C.T.


Dr Puechmaille is a postdoctoral fellow at the School of Biology and Environmental Science at University College Dublin. His research interests focus on evolution, speciation, ecology, and conservation with an emphasis on bats.

## References

1. Turner GG, Reeder DM. Update of white nose syndrome in bats, September 2009. *Bat Research News*. 2009;50:47–53.
2. Simmons NB. Order Chiroptera. In: Wilson DE, Reeder DM, editors. *Mammal species of the world: a taxonomic and geographic reference*, 3rd ed. Baltimore: Johns Hopkins University Press; 2005. p. 312–529.
3. Consensus statement. Austin (TX): Second WNS Emergency Science Strategy Meeting; 2009 May 27–28 [cited 2009 Nov 30]. <http://www.batcon.org/pdfs/whitenose/ConsensusStatement2009.pdf>
4. Meteyer CU, Buckles EL, Blehert DS, Hicks AC, Green DE, Shearn-Bochsler V, et al. Histopathologic criteria to confirm white-nose syndrome in bats. *J Vet Diagn Invest*. 2009;21:411–4.
5. Blehert DS, Hicks AC, Behr M, Meteyer CU, Berlowski-Zier BM, Buckles EL, et al. Bat white-nose syndrome: an emerging fungal pathogen? *Science*. 2009;323:227. DOI: 10.1126/science.1163874
6. Gargas A, Trest MT, Christiansen M, Volk TJ, Blehert DS. *Geomyces destructans* sp. nov. associated with bat white-nose syndrome. *Mycotaxon*. 2009;108:147–54.
7. Gianni C, Caretta G, Romano C. Skin infection due to *Geomyces pannorum* var. *pannorum*. *Mycoses*. 2003;46:430–2. DOI: 10.1046/j.1439-0507.2003.00897.x
8. Kunz TH, Weazen JA, Burnett CD. Changes in body mass and fat reserves in pre-hibernating little brown bats (*Myotis lucifugus*). *Ecology*. 1998;79:8–17.
9. Johnson SA, Brack VJ, Rolley RE. Overwinter weight loss of Indiana bats (*Myotis sodalis*) from hibernacula subject to human visitation. *American Midland Naturalist*. 1998;139:255–61. DOI: 10.1674/0003-0031(1998)139[0255:OWLOIB]2.0.CO;2
10. Dietz C, Von Helversen O, Dietmar N. *Bats of Britain, Europe and northwest Africa*. London: A and C Black Publishers Ltd.; 2009.
11. Voyles J, Young S, Berger L, Campbell C, Voyles WF, Dinudom A, et al. Pathogenesis of chytridiomycosis, a cause of catastrophic amphibian declines. *Science*. 2009;326:582–5. DOI: 10.1126/science.1176765
12. Fisher MC, Garner TW, Walker SF. Global emergence of *Batrachochytrium dendrobatidis* and amphibian chytridiomycosis in space, time, and host. *Annu Rev Microbiol*. 2009;63:291–310. DOI: 10.1146/annurev.micro.091208.073435
13. Williams-Guillen K, Perfecto I, Vandermeer J. Bats limit insects in a neotropical agroforestry system. *Science*. 2008;320:70. DOI: 10.1126/science.1152944

14. Federico P, Hallam TG, McCracken GF, Purucker ST, Grant WE, Correa-Sandoval AN, et al. Brazilian free-tailed bats as insect pest regulators in transgenic and conventional cotton crops. *Ecol Appl*. 2008;18:826–37. DOI: 10.1890/07-0556.1
15. Kalka MB, Smith AR, Kalko EK. Bats limit arthropods and herbivory in a tropical forest. *Science*. 2008;320:71. DOI: 10.1126/science.1153352

Address for correspondence: Emma C. Teeling, School of Biology and Environmental Science, University College Dublin, Belfield, Dublin 4, Ireland; email: emma.teeling@ucd.ie



# White-Nose Syndrome Fungus (*Geomyces destructans*) in Bat, France

## Technical Appendix

Colonies of the fungus on malt extract agar were initially white but with spore production and age, quickly darkened from the center to a dull gray, often with a faint green hue. Colonies were compact and domed, reaching diameters of  $\approx 16$  mm in 24 days at 10°C. Hyphae were 2–3  $\mu\text{m}$  in diameter and septate. Older colonies had a strong moldy smell, and brown droplets were exuded on their surfaces. The under surface of colonies also darkened centrally with age, from brown to dark brown. Spore production was prolific from an early stage of growth, giving the colonies a powdery texture; clumps of white mycelia were also occasionally present on colonies.

Spores were typically produced from the tip of groups of short branches, sometimes in loose verticils, at the apices of conidiophores. Spores also developed from side branches and directly from the conidiophore surface. The spores were produced either in small clusters, in short chains of 2–3 spores, or singly. They were hyaline, irregularly curved, broadly crescent-shaped (typically 6–8  $\mu\text{m}$  long and 3–4  $\mu\text{m}$  wide) narrowed at each end, one of which was broadly truncate, often with an annular frill. In side view, many spores appeared obovoid; intercalary spores were barrel-shaped. Spore walls were relatively thick and an outer surface granulation was evident on some.

Microscopic examination of the original swab sample had shown numerous spores having the above-mentioned features. The psychrophilic nature of the fungus and its morphologic features, lead us to conclude that this fungus is *Geomyces destructans*, which was recently isolated from bats in the northeastern United States that had white-nose syndrome.




Figure. Phylogenetic relationship between *Geomyces destructans* and other fungus species. A) Bayesian tree for internal transcribed spacer (ITS) alignment (474 nt, Hasegawa, Kishino, and Yano +  $\Gamma$  model). B) Bayesian tree for small subunit (SSU) rRNA alignment (1,865 nt, general time reversible +  $\Gamma$  + invariant sites model). Dendrograms were constructed by using Bayesian analyses with BEAST software ([http://beast.bio.ed.ac.uk/Main\\_Page](http://beast.bio.ed.ac.uk/Main_Page)). Most appropriate substitution models were selected according to ModelTest version 3.7 (<http://darwin.uvigo.es/software/modeltest.html>). A strict molecular clock model was applied. No outgroup was specified and the constant population size coalescent was used as a tree prior. The program was run for 10,000,000 generations and sampled every 500. The first 1,000,000 generations were discarded as burn-in. Effective sample sizes for the estimated parameters and posterior

probability as calculated with the program Tracer version 1.4 (<http://tree.bio.ed.ac.uk/software/tracer>) were >800. Bayesian posterior probability values are shown near each supported node (>0.5). The fungus strain identified in this study is shown in **boldface**. Scale bars indicate nucleotide substitutions per site.