

Appendix Table 1. Household secondary attack rates for ARI, ILI, and laboratory-confirmed pandemic (H1N1) 2009, by household size, Region 8, Texas, April–May 2009*

Household size, no. persons	Index case-patients	Household contacts		Total household contacts, A + B	Ill household members not included	Secondary attack rate, A/A + B, %
		Secondary case-patients, A	Not ill, B			
ARI						
2–3	21	7	24	31	1	23
4	23	11	58	69	0	16
5–6	26	10	101	111	4	9
7–9	7	4	41	45	3	9
All households	77	32	224	256	8	13
ILI						
2–3	21	7	24	31	1	23
4	23	8	61	69	0	12
5–6	26	5	107	112	3	4
7–9	7	4	42	46	2	9
All households	77	24	234	258	6	9
Laboratory-confirmed pandemic (H1N1) 2009						
2–3	21	2	30	32	0	6
4	23	4	65	69	0	6
5–6	26	4	110	114	1	4
7–9	7	1	47	48	0	2
All households	77	11	252	263	1	4

*ARI, acute respiratory infection; ILI, influenza-like illness (fever measured or subjective and cough or sore throat). Ill household members were not included in the calculation of the secondary attack rate if they had the same symptom onset as the index case or if symptom onset was not known.