

Novel Betaherpesvirus in Bats

Shumpei Watanabe,¹ Ken Maeda,¹ Kazuo Suzuki,
Naoya Ueda, Koichiro Iha, Satoshi Taniguchi,
Hiroshi Shimoda, Kentaro Kato,
Yasuhiro Yoshikawa, Shigeru Morikawa,
Ichiro Kurane, Hiroomi Akashi,
and Tetsuya Mizutani

Because bats are associated with emerging zoonoses, identification and characterization of novel viruses from bats is needed. Using a modified rapid determination system for viral RNA/DNA sequences, we identified a novel bat betaherpesvirus 2 not detected by herpesvirus consensus PCR. This modified system is useful for detecting unknown viruses.

Since the 1990s, bats have been associated with several emerging zoonotic agents, including Hendra, Nipah, Ebola, lyssa, and severe acute respiratory syndrome coronavirus-like viruses (1). Bats seem to have great potential as reservoirs for emerging viruses. Therefore, to understand the role of bats as a host species, identification and characterization of novel viruses from bats is needed. For virus isolation, we have been attempting to establish primary cell cultures from various bats (2,3). Using a rapid determination system for viral RNA sequences (RDV), we discovered a novel adenovirus and gammaherpesvirus in bats (2,4). This system, which we simplified to a less laborious one (5), is useful for detecting viruses, regardless of virus species (6).

The Study

During June–August, 2008, with the permission of the governor of Wakayama Prefecture, Japan, we caught 8 insectivorous vespertilionid bats, *Miniopterus fuliginosus*, and used their spleens and kidneys to establish primary cell cultures. During passage of the primary spleen adherent cells, cytopathic effect (cell death) was noted at third passage. The collected supernatant was injected into fresh primary kidney cells and caused apparent cytopathic effect at first passage.

Author affiliations: The University of Tokyo, Tokyo, Japan (S. Watanabe, N. Ueda, K. Iha, S. Taniguchi, K. Kato, Y. Yoshikawa, H. Akashi); Yamaguchi University, Yamaguchi, Japan (K. Maeda, H. Shimoda); Hikiwa Park Center, Wakayama, Japan (K. Suzuki); and National Institute of Infectious Diseases, Tokyo (S. Morikawa, I. Kurane, T. Mizutani)

DOI: 10.3201/eid1606.091567

Before using the RDV method, we had attempted to detect herpesvirus by nested PCR with the consensus primer sets DFA, ILK, KG1, TGV, and IYG, which were designed according to the consensus-degenerate hybrid oligonucleotide primers program (7). These consensus degenerate primers are effective for detecting many herpesviruses from vertebrate hosts. However, in this study they failed to detect any herpesviruses.

We then attempted to detect herpesvirus by using RDV version 3.1, our modification from version 3.0 (5). The adapters and primers for construction of the second cDNA library in RDV version 3.1 were newly designed and replaced those used in RDV version 3.0 (online Technical Appendix 1, www.cdc.gov/EID/content/16/6/986-Techapp1.pdf). Both adapters have sticky-end structures digested with *Sau3AI* or *HpyCH4 IV*. RDV version 3.1 can determine an unknown viral cDNA fragment with 64 primer pairs, which we used for constructing the second cDNA library.

With RDV version 3.1, we obtained 4 unknown cDNA fragments, which had no matches in a BLASTn (www.ncbi.nlm.nih.gov/blast/Blast.cgi) search. In a BLASTx search, 1 cDNA fragment (deduced sequence of 29 aa) was homologous to the glycoprotein B (gB) amino acid sequence of the tupaiid herpesvirus 1 (TuHV-1) (79% identity), which belongs to subfamily *Betaherpesvirinae*. We designed new consensus-degenerate hybrid oligonucleotide primers (<http://blocks.fhrc.org/codehop.html>) selective for the betaherpesvirus gB and DNA polymerase (DPO) genes, and we determined the complete gB sequence and the partial DPO sequence of the isolated virus (5,029 bp, DNA Data Bank of Japan accession no. AB517983). BLAST search indicated that the complete gB sequence was novel and most similar to that of TuHV-1 (59% aa sequence identity) (online Appendix Figure, www.cdc.gov/EID/content/16/6/986-appF.htm). We named the isolated virus bat betaherpesvirus 2 (BatBHV-2).

We constructed a phylogenetic tree by using the neighbor-joining method with the gB amino acid sequence and the available sequences of known herpesviruses (Figure). The phylogenetic tree based on betaherpesvirus gB genes showed that BatBHV-2 is most closely related to TuHV-1 and caviid herpesvirus 2 (guinea pig cytomegalovirus). The subfamily *Betaherpesvirinae* consists of the genera *Cytomegalovirus*, *Muromegalovirus*, and *Roseolovirus*. TuHV-2 and caviid herpesvirus 2 are species unassigned to any genus in the subfamily *Betaherpesvirinae*.

In May 2009, we collected, again with permission, another 50 bats belonging to 1 species, *M. fuliginosus*, from the same location for an epizootologic study (online Technical Appendix 2, www.cdc.gov/EID/content/16/6/986-Techapp2.pdf). Spleens and blood were collected from all

¹These authors contributed equally to this article.

bats, and other organs (liver, kidney, lung, brain, intestine, trachea, and urinary bladder) were collected from 10 bats. Nested PCR was performed by using specific primers selective for the DPOL gene of BatBHV-2, and PCR products were subjected to direct sequencing. Viral nucleotide sequences were obtained from 4 of the 50 spleen samples. Each nucleotide sequence showed complete identity to the partial DPOL sequence of the BatBHV-2. Other organs and serum collected from 2 of the bats were also tested by nested PCR, and viral DNA was detected in the liver, kidneys, and lungs of both bats.

Conclusions

Although PCRs with consensus primers effectively detect known and unknown viruses, they failed to detect BatBHV-2, possibly because of minor mismatches between the sequences of BatBHV-2 and the primer sets (TGV, IYG, and KG1). The variety of virus sequences and gene mutations often prevents successful amplification of virus genes. RDV, however, can detect viral cDNA fragments




Figure. Phylogenetic tree based on the deduced amino acid sequences of complete glycoprotein B. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1,000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The tree was rooted to herpes simplex virus type 1 (X14112). The evolutionary distances were computed by using the Poisson correction method and are in units of the number of amino acid substitutions per site. All positions containing gaps and missing data were eliminated from the dataset. The final dataset included a total of 698 positions. Phylogenetic analyses were conducted in MEGA4 (8). The herpesviruses used for comparison and their accession numbers are as follows: Epstein-Barr virus 1 (EBV, NC_007605), caviid herpesvirus (CavHV-2, FJ355434); mouse cytomegalovirus (MuHV-1, NC_004065), human cytomegalovirus (HCMV, X17403), human herpesvirus 6 (HHV-6, AF157706), human herpesvirus 7 (HHV-7, AF037218), human herpesvirus 8 (HHV-8, AF148805), rat cytomegalovirus (MuHV-2, NC_002512), cercopithecine herpesvirus 8 (CeHV-8, AY186194), saimiriine herpesvirus 2 (SaHV-2, NC_001350), and tupaiid herpesvirus 1 (TuHV-1, AF281817). Scale bar indicates evolutionary distance.

independent of virus species and thus is useful as a first-choice tool for identifying emerging known and unknown viruses in animals and humans.

BLAST search showed that the complete gB sequence of the isolated virus was novel and most similar to that of TuHV-1. Recently, bats have been described as hosts for herpesviruses in several countries in Europe, America, Africa, and Asia (4,9,10). Wibbelt et al. reported that the partial DPOL sequence (175 bp) of a betaherpesvirus, bat betaherpesvirus 1 (BatBHV-1), was obtained from several insectivorous bat species (10). Although the length of the BatBHV-1 sequence was short, similarity between BatBHV-1 and BatBHV-2 was relatively high (58%). BatBHV-1 is most similar to TuHV-1(61%). These findings suggest that BatBHV-2 is a different species than BatBHV-1.

Our epizootologic study found relatively high (8%) prevalence of BatBHV-2 in insectivorous bats. Although the virus genome was detected in a few parenchymal organs by nested PCR, no amplification was possible for serum, intestine, or urinary bladder samples, which may exclude apparent virus shedding by the bats. In addition, all 50 bats collected appeared clinically healthy. To understand the life cycle of this virus, the possibility of a latent infection in these insectivorous bats must be explored.

Acknowledgments

We thank Momoko Ogata and Miho Nishimura for their assistance.

This study was supported in part by grants from the Japan Society for the Promotion of Science; the Ministry of Health, Labor, and Welfare; and the Ministry of Education, Culture, Sports, Science, and Technology, Japan.


Mr Watanabe is a PhD student in the Department of Veterinary Microbiology, Graduate School of Agricultural and Life Sciences, The University of Tokyo. His research interests include the epidemiology and pathogenic mechanisms of emerging viruses from bats.

References

1. Wong S, Lau S, Woo P, Yuen KY. Bats as a continuing source of emerging infections in humans. *Rev Med Virol*. 2007;17:67–91. DOI: 10.1002/rmv.520
2. Maeda K, Hondo E, Terakawa J, Kiso Y, Nakaichi N, Endoh D, et al. Isolation of novel adenovirus from fruit bat (*Pteropus dasymallus yayeyamae*). *Emerg Infect Dis*. 2008;14:347–9. DOI: 10.3201/eid1402.070932
3. Omatsu T, Watanabe S, Akashi H, Yoshikawa Y. Biological characters of bats in relation to natural reservoir of emerging viruses. *Comp Immunol Microbiol Infect Dis*. 2007;30:357–74. DOI: 10.1016/j.cimid.2007.05.006
4. Watanabe S, Ueda N, Iha K, Masangkay JS, Fujii H, Alviola P, et al. Detection of a new bat gammaherpesvirus in the Philippines. *Virus Genes*. 2009;39:90–3. DOI: 10.1007/s11262-009-0368-8

5. Watanabe S, Mizutani T, Sakai K, Kato K, Tohya Y, Fukushi S, et al. Ligation-mediated amplification for effective rapid determination of viral RNA sequences (RDV). *J Clin Virol*. 2008;43:56–9. DOI: 10.1016/j.jcv.2008.05.004
6. Mizutani T, Endoh D, Okamoto M, Shirato K, Shimizu H, Arita M, et al. Rapid genome sequencing of RNA viruses. *Emerg Infect Dis*. 2007;13:322–4. DOI: 10.3201/eid1302.061032
7. VanDevanter DR, Warrener P, Bennett L, Schultz ER, Coulter S, Garber RL, et al. Detection and analysis of diverse herpesviral species by consensus primer PCR. *J Clin Microbiol*. 1996;34:1666–71.
8. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*. 2007;24:1596–9. DOI: 10.1093/molbev/msm092
9. Razafindratsimandresy R, Jeanmaire EM, Counor D, Vasconcelos PF, Sall AA, Reynes JM. Partial molecular characterization of alpha-herpesviruses isolated from tropical bats. *J Gen Virol*. 2009;90:44–7. DOI: 10.1099/vir.0.006825-0
10. Wibbelt G, Kurth A, Yasmum N, Bannert M, Nagel S, Nitsche A, et al. Discovery of herpesviruses in bats. *J Gen Virol*. 2007;88:2651–5. DOI: 10.1099/vir.0.83045-0

Address for correspondence: Tetsuya Mizutani, Department of Virology 1, National Institute of Infectious Diseases, Gakuen 4-7-1, Musashimurayama, Tokyo 208-0011, Japan; email: tmizutan@nih.go.jp



Search
past issues

EID
online
www.cdc.gov/eid

Novel Betaherpesvirus in Bats

Technical Appendix 1

System for Rapid Determination of Viral RNA Sequences, Version 3.1

The system for rapid determination of viral RNA sequences (RDV) version 3.0 (1) was modified to simplify the procedure. The modified method was designated version 3.1. Adapters and primers for construction of second cDNA library in RDV version 3.1 were newly designed and used instead of those in RDV version 3.0 (see Figure below. A: adapters, B: primers). Both adapters have sticky-end structures digested with *Sau3AI* or *HpyCH4IV*. The RDV version 3.1 method includes the 4 procedures described below.

1. RNA extraction. To eliminate contaminating cellular RNA and DNA from the samples, culture supernatant was treated with DNase I and RNase A as described previously (2), following RNA extraction by using a Total RNA isolation mini kit (Agilent Technologies, Santa Clara, CA, USA).
2. Construction of first cDNA library. A whole-transcriptome amplification system (WTA; Sigma-Aldrich, Saint Louis, MO, USA) was used to amplify double-stranded viral cDNA as described previously (3).
3. Construction of second cDNA library. After the first cDNA library purification with the MonoFas DNA isolation system (GL Sciences, Tokyo, Japan), DNA was

digested with 20 U of *Hpy*CH4IV (New England Biolabs, Ipswich, MA, USA) and *Sau*3AI (New England Biolabs) at 37°C for 60 min, and the digested DNA was again purified by using MonoFas. For construction of the second cDNA library, aliquots of 2.5 uL of DNA solution, 2.5 uL of distilled water, 2.5 uL of RDV-Adaptor-*Sau*3AI adaptor (10 uM) and RDV-Adaptor-*Hpy*CH4IV adaptor (10 uM) were mixed. The sequences and structures of the adapters are shown in figure below. A ligation-convenience kit (Nippon Gene, Toyama, Japan) was used for adaptor ligation. The DNA solution and 10 uL of ligation mix were reacted at 16°C for 30 min, and the DNA was isolated by using MonoFas. The second cDNA library was amplified by PCR with specially designed primer sets; the forward primers had 5 nt including GATC (*Sau*3AI-digested sequence) and 1 variable nucleotide added to the 3' end of the RDV-Adaptor-*Sau*3AI sequence, and the reverse primers had 6 nt including ACGT (*Hpy*CH4IV-digested sequence) and 2 variable nucleotides added to the 3' end of the RDV-Adaptor-*Hpy*CH4IV sequence. The PCR mixture was prepared by mixing 15 uL of AmpliTaq Gold PCR Master Mix containing AmpliTaq Gold, 0.5 uL of forward primer for RDV-Adaptor-*Sau*3AI (10 uM), 0.5 uL of reverse primer for RDV-Adaptor-*Hpy*CH4IV (10 uM), 1 uL of DNA solution, and 13 uL of distilled water. PCR was performed by using 64 primers under the conditions described previously (1).

4. Direct sequencing. After electrophoresis of PCR products on agarose gels, bands >120 bp were excised, and DNA was extracted from the gel by using MonoFas. Direct sequencing was performed by using the forward or reverse primer.

References

1. Omatsu T, Watanabe S, Akashi H, Yoshikawa Y. Biological characters of bats in relation to natural reservoir of emerging viruses. *Comp Immunol Microbiol Infect Dis.* 2007;30:357–74.
[PubMed DOI: 10.1016/j.cimid.2007.05.006](https://pubmed.ncbi.nlm.nih.gov/doi/10.1016/j.cimid.2007.05.006)
2. Watanabe S, Ueda N, Iha K, Masangkay JS, Fujii H, Alviola P, et al. Detection of a new bat gammaherpesvirus in the Philippines. *Virus Genes.* 2009;39:90–3. [PubMed DOI: 10.1007/s11262-009-0368-8](https://pubmed.ncbi.nlm.nih.gov/doi/10.1007/s11262-009-0368-8)
3. Sakai K, Mizutani T, Fukushi S, Saijo M, Endoh D, Kurane I, et al. An improved procedure for rapid determination of viral RNA sequences of avian RNA viruses. *Arch Virol.* 2007;52:1763–5. [DOI: 10.1007/s00705-007-0999-9](https://pubmed.ncbi.nlm.nih.gov/doi/10.1007/s00705-007-0999-9)

A

RDV-Adaptor-Sau3AI

```

5'   TGTCACGACTGAACCGAAC   3'
    CAGGTGCTGACTTGGCTTGCTAG
3'

```

RDV-Adaptor-HpyCH4IV

```

5'   CGGAGAGCATACCCTTACGAA   3'
    TGCCTCTCGTATGGGAATGCTTGC
3'

```

B

```

S1 5'- tgtcacgactgaaccgaacgac a - 3'
S2  tgtcacgactgaaccgaacgac t
S3  tgtcacgactgaaccgaacgac g
S4  tgtcacgactgaaccgaacgac c

```

```

H1 5'- cggagagcatacccttacgaacgt aa - 3'
H2  cggagagcatacccttacgaacgt af
H3  cggagagcatacccttacgaacgt aj
H4  cggagagcatacccttacgaacgt ac
H5  cggagagcatacccttacgaacgt ta
H6  cggagagcatacccttacgaacgt ff
H7  cggagagcatacccttacgaacgt ty
H8  cggagagcatacccttacgaacgt te
H9  cggagagcatacccttacgaacgt ga
H10 cggagagcatacccttacgaacgt gf
H11 cggagagcatacccttacgaacgt gg
H12 cggagagcatacccttacgaacgt gc
H13 cggagagcatacccttacgaacgt ca
H14 cggagagcatacccttacgaacgt cf
H15 cggagagcatacccttacgaacgt cy
H16 cggagagcatacccttacgaacgt cc

```

Adapter sequence *HpyCH4IV*-digested sequence

Novel Betaherpesvirus in Bats

Technical Appendix 2

Epizootologic Study

Fifty bats (nos. Mf01–Mf50) belonging to 1 species, *M. fuliginosus*, were collected; 28 were female, 25 of which were pregnant. Spleens and serum samples were collected from all bats, and other organs (liver, kidney, lung, brain, intestine, trachea, and urinary bladder) were collected from 10 bats (nos. Mf1–Mf10). DNA was extracted from each spleen sample and subjected to viral genome detection. Nested PCR was performed by using specific primers targeting the DNA polymerase (DPOL) gene of bat betaherpesvirus 2 (BatBHV-2), and PCR products were subjected to direct sequencing. Viral nucleotide sequences were obtained from 4 (bats. Mf4, Mf7, Mf14, Mf33) of the 50 spleen samples. Bats Mf4, Mf7 and Mf33 were pregnant. The obtained nucleotide sequences showed complete identity to the partial DPOL sequence of the isolated virus. Other organs and serum collected from bats Mf4 and Mf7 were also tested by nested PCR. Viral DNA was detected in organs (liver, kidney, and lung) of each of these bats. The pattern of viral distribution in organs was identical between bats Mf4 and Mf7. These results were summarized below.

Results of nested PCR

| Bat no. | Sex | Spleen | Intestine | Liver | Kidney | Bladder | Lung | Trachea | Brain | Sera |
|---------|----------|--------|-----------|-------|--------|---------|------|---------|-------|------|
| M14 | F | + | - | + | + | - | + | - | - | - |
| M17 | F | + | - | + | + | - | + | - | - | - |
| M14 | F | + | ND | ND | ND | ND | ND | ND | ND | ND |
| M133 | M | + | ND | ND | ND | ND | ND | ND | ND | ND |
| ND | Not done | | | | | | | | | |