

Evolution of Seventh Cholera Pandemic and Origin of 1991 Epidemic, Latin America

Connie Lam, Sophie Octavia, Peter Reeves, Lei Wang, and Ruiting Lan

Thirty single-nucleotide polymorphisms were used to track the spread of the seventh pandemic caused by *Vibrio cholerae*. Isolates from the 1991 epidemic in Latin America shared a profile with 1970s isolates from Africa, suggesting a possible origin in Africa. Data also showed that the observed genotypes spread easily and widely.

The seventh cholera pandemic began in 1961, and by 1966, it had affected most of Asia. Cholera incidence then decreased slightly until 1971, when an upsurge was observed in Africa and Europe, which had been free of cholera for >100 years (1). Cholera rates remained relatively low during the 1980s, with the disease confined to Asia and Africa. However, 2 major cholera outbreaks appeared in the 1990s: first, a resurgence of cholera in Africa, and, second, outbreaks that started in Peru became the first cholera epidemic in Latin America since 1895 (2). In addition, a novel serotype caused major outbreaks on the Indian subcontinent in 1992. That strain was referred to as O139 Bengal and was later shown to be a variant of the seventh pandemic clone with its replacement of the O antigen (1). Pulsed-field gel electrophoresis (3), amplified fragment length polymorphism analysis (4), and ribotyping (1) have been applied to seventh pandemic isolates but did not fully resolve the relationships of the various outbreaks. In this study, we used genome-wide single-nucleotide polymorphisms (SNPs) to track the evolution and spread of the seventh cholera pandemic, including the O139 Bengal strain.

The Study

The availability of complete genome sequences of a pre-seventh pandemic isolate, M66-2 (5), a seventh

pandemic isolate, N16961 (6), and the partial genome sequence of an O139 Bengal isolate, MO10 (7), enabled identification and use of SNPs as evolutionary markers in *Vibrio cholerae*. A set of 18 SNPs was chosen from 125 N16961 SNPs (5) and 12 SNPs selected from 59 identified by comparison of the N16961 and MO10 genome sequences. The SNPs selected were mostly from genes with known function and were distributed throughout the 2 chromosomes for the N16961 SNPs and the large chromosome for the MO10 SNPs. We have previously shown that recombinant regions could be identified by the differences in distribution of SNPs in such regions (5); for this study, only mutational SNPs were selected.

The 30 SNPs (online Technical Appendix, www.cdc.gov/EID/content/16/7/1130-Techapp.pdf) were used to type a collection of 64 seventh pandemic *V. cholerae* isolates. SNPs were detected by using hairpin primer real-time PCR. SNP data for 3 complete *V. cholerae* genomes (M66-2, N16961, MJ-1236) and 4 partially sequenced genomes (MO10, RC9, B33, CIRS 101) (7) were obtained from the National Center for Biotechnology Information (Rockville, MD, USA) and included in the analysis. The 71 isolates were divided into 10 SNP profiles by using the 30 SNPs (online Appendix Table, www.cdc.gov/EID/content/16/7/1130-appT.htm). Three profiles were represented by 1 isolate only, whereas the remaining profiles contained 4–17 isolates. The Simpson index of diversity for all SNPs combined was 0.929.

A maximum-parsimony tree (Figure) was constructed to show the relationships of the SNP profiles. The tree was fully resolved with no reverse or parallel changes in the seventh pandemic isolates. The pre-seventh pandemic strains were used as an outgroup and placed at the base of the tree. Six groups could be distinguished, with each group containing SNP profiles differing by no more than 1 SNP. The ladderized tree shows the stepwise evolution of the SNP profiles and groups. Group I at the bottom of the tree originated in Indonesia in 1961. It contains mostly isolates from Asia from the 1960s but continued to be isolated in Southeast Asia. The other groups evolved sequentially. Group II contains isolates from Africa from the 1970s to the 1990s and all 4 isolates from Latin America; group III contains earlier 1970s isolates from Asia and 1980s isolates from Africa; group IV contains late 1970s and 1980s isolates from Asia only; while group V contains 1990s isolates from Asia and Africa. Group VI contains only O139 isolates with the same SNP profile.

Conclusions

The presence of isolates from Africa in 3 groups can be explained by multiple introductions of cholera into Africa from cholera-endemic regions in Asia. The isolates in the first introduction in the 1970s shared a single origin (group

Author affiliations: University of New South Wales, Sydney, New South Wales, Australia (C. Lam, S. Octavia, R. Lan); University of Sydney, Sydney (P. Reeves); and Nankai University, Tianjin, People's Republic of China (L. Wang)

DOI: 10.3201/eid1607.100131

II). However, during the late 1980s and early 1990s, cholera outbreaks appeared to be caused by strains from 3 related sources. The first source came from group II, which was already established in Africa, and the second and third sources came from groups III and V in Asia. Because both groups were supported by multiple SNPs, it is less likely

that the 1970s isolates from Africa and Asia evolved in parallel to fall into the same groups. Additionally, B33 in Group V carries a classical CTX prophage (8), which indicates that this strain likely originated in Asia.

The cholera epidemic in Latin America was originally suspected to have come from Asia and to have been facilitated by the discharge of contaminated ballast water into Peruvian ports by international trade ships (2). However, the isolates from Latin America analyzed in this study were closely related to isolates found in Africa in the 1970s and 1990s. Four isolates, 2 from Peru and 1 each from Brazil and French Guiana, had an SNP profile identical to the 12 isolates from Africa that originated during that period. No isolates from Asia fell into this group. This finding suggests that the strain that caused the epidemic in Latin America came from Africa rather than Asia.

The outbreak in Peru occurred in parallel with the upsurge of cholera generally in Africa (1) and could have been imported at that time. However, the epidemic strain may have reached Latin America well before it caused the epidemic in 1990s, given the ability of the organism to persist in the marine environment for long periods (2). The strain could have been brought into the region during the mass migration from Africa to Latin America in the 1970s (9). The isolates from Latin America differ by 1 locus from the other seventh pandemic strains (Asia and Africa) by multilocus enzyme electrophoresis (10) and also differ in the *Vibrio* spp. seventh pandemic island-II gene cluster (11), which suggests that further evolution occurred after the strain separated from its likely ancestral strain from Africa and supports this latter scenario. The epidemic strain in Latin America could not have originated from the 1990s isolates from Asia in Groups III–V because they arose later than Group II isolates. However, a 1970s lineage in Asia that spread to Africa may have remained in Asia until the 1990s but was not represented in the isolates sampled. Further investigation is needed to resolve this hypothesis. Furthermore, although the SNP profiles of the isolates from Africa and Latin America are identical, they may have diverged substantially because the SNPs used can only determine node positions but not branch length caused by phylogenetic discovery bias (12).

Our SNP data clearly show that O139 Bengal was a derivative of the seventh pandemic, as previously suggested (13). Nine of the 12 O139 SNPs can now be seen to have arisen in its O1 precursor strain because they were present in seventh pandemic isolates as early as 1979 (online Technical Appendix). These SNPs also resolved the relationships of Groups IV–V. Some studies have suggested that the O139 variant may have multiple origins (14). However, our results suggest that these O139 isolates from the then new epidemic have a single origin, which is consistent with earlier ribotyping data (15).




Figure. Maximum parsimony tree of 68 seventh cholera pandemic and 3 pre-seventh cholera pandemic isolates. The tree was based on 18 N16961 seventh pandemic single-nucleotide polymorphisms (SNPs) and 12 MO10 O139 SNPs. The 3 pre-seventh pandemic isolates were used as an outgroup. Each strain name is followed by the year and location of isolation. All 15 O139 isolates had the same SNP profile and are shown as group VI. The numbers on each node represent the number of supporting SNPs. M821 and M819 from France and Germany are likely imported from either Africa or Asia. SNP data for the following isolates were obtained from GenBank: accession nos. RC9, ACHX00000000; MJ-1236, CP001385/CP001486; B33, ACHZ00000000; CIRS 101, ACVA00000000; MO10, AAKF00000000; N16961, AE003852; and M66–2, CP001233. Scale bar indicates number of nucleotide substitutions.

Our data show each of the groups/genotypes spread easily and widely to multiple countries or regions. This finding suggests that cholera epidemics or upsurges, which often occurred at the same time in many countries, were caused by the spread of newly arisen genotypes. Additionally, a genotype can also persist for long periods. Thus, in cholera-endemic regions such as Southeast Asia and Africa, cholera can be caused not only by an endemic genotype, but also by new epidemic genotypes. This finding is useful for control of cholera epidemics.

Acknowledgments

We thank Yi Ren and Gordon Stevenson for technical assistance. We also thank all strain donors, including J. Albert, A. Dodin, P. Eccheveria, J. Kaper, T. Popovic, B. Sack, C. Salles, W.C. Yam, and the reviewers for their helpful suggestions.

This research was supported by a Goldstar award from the University of New South Wales.

Ms Lam is a PhD student in medical microbiology at the University of New South Wales. Her research interests include molecular epidemiology and evolution of pathogenic bacteria.

References

1. Reeves PR, Lan R. Cholera in the 1990s. *Br Med Bull.* 1998;54:611–23.
2. Seas C, Miranda J, Gil AI, Leon-Barua R, Patz J, Huq A, et al. New insights on the emergence of cholera in Latin America during 1991: the Peruvian experience. *Am J Trop Med Hyg.* 2000;62:513–7.
3. Singh DV, Matte MH, Matte GR, Jiang S, Sabeena F, Shukla BN, et al. Molecular analysis of *Vibrio cholerae* O1, O139, non-O1, and non-O139 strains: clonal relationships between clinical and environmental isolates. *Appl Environ Microbiol.* 2001;67:910–21. DOI: 10.1128/AEM.67.2.910-921.2001
4. Lan R, Reeves PR. Pandemic spread of cholera: genetic diversity and relationships within the seventh pandemic clone of *Vibrio cholerae* determined by amplified fragment length polymorphism. *J Clin Microbiol.* 2002;40:172–81. DOI: 10.1128/JCM.40.1.172-181.2002
5. Feng L, Reeves PR, Lan R, Ren Y, Gao C, Zhou Z, et al. A recalibrated molecular clock and independent origins for the cholera pandemic clones. *PLoS One.* 2008;3:e4053. DOI: 10.1371/journal.pone.0004053
6. Heidelberg JF, Eisen JA, Nelson WC, Clayton RA, Gwinn ML, Dodson RJ, et al. DNA sequence of both chromosomes of the cholera pathogen *Vibrio cholerae*. *Nature.* 2000;406:477–83. DOI: 10.1038/35020000
7. Chun J, Grim CJ, Hasan NA, Lee JH, Choi SY, Haley BJ, et al. Comparative genomics reveals mechanism for short-term and long-term clonal transitions in pandemic *Vibrio cholerae*. *Proc Natl Acad Sci U S A.* 2009;106:15442–7. DOI: 10.1073/pnas.0907787106
8. Faruque SM, Tam VC, Chowdhury N, Diraphat P, Dziejman M, Heidelberg JF, et al. Genomic analysis of the Mozambique strain of *Vibrio cholerae* O1 reveals the origin of El Tor strains carrying classical CTX prophage. *Proc Natl Acad Sci U S A.* 2007;104:5151–6. DOI: 10.1073/pnas.0700365104
9. Tauxe RV, Blake PA. Epidemic cholera in Latin America. *JAMA.* 1992;267:1388–90. DOI: 10.1001/jama.267.10.1388
10. Evins GM, Cameron DN, Wells JG, Greene KD, Popovic T, Giono-Cerezo S, et al. The emerging diversity of the electrophoretic types of *Vibrio cholerae* in the Western Hemisphere. *J Infect Dis.* 1995;172:173–9.
11. Nusrin S, Gil AI, Bhuiyan NA, Safa A, Asakura M, Lanata CF, et al. Peruvian *Vibrio cholerae* O1 El Tor strains possess a distinct region in the *Vibrio* seventh pandemic island-II that differentiates them from the prototype pandemic El Tor strains. *J Med Microbiol.* 2009;58:342–54. DOI: 10.1099/jmm.0.005397-0
12. Pearson T, Busch JD, Ravel J, Read TD, Rhoton SD, U'Ren JM, et al. Phylogenetic discovery bias in *Bacillus anthracis* using single-nucleotide polymorphisms from whole-genome sequencing. *Proc Natl Acad Sci U S A.* 2004;101:13536–41. DOI: 10.1073/pnas.0403844101
13. Karaolis DK, Lan R, Reeves PR. The sixth and seventh cholera pandemics are due to independent clones separately derived from environmental, nontoxigenic, non-O1 *Vibrio cholerae*. *J Bacteriol.* 1995;177:3191–8.
14. Faruque SM, Saha MN, Asadulghani, Sack DA, Sack RB, Takeda Y, et al. The O139 serogroup of *Vibrio cholerae* comprises diverse clones of epidemic and nonepidemic strains derived from multiple *V. cholerae* O1 or non-O1 progenitors. *J Infect Dis.* 2000;182:1161–8. DOI: 10.1086/315807
15. Faruque SM, Abdul Alim AR, Roy SK, Khan F, Nair GB, Sack RB, et al. Molecular analysis of rRNA and cholera toxin genes carried by the new epidemic strain of toxigenic *Vibrio cholerae* O139 synonym Bengal. *J Clin Microbiol.* 1994;32:1050–3.

Address for correspondence: Ruiting Lan, University of New South Wales, School of Biotechnology and Biomolecular Sciences, Sydney, New South Wales 2052, Australia; email: r.lan@unsw.edu.au



Manage your email to focus on content of interest to you.

GovDelivery

www.cdc.eid/ncidod/eid/subscrib.htm

Appendix Table. Single nucleotide polymorphism profiles of 71 isolates of pandemic *Vibrio cholerae**

Group	SNP profile (no. isolates)	N16961 SNPs†*																MO10 SNPs†													
		vc 067	vc 083	vc 083	vc 098	vc 108	vc 109	vc 124	vca 094	vc 204	vc 208	vc 209	vc 267	vc 196	vca 024	vca 107	vc 032	vc 157	vc 189	vc 236	vc 256	vc 187	vc 095	vc 108	vc 186	vc 207	vc 131	vc 000	vc 084	vc 170	vc 259
Pre-7th	1 (2)	T	A	G	C	A	C	C	T	G	C	G	G	A	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
	2 (1)	T	A	G	C	A	C	C	T	G	C	G	G	A	C	C	C	C	A	A	C	C	C	C	T	C	A	G	G	G	C
I	3 (1)	G	G	A	T	T	T	T	C	A	T	T	A	A	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
	4 (13)	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
II	5 (17)	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
III	6 (7)	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
IV	7 (7)	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C
V	8 (1)	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	G	T	A	T	T	G	T	T	G	G	G	C
	9 (7)	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
VI	10 (15)	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T

*SNP, single-nucleotide polymorphism. Complete details for each strain and SNP profile can be found in the online Technical Appendix (www.cdc.gov/EID/content/16/7/1130-Techapp.pdf). SNP data for 7 isolates were obtained from GenBank (accession nos. RC9- ACHX00000000, MJ-1236- CP001385/CP001486, B33-ACHZ00000000, CIRS 101-ACVA00000000, MO10- AAKF00000000, N16961-AE003852, M66-2-CP001233).

†SNPs were selected from a comparison of N16961 with M66-2, and N16961 with MO10. The exact location of each SNP can be found in the online Technical Appendix. SNP mutations are shaded; ancestral SNPs have been left unshaded. The SNPs are grouped in the order in which the mutations are inferred to have occurred.

Evolution of Seventh Cholera Pandemic and Origin of 1991 Epidemic, Latin America

Technical Appendix

Methods

Primers and Location of Single Nucleotide Polymorphisms studied

The detection of each single nucleotide polymorphism (SNP) required 2 forward primers. The first forward primer contained the SNP of the seventh pandemic at the 3' end, while the second primer contained the SNP of either MO10 O139 Bengal or M66-2 pre-seventh pandemic. A complementary tail was added to the 5' end of each forward primer to form a hairpin structure. Alterations to the original primer sequence were also made to facilitate the folding of the primer into a hairpin structure. Table A-1 and Table A-2 contain the name, location, forward and reverse primer for each SNP. The same reverse primer was used for each pair of SNP reactions.

Hairpin Real-Time PCR (HP RT-PCR)

All RT-PCRs were carried out in a Rotor-Gene 6000 instrument (Corbett Life Science, Mortlake, New South Wales, Australia) with a 72-well rotor disk. Each RT-PCR reaction consisted of ≈ 100 ng DNA, 2.5 μmol each of forward and reverse primers and 5 μL SensiMix*Plus* SYBR Green (Quantace, Alexandria, New South Wales, Australia) (includes 2 \times Mix containing reaction buffer), Heat-Activated Taq DNA polymerase, dNTPs 6 mM MgCl₂, SYBR Green I. MilliQ water was added to adjust the final volume to 10 μL . The thermal cycling conditions were set up as follows: stage 1, 95°C for 10 min to activate Taq polymerase, stage 2, 95°C for 15 s, 69°C for 30 s, repeated 10 \times followed by stage 3, 95°C for 15 s, 60°C for 30 s, repeated 40 \times . On completion of each run, data were collected and analyzed with Rotor-Gene operating software v1.7.87 (Corbett Life Science). The fluorescent signal for each reaction was measured at the end of each cycle and plotted on a fluorescence curve. The cycle threshold (Ct) was set across the amplification curves during the exponential fluorescence phase.

Table A-1: Name, location and primers of 7th cholera pandemic and pre-7th pandemic single-nucleotide polymorphisms*

Locus	Gene	Annotation	Location in N16961	7th Pandemic		M66-2		Reverse primer (5'→3')
				Forward primer (5'→3')	SNP	Forward primer (5'→3')	SNP	
Large chromosome								
vc0329	<i>rpoC</i>	DNA-directed RNA polymerase, beta subunit	393808	atctttacc <u>T</u> GTCTCTGCAGCAGGTAAAGA <u>I</u>	T	gtctttacc <u>T</u> GTCTCTGCAGCAGGTAAAGA <u>C</u>	C	GAAGTATTGAGCTGGCATGTC
vc0672	<i>ptsP</i>	Phosphoenolpyruvate-protein phosphotransferase	765286	ttctctac <u>C</u> GCCTTGGCAGTAGAGAA <u>A</u>	A	ctctctac <u>C</u> GCCTTGGCAGTAGAGAG <u>G</u>	G	TCCCCATCGAGCTTTTTTA
vc0835	<i>tcpT</i>	Toxin co-regulated pilus biosynthesis protein T	943582	caaacattAGCCGA <u>C</u> GCCTAATGTTTT <u>G</u>	G	aaacattAGCCGA <u>C</u> GCCTAATGTTTT <u>I</u>	T	TGAGGTAGTTTTCGGTTCCACG
vc0837	<i>tcpF</i>	Toxin co-regulated pilus biosynthesis protein F	946533	ccactag <u>G</u> AACCATATCAGCCTAGTG <u>G</u>	G	tcactag <u>G</u> AACCATATCAGCCTAGTG <u>A</u>	A	GTATTTGACCACTTGTAAACCAT
vc0987	<i>hemH</i>	Ferrochelatase	1099038	ttgcagc <u>G</u> AGGAAGAGTGGCTGCA <u>A</u>	A	ctgcagc <u>G</u> AGGAAGAGTGGCTGCA <u>G</u>	G	TTTTAATGCCTTGGCGAG
vc1088		Sensor histidine kinase	1202761	agacaa <u>A</u> GCACTCGGGTGCCTTGTC <u>I</u>	T	ggacaa <u>A</u> GCACTCGGGTGCCTTGTC <u>C</u>	C	GCCTACAGCAGCATCAAAAAAT
vc1091	<i>oppA</i>	Oligopeptide ABC transporter, periplasmic oligopeptide-binding protein	1206798	agcggta <u>T</u> TCCGGAAATCACCGC <u>I</u>	T	tgcggt <u>A</u> TCCGGAAATCACCGC <u>A</u>	A	CTTCGCTTCTGCAATACGCTCT
vc1248		Methyl-accepting chemotaxis protein	1435093	aacgctc <u>A</u> ACCTATCATAACGAGCGT <u>I</u>	T	gacgctc <u>A</u> ACCTATCATAACGAGCGT <u>C</u>	C	GGCTCAGGCATGTTGTTGG
vc1579		Enterobactin synthetase component F-related protein	1770713	atcaccatc <u>T</u> GCCGTTAATTGATGGTG <u>A</u>	T	gtcaccatc <u>T</u> GCCGTTAATTGATGGTG <u>A</u>	C	TGCTTGTATATGTTGTGCCT
vc1898		Methyl-accepting chemotaxis protein	2128112	aatgat <u>C</u> AGTTTTGCGCTGATCAT <u>I</u>	T	gatgat <u>C</u> AGTTTTGCGCTGATCAT <u>C</u>	C	AATGCAGCGGTTGAAACACT
vc1967		Methyl-accepting chemotaxis protein	2201875	tgtttctt <u>G</u> CATCAACAATCAAGAA <u>A</u> C <u>A</u>	A	cgtttctt <u>G</u> CATCAACAATCAAGAA <u>A</u> C <u>G</u>	G	ACTTGCCTTGCCTTATCGTAGG
vc2046		Conserved hypothetical protein	2287749	aagcttt <u>G</u> TGGCCAGTGCAAAGCT <u>I</u>	T	gagcttt <u>G</u> TGGCCAGTGCAAAGCT <u>C</u>	C	AACCTTGAGTATCCTGTGG
vc2080		Transcriptional regulator, AraC/XylS family	2322891	agctctt <u>A</u> TCTCCATCCGAGTTAAGAGC <u>I</u>	T	cgctctt <u>A</u> TCTCCATCCGAGTTAAGAGC <u>G</u>	G	GCATTATCTAACGACGGA
vc2091	<i>sdhC</i>	Succinate dehydrogenase, cytochrome b556 subunit	2336169	actctc <u>T</u> GACAGGTGAGGAGAG <u>I</u>	T	tctctc <u>T</u> GACAGGTGAGGAGAG <u>A</u>	A	CAGCAATCGCATCCATCCT
vc2674	<i>hslU</i>	Protease HslVU, ATPase subunit HslU	2927259	tgccct <u>T</u> AACCTTGATGAAAGGC <u>A</u>	A	cgccct <u>T</u> AACCTTGATGAAAGGC <u>G</u>	G	TGTTGAAGTGACCCCGAA
Small chromosome								
vca0247		Transcriptional regulator, DeoR family	302905	actttgc <u>G</u> ATACAATCGAGCGCAAAG <u>I</u>	T	gctttgc <u>G</u> ATACAATCGAGCGCAAAG <u>C</u>	C	TCGGTTAGCCCTTTGCCAGA
vca0946	<i>malK</i>	Maltose/maltodextrin ABC transporter, ATP-binding protein	830773	gccgcta <u>G</u> AACCGTCCAA <u>T</u> AGCGG <u>C</u>	C	accgcta <u>G</u> AACCGTCCAA <u>T</u> AGCGG <u>I</u>	T	GTCGTACAAATTGAGGTGCGGG
vca1073		Bifunctional protein putA	956775	agtggt <u>C</u> GATGCAGATGTGGCACACT <u>I</u>	T	ggtggt <u>C</u> GATGCAGATGTGGCACACC <u>C</u>	C	ACTGCGTGTGCTGTTTGT

*SNP, single nucleotide polymorphism. Nucleotides in lower case indicate a complementary tail which was added to the primer to form a hairpin structure. Bold and italic nucleotides indicate that a deliberate change has been made to the sequence to facilitate the folding of the primer into a hairpin structure. Bold and underlined nucleotides indicate the corresponding SNP of either the 7th pandemic or M-662.

Table A-2: Name, location and primers of the *Vibrio cholerae* O139 MO10 single-nucleotide polymorphisms*

Locus	Gene	Annotation	Location in N16961	7th pandemic		MO10		Reverse primer (5'→3')
				Forward primer (5'→3')	SNP	Forward primer (5'→3')	SNP	
vc0008		Amino acid ABC transporter ATP binding protein	5414	cgataacc CGGTATGTTTTGGGTATCG	G	tgataacc CGGTATGTTTTGGGTATCA	A	GCGTGAACCTTTCTTGAGC
vc0847		Phage family integrase	913179	caacagc CTTGCCGTTTGGCTGTTG	G	taacagc CTTGCCGTTTGGCTGTTA	A	GCCATCGTGATTTTATTT
vc0959		Haemolysin (putative)	1024406	gccgaaAG TTCTTGGCGATCTTTCGGC	C	accgaaAG TTCTTGGCGATCTTTCGGT	T	GGTCCGAGTAGAAAGTCC
vc1082		Hypothetical protein	1149897	ggcctt CTTCTGGTTGAGAAGGCC	C	agcctt CTTCTGGTTGAGAAGGCT	T	AGATGGGCAAATACCTTA
vc1318	<i>ompV</i>	outer membrane protein OmpV	1401874	tggaac CAATATCGCCTGTGTTGCCA	A	aggcaac CAATATCGCCTGTGTTGCCT	T	TACCAGCAAGGGCACAATCA
vc1707		Hypothetical protein	1838824	cgtttga ACTGTCACATTCCAAACG	G	tgtttga ACTGTCACATTCCAAACA	A	AAACTTCGATAGCGTGAT
vc1865		Hypothetical protein	2005889	accagc AATTTAACTTGGCGCTGGT	T	cccagc AATTTAACTTGGCGCTGGG	G	CCCACCCCAAGGCAAGC
vc1877	<i>1pxk</i>	Tetra-acyldisaccharide 4'-kinase	2021732	gtcgtg GATGTTACCCACCACGAC	C	ttcgtg GATGTTACCCACCACGAA	A	TATCAAACGGGCGACAAA
vc2077		Ferrous iron transport protein B	2234253	gcacac CTCTGCATCAGGTGTGC	C	acacac CTCTGCATCAGGTGTGT	T	AAAGAAGCGGTTGTGGGG
vc2362		Threonine Synthetase	2518900	tttttc CGATTGTGCCGAAAAAG	A	cttttc CGATTGTGCCGAAAAAG	G	GGTCAAGCCGTTCCGCAA
vc2562	<i>cpdB</i>	Bi-functional 2' 3'- cyclic nucleotide 2'-phosphodiesterase/3' nucleotidase periplasmic precursor protein	2744542	gtgtacct GCGATCATCAAGGTACAC	C	atgtacct GCGATCATCAAGGTACAT	T	ACATCACGTCGTTGCTT
vc2599		Ribonuclease R	2766113	gtgaag GCTTGCTACGGCCTTCAC	C	atgaag GCTTGCTACGGCCTTCAT	T	CACCAACGAAATCAGAGT

*SNP, single nucleotide polymorphism. Nucleotides in lower case indicate a complementary tail which was added to the primer to form a hairpin structure.

Bold and italic nucleotides indicate that a deliberate change has been made to the sequence to facilitate the folding of the primer into a hairpin structure. Bold and underlined nucleotides indicate the corresponding SNP of either the 7th pandemic or MO10.

Table A-3: Single-nucleotide polymorphism (SNP) profiles of 71 isolates of pandemic *Vibrio cholerae*

Group	SNP profile	Isolate	Year	Location	N16961 SNPs																	MO10 SNPs												
					vc0672	vc0835	vc0837	vc0987	vc1088	vc1091	vc1248	vca0946	vc2046	vc2080	vc2091	vc2674	vc1967	vca0247	vca1073	vc0329	vc1579	vc1898	vc2363	vc2562	vc1877	vc0959	vc1082	vc1865	vc2077	vc1318	vc0008	vc0847	vc1707	vc2599
Pre-7th	1	M66-2†	1937	Indonesia	T	A	G	C	A	C	C	T	G	C	G	G	A	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M543	1938	Iraq	T	A	G	C	A	C	C	T	G	C	G	G	A	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
	2	M640	1954	Egypt	T	A	G	C	A	C	C	T	G	C	G	G	A	C	C	C	C	A	A	C	C	C	C	T	C	A	G	G	G	C
I	3	M793	1961	Indonesia	G	G	A	T	T	T	T	C	A	T	T	A	A	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M686	1968	Thailand	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M799	1989	Hong Kong	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M803	1961	Hong Kong	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M804	1962	India	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M805	1963	Cambodia	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M806	1964	India	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M807	1966	Vietnam	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M808	1969	Vietnam	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C	

II	5	M811	1971	Burma	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M815	1973	Philippines	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M820	1978	Malaysia	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M662	1993	Indonesia	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M663	1992	Indonesia	G	G	A	T	T	T	T	C	A	T	T	A	T	C	C	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M809	1970	Sierra Leone	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M821	1982	France	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M823	1984	Algeria	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M826	1990	Malawi	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M2314	1991	Peru	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M2315	1999	Brazil	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M2316	1998	Peru	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M829	1992	Malawi	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M830	1993	French Guiana	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M812	1971	Chad	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M817	1974	Chad	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M810	1970	Ethiopia	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M813	1972	Senegal	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M814	1972	Morocco	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
		M816	1974	Senegal	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C
M819	1975	Germany	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C		
M818	1975	Comoros Islands	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	C	C	G	A	C	C	C	C	T	C	A	G	G	G	C		
III	6	M650	1976	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C	
		M647	1970	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
		M795	1976	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
		M797	1986	Hong Kong	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
		N16961†	1971	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
		RC9†	1985	Kenya	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
		M825	1988	Zaire	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	T	A	A	C	C	C	C	T	C	A	G	G	G	C
IV	7	M646	1979	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
		M652	1981	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
		M714	1979	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
		M723	1982	Thailand	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
		M740	1985	Thailand	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
		M764	1989	Thailand	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
		M822	1983	Vietnam	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	T	C	A	G	G	G	C	
V	8	M654	1991	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	G	G	G	C	
		9	M791	1991	Thailand	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C

		M824	1987	Algeria	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
		MJ1236†	1994	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
		CIRS101	2002	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
		†																															
		B33†	2004	Mozambique	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
		M827	1990	Guinea	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
		M828	1991	Morocco	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	G	G	C
VI	10	M834	1993	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M833	1993	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M985	1992	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M987	1992	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M989	1993	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M988	1993	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M986	1992	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M984	1992	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M835	1993	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M537	1993	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M540	1993	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M542	1993	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M545	1993	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		M831	1993	Bangladesh	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T
		MO10†	1992	India	G	G	A	T	T	T	T	C	A	T	T	A	T	T	T	T	A	G	T	A	T	T	G	T	T	A	A	A	T

§ SNPs were selected from comparison between N16961 with M66-2 ,and N16961 with MO10. SNP mutations are shaded in blue, ancestral SNPs have been left unshaded. The SNPs are grouped in the order in which the mutations are inferred to have occurred. Horizontal lines separate individual SNP profiles

† SNP data for these isolates was obtained from GenBank.(accession nos: RC9- ACHX00000000, MJ-1236- CP001385/CP001486, B33- ACHZ00000000, CIRS 101- ACVA00000000, MO10- AAKF00000000, N16961- AE003852, M66-2- CP001233)