

# Worldwide Occurrence and Impact of Human Trichinellosis, 1986–2009

K. Darwin Murrell and Edoardo Pozio

## Medscape **ACTIVITY** EDUCATION

Medscape, LLC is pleased to provide online continuing medical education (CME) for this journal article, allowing clinicians the opportunity to earn CME credit.

This activity has been planned and implemented in accordance with the Essential Areas and policies of the Accreditation Council for Continuing Medical Education through the joint sponsorship of Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is accredited by the ACCME to provide continuing medical education for physicians.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1 *AMA PRA Category 1 Credit(s)*<sup>™</sup>. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 70% minimum passing score and complete the evaluation at [www.medscape.org/journal/eid](http://www.medscape.org/journal/eid); (4) view/print certificate.

**Release date: November 23, 2011; Expiration date: November 23, 2012**

### Learning Objectives

Upon completion of this activity, participants will be able to:

- Evaluate epidemiologic patterns of trichinellosis
- Analyze the clinical presentation of trichinellosis
- Distinguish the most common animal source of trichinellosis

### Editor

**Caran R. Wilbanks**, Technical Writer/Editor, *Emerging Infectious Diseases*. *Disclosure: Caran R. Wilbanks has disclosed the following relevant financial relationship: partner is employed by McKesson Corporation.*

### CME Author

**Charles P. Vega, MD**, Associate Professor; Residency Director, Department of Family Medicine, University of California, Irvine. *Disclosure: Charles P. Vega, MD, has disclosed no relevant financial relationships.*

### Authors

*Disclosures: K. Darwin Murrell, PhD; and Edoardo Pozio, PhD, have disclosed no relevant financial relationships.*

To assess the global incidence and clinical effects of human trichinellosis, we analyzed outbreak report data for 1986–2009. Searches of 6 international databases yielded 494 reports. After applying strict criteria for relevance and reliability, we selected 261 reports for data extraction. From 1986 through 2009, there were 65,818 cases and 42 deaths reported from 41 countries. The World Health Organization European Region accounted for 87% of cases; 50% of those occurred in Romania, mainly during 1990–1999. Incidence in the region ranged from 1.1 to 8.5 cases per 100,000 population. Trichinellosis affected primarily adults (median age 33.1 years) and about equally affected men (51%) and women. Major clinical effects, according to 5,377 well-described cases, were myalgia, diarrhea, fever, facial

edema, and headaches. Pork was the major source of infection; wild game sources were also frequently reported. These data will be valuable for estimating the illness worldwide.

Since the mid-19th century, trichinellosis has been a well-recognized meat-borne zoonosis; however, despite concerted control efforts, it remains a threat in many countries. Veterinary control over the slaughter of food animals to ensure food safety, particularly meat inspection, was introduced in Germany in 1866 specifically to prevent trichinellosis from pork infected with the muscle larvae of *Trichinella spiralis* (1). In the European Union, the estimated annual cost incurred from meat inspection of 167 million pigs (2) ranges from €25 million to €400 million (3). Even in countries without mandatory meat inspection (e.g., United States), the economic cost of selling pork in international and national markets is substantial (4).

Author affiliations: University of Copenhagen, Copenhagen, Denmark (K.D. Murrell); and Istituto Superiore di Sanità, Rome, Italy (E. Pozio)

DOI: <http://dx.doi.org/10.3201/eid1712.110896>

The epidemiology and systematics (i.e., the study of the diversification) of this zoonosis are now recognized to involve, in addition to *T. spiralis*, 7 other species in 4 genotypes, all of which are more commonly found in wild animals than in domestic pigs (5). *Trichinella* spp. have been found in domestic and wild animals in 66 countries (6). Human trichinellosis has been documented in 55 countries, particularly those with well-established food behavior that includes consuming meat dishes with raw or undercooked meat (6). Whether trichinellosis is a low-prevalence disease or is frequently misdiagnosed is not clearly understood; its detection can be difficult in low-level infections and its clinical manifestations overlap those of other diseases, such as influenza and chronic fatigue syndrome (7). Human infection is classically characterized by gastroenteritis; myalgia; malaise; facial edema; headache; subungual or conjunctival hemorrhages; and increased eosinophils, leukocytes, and muscle enzymes (7).

Reliable estimates of the incidence of trichinellosis among humans and its effect on health are not available; these estimates are necessary for setting priorities. In the 1990s, the global prevalence of trichinellosis was  $\approx 10$  million, and a recent incidence estimate suggested  $\approx 10,000$  infections per year (6). However, because of problems related to incomplete data from some regions and to the quality of diagnostic criteria for infection, the Foodborne Disease Burden Epidemiology Reference Group of the World Health Organization (WHO) requested a systematic review of the global incidence, burden of disease, and major sources of infection that used strict criteria for data selection and extraction. Our analyses and summaries of the epidemiologic and clinical data selected provide a basis for an assessment of trichinellosis as a public health problem.

### Data Sources and Selection Criteria

We retrospectively reviewed trichinellosis outbreak investigations conducted worldwide during 1986–2009. The data analysis focused on incidence, age and sex of patients, infection rates, major clinical aspects including sequelae, and meat sources of infection. The database we developed was geographically organized according to the WHO regions ([www.who.int/choice/demography/regions/en](http://www.who.int/choice/demography/regions/en)): African Region, 46 countries; Region of the Americas, 12 countries; Eastern Mediterranean Region, 22 north African and Middle Eastern countries; European Region, 44 European and 6 Asian countries; South-East Asian Region, 11 east Asian countries; and Western Pacific Region, 27 countries. Data searches of literature included PubMed, Centres for Agricultural Bioscience International (CABI) abstracts, WHO library, System for Information on Gray Literature, Pan America Health Organization Virtual Library, and Index Medicus for South-East Asian Region.

The search terms used were trichinosis, trichinellosis, trichinelliasis, and trichinella. These were combined with the terms prevalence, outbreaks, epidemiology, clinical symptoms, and duration. The search terms were also combined with pork, pig, wild boar, wild pig, warthog, horse, badger, jackal, cougar, walrus, armadillo, turtle, and bear meat. Published abstracts were screened for retention by using the criteria of relevance to human outbreaks or single cases occurring from 1986 through 2009 and by determining whether the report was based on original data (primary source or unpublished data managed by national government agencies). The full paper versions of selected abstracts were then obtained where possible and further screened and evaluated. Outbreak reports published  $>1$  time were occasionally encountered, and care was taken to prevent duplication of data in the database; preference was given to published international, peer-reviewed versions. In some instances, data were obtained through contact with scientists in countries of interest who had access to unpublished and detailed information about outbreaks; these sources are indicated in the reference lists in the online Technical Appendix ([wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf)). In addition, information about isolated outbreaks maintained by a national health system was obtained for some countries through personal contacts (see Acknowledgments). For reports in which English versions were not available, translations were obtained through the generous help of colleagues (see Acknowledgments).

### Definitions

Reports, published or unpublished, were excluded from the database if the diagnosis of *Trichinella* spp. infection was not based on a diagnostic procedure that we defined as confirmatory. Although direct demonstration of muscle larvae infection in biopsy samples is now infrequent, reliance solely on serologic testing to confirm infection can be problematic (8). Therefore, a serologically positive case was included in the database only if the sample was confirmed by a Western blot test or if the patient's illness could be classified as highly probable according to the clinical diagnostic algorithm published by Dupouy-Camet and Bruschi (7), in which a patient with a positive serologic test result must also exhibit  $\geq 1$  classical trichinellosis signs and symptoms (e.g., myalgia, facial edema, headaches, diarrhea, eosinophilia).

We took a conservative approach in extracting clinical data for the analysis of frequency of major signs and symptoms in patients with *Trichinella* spp. infections and excluded report data if there was lack of clarity and reliability of clinical procedures and laboratory tests used to confirm infection. However, in some instances of inadequate clinical descriptions, the report was retained if it otherwise presented useful epidemiologic data.

Consequently, the total number of human infections (Tables 1–3) exceeds the number of cases used to summarize the frequency of major signs and symptoms (Table 4). Except in rare cases, clinical data were extracted from outbreak reports only if the data were from multiple cases that met these criteria; exceptions were reports of single cases from countries with rare occurrences of trichinellosis but that had good clinical and laboratory confirmation data (e.g., Korea, Japan, India).

From the original 494 abstracts identified from literature searches, 378 were judged to potentially meet the criteria for data extraction, and full articles were obtained for most abstracts. From these, 261 reports were retained for data extraction and inclusion. A major reason for rejection of articles was failure to meet the criteria for confirmation of infection, especially interpretation of serologic results.

### Incidence

In Table 2, the incidence (100,000 person-years) is reported for specific periods because the data were obtained over a shorter period than the formal study interval (1986–2009). For some countries, when incidence figures were not reported in published papers or national health reports, we calculated incidence from data available in the referenced reports by using the WHO World Population Prospects (the 2008 revision) (<http://esa.un.org/wpp/unpp/p2k0data.asp>). Overall, from 1986 through 2009, there were 65,818 cases and 42 deaths from trichinellosis reported from 41 countries (Tables 1–3). The European Region accounted for 86% of cases (56,912), of which 28,564 (50%) occurred in Romania, mainly during 1990–1999. The full references for specific country reports summarized in Tables 2–5 are available in the online Technical Appendix.

Of 46 countries in the African Region, trichinellosis has been documented only among soldiers in the Gojjam region and policemen in the Arsi region of Ethiopia, a country where the Christian population accounts for ≈60% of the total population. In the Eastern Mediterranean Region, trichinellosis was documented only in the Christian population of Lebanon and in Iran from the consumption of wild boar meat (Table 2). In Algeria and Senegal, where most of the population is Muslim, trichinellosis has been documented only in Europeans (6).

In the European Region, 4 epidemiologic patterns are discernable: 1) countries of eastern Europe where incidence rates are >1 case/100,000 inhabitants (Bosnia-Herzegovina, 4.1; Bulgaria, 2.4–2.9; Croatia, 1.7–4.8; Latvia, 1.1–1.3; Lithuania, 1.2–6.6; Romania, 2.9–8.5; and Serbia, 5.0); 2) countries with a low number of inhabitants where the occurrence of a large outbreak results in a high incidence rate (e.g., Israel, 3.0; Slovakia, 6.2; and Slovenia, 10.5); 3) 19 countries with a low incidence rate caused either by sporadic infections or by a large general population that reduces the incidence per 100,000 inhabitants even when a large outbreak occurs; and 4) 21 countries where no autochthonous infections were reported during the period. Incidence in eastern Europe spiked during the late 1980s and early 1990s and then decreased over the past decade. This pattern may be linked to the political, social, and economic changes that occurred with the breakup of the former Soviet Union as described by Djordjevic et al. (9). The gradual restoration of infrastructure related to food safety (e.g., meat inspection, pig production management, veterinary services) probably contributed substantially to the decrease in incidence in these countries.

The number of cases in the Region of the Americas was comparatively low (Table 1), except in Argentina (Table 2). National incidence estimates are limited for Region of the Americas countries and published only for the United States, Chile, and Argentina; data from Canada, Mexico, and Argentina pertain only to selected states, provinces, or districts that had large outbreaks. In Canada during the period, a few large outbreaks in northern communities among native people who consumed wild game accounted for most of the outbreaks in the country. As an example of the problem of informal or clandestine meat transportation, 2 outbreaks occurred among foreign hunters; in 1 outbreak, the hunters transported infected bear home (France) and unknowingly exposed friends and family (17 total cases). For Canada and Greenland, trichinellosis was caused by consumption of wild game harboring *T. nativa*, which does not infect swine; no pork-transmitted *Trichinella* spp. have been recorded in Canada for many years.

The risk for trichinellosis has decreased markedly in the United States and Chile since the 1990s. The large number of cases associated with Argentina contrasts with

Table 1. Clinically confirmed cases of trichinellosis in humans documented in World Health Organization regions, 1986–2009

Region (no. countries)	No. (%) countries with trichinellosis	No. (%) documented human infections	No. (%) deaths
African Region (46)	1 (2.17)	28 (0.04)	1 (3.57)
Region of the Americas (12)	5 (42.67)	7,179 (10.90)	10 (0.10)
Eastern Mediterranean Region (22)	2 (9.09)	50 (0.07)	0
European Region (50)	29 (58.00)	56,912 (86.47)	24 (0.04)
South-East Asian Region (11)	1 (9.69)	219 (0.33)	1 (0.50)
Western Pacific Region (27)	3 (11.11)	1,344 (2.04)	6 (0.40)
Other*	NA	86 (0.13)	0
Total (168)	41 (24)	65,818 (100.00)	42 (0.40)

\*Infections acquired in countries other than the one in which diagnosis occurred. NA, not applicable

the situation among other countries in South America. The cases in Argentina may be related to the European origins of persons immigrating there and the risky food behavior

they brought with them (10). Although incidence data from Mexico and Argentina are limited, trichinellosis outbreaks are reported frequently in Argentina from domestic pork,

Table 2. Total cases and incidence of *Trichinella* spp. infections, by World Health Organization region and country, 1986–2009\*

Region/country	Years	No. cases	Average incidence†
African Region, Ethiopia	1986	8	0.02
	1990	20	0.04
Region of the Americas		7,179	
Argentina	1990–2005	5,221	1.48
Canada	1987–2009	257	0.03
Chile	1991–2004	698	0.36
Mexico	1986–2001	351	0.02
United States	1987–2007	652	0.016–0.004
Eastern Mediterranean Region		50	
Iran	2007	6	0.008
Lebanon	1995	44	1.25
European Region		56,912	
Belarus	1988, 1989	16	0.08, 0.55
Bosnia and Herzegovina	1993–2003	1,600	0.1–8.0
Bulgaria	1990–2006	4,108	2.9
Croatia	1994–2009	2,110	0.02–12.3
Czech Republic	1986–2009	31	0.01
Estonia	1986–2009	91	0.0–2.9
France	1986–2009	1,203	0.00–0.95
Georgia	1988	3	0.05
Germany	1986–2009	185	0.00–0.01
Greece	2009	1	0.008
Hungary	1986–2009	158	0.18–0.057
Ireland	2007	2	0.04
Israel	2002, 2004	230	0.5, 3.0
Italy	1986–2009	1,181	0.0–0.9
Kyrgyzstan	1996	10	0.2
Latvia	1986–2009	636	0.07–3.8
Lithuania	1989–2009	3,979	0.4–21.8
Macedonia	1992	6	0.3
Poland	1986–2007	3,084	0.05–1.5
Romania	1986–2007	28,564	1.7–16.1
Russia	1996–2002	971	0.3–0.6
Serbia	1994–2003	5,210	1.8–7.8
Slovakia	1986–2008	440	0.0–6.2
Slovenia	1989–2006	203	0.00–10.5
Spain	1986–2009	1,244	0.0–0.4
Switzerland	1994, 2009	4	0.01, 0.04
Turkey	2003, 2004	425	0.01, 0.59
United Kingdom	1999	7	0.01
Ukraine	1986–2009	1,210	0.00–0.30
South-East Asian Region		219	
India	1996–2002	3	0.0003
Thailand	1993–2007	216	0.35
Western Pacific Region		1,344	
Japan	1999–2005	4	NA
South Korea	1999–2003	8	0.016
Laos	2004–2006	123	2.09
People's Republic of China	1995–2009	1,137	NA
Singapore	1998	25	0.64
Vietnam	2001–2004	47	0.058

\*The detailed data and references for each country are available in the online Technical Appendix ([wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf)), section A. NA, insufficient data for incidence calculation.

†Incidence/100,000 person-years. For some countries, incidence was not reported and was calculated from data available in the report referenced.



Table 3. Trichinellosis acquired in locations different from those where the disease developed and was diagnosed, 1986–2009\*

Country where infection developed and was diagnosed	Country where infection was acquired (no. clinical cases)
Austria	Yugoslavia (10)
Belgium	Canada (1)
Czech Republic	Poland (2), Ukraine (2), France (1)
Denmark	Poland (12)
France	Algeria (6), Cameroon (1), Canada (13), Croatia (1), Greenland (2), Kenya (2), Laos (5), Senegal (5), Spain (1), Thailand (1), Turkey (3), Yugoslavia (1)
Germany	Canada (1), Poland (3)
Italy	Romania (4)
The Netherlands	Yugoslavia (3), Montenegro (5)
Hong Kong	Canada (1)

\*Complete data and references are available in the online Technical Appendix ([wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf)), section B.

indicating that a substantial pig husbandry risk persists in that country.

The Asian countries of the Western Pacific Region and the South-East Asian Region reported few outbreaks during the period (Tables 1, 2). Although large outbreaks in the People's Republic of China have been reported (11), the criteria for selection of reports and data extraction eliminated some reports because of insufficient diagnostic detail to meet the confirmation criteria. Most of the outbreaks reported from Thailand, Laos, and Vietnam, occurred in the northern mountainous regions among the indigenous people who practice free-roaming pig husbandry (12–14). After a 30-year period of no reports of trichinellosis cases, Laos recently experienced several outbreaks (12). The estimated incidence in rural areas of that country is high, which suggests a possible emerging problem there.

Globally, reporting of trichinellosis varies greatly. A major factor affecting the collection of epidemiologic and clinical data is an absent or inadequate national reporting system. For example, in some countries of eastern Europe (e.g., Bosnia-Herzegovina, Byelorussia, Georgia, Moldavia,

Romania, Russia, Ukraine) trichinellosis occurs frequently in villages during the winter, and infection might not be diagnosed and subsequently reported unless infection is sufficiently severe to require hospitalization or the cases are part of a larger outbreak that requires attention from public health authorities (A. Marinculic, M.C. Cretu, W. Kociecka, N. Iashvili, N. Bogatko, pers. comm.). For example, in Romania, most of the 20,059 cases documented during 1990–1999 pertain to hospitalized persons only. However, for each hospitalized person, there are probably others in whom a moderate or mild infection developed that did not justify the travel and costs that would be incurred in seeking attention for diagnosis and treatment. Consequently, they are not usually officially recorded as having trichinellosis (M.C. Cretu, pers. comm.). In countries where most of the population is Muslim, *Trichinella* spp. infection is rare and may not be reported at all because of a scarcity of physicians, lack of good diagnostic tools, and occurrence in remote areas. In contrast, in industrialized countries such as those of Western Europe, United States, and Canada, nearly all cases are more likely to be detected and recorded, including asymptomatic cases associated with large outbreaks. For these reasons, the data we present may underrepresent the incidence of trichinellosis in lesser developed countries in comparison to that in industrialized and affluent countries.

### Sex- and Age-specific Infection

Data from clinical reports (Table 5) demonstrate that trichinellosis is a disease primarily of adults, occurring about equally among both sexes (2,631 [51%] of 5,154 infections occurred in male patients). Infection in male patients did occur more frequently, however, in Ethiopia (100%), Vietnam (91%), Japan and South Korea (75%), Thailand (64%), and China (57%). Age-specific infection data (Table 5) show the highest proportion of cases, for both sexes, was among persons 20–50 years of age (median 33.1 years). Data on age-specific prevalence rates were rarely reported; however, recent improvements in diagnosis of trichinellosis, particularly immunodiagnostic methods, may encourage more human prevalence surveys

Table 4. Frequency of major clinical signs associated with trichinellosis among World Health Organization regions, 1986–2009\*

Region	Total no. cases†	Clinical sign, no. cases						
		Diarrhea	Myalgia	Fever	Facial and/or eyelid edema	Headache	Eosinophilia	Deaths
African Region	28	28	8	11	8	6	6	0
Region of the Americas	1,229	400	969	821	790	410	606	10
Eastern Mediterranean Region	45	43	42	41	Not reported	30	0	4
European Region	3,118	798	1,971	1,387	1,617	351	1,850	24
South-East Asian Region	210	82	206	103	102	71	97	1
Western Pacific Region	747	79	409	474	429	104	180	8
Total no. (%)	5,377 (100.0)	1,430 (27.0)	3,605 (67.0)	2,837 (53.0)	2,946 (55.0)	972 (18.0)	2,739 (51.0)	35 (1.0)

\*Report references are available in the online Technical Appendix ([wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf)), section C.

†Cases included in this table were selected from all reports on the basis of detailed descriptions of clinical data in the reports.

Table 5. Demographic data on trichinellosis patients, by World Health Organization region and country, 1986–2009\*

Region/country	% Male patients (total no. cases)†	Age, y, of infected persons (no. cases)
African Region: Ethiopia	100 (28)	Range 23–25 (3); mean 24 (3)
Region of the Americas		
Canada	62.1 (150)	Range 21–67 (85); mean 34.4 (65)
Chile	60 (667)	Range 5–70 (667)
Mexico	35 (59)	Range 25–44 (59)
United States	57.5 (632)	Range 1–87 (412); mean 42.0 (126); median 37.1 (232)
Eastern Mediterranean Region: Lebanon	54 (44)	Range 10–70 (44); mean 33 (44)
European Region		
Bulgaria	49 (228)	Range 1–70 (228)
Croatia	57 (200)	Range 3–67 (200); mean 35 (200)
Czech Republic	41.9 (31)	Range 9–68 (31); mean 35.9 (31)
France	51.4 (586)	Range 1–84 (581); mean 43.8 (581)
Germany	51.9 (104)	Range 1–73 (101); mean 34.8 (101)
Israel	100 (26)	Mean 32 (26)
Italy	50.3 (382)	Range 1–90 (368); mean 36.7 (368)
Romania	53.2 (521)	Range 1–>60 (521); mean 31.4 (521)
Slovakia	63.6 (11)	Range 16–80 (21); mean 40.5 (21)
Spain	57.5 (237)	Range 2–86 (140); mean 40.7 (177)
Turkey	52.6 (418)	Range 1.5–73 (418); mean 31.1 (418)
South-East Asian Region: Thailand	71 (165)	Range 7–70 (210); mean 35.6 (208); median 34.5 (140)
Western Pacific Region		
Laos	47 (111)	Range 5–69 (111); mean 30.4 (90); median 34 (21)
People's Republic of China	58.2 (802)	Range 1–90 (482)
Vietnam	92 (42)	Range 20–60 (42); mean 45.4 (42)
Singapore	56 (25)	Mean 22.5 (25)

\*Clinical details and report references are available in the online Technical Appendix ([wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf)), section D.

†Data are from reports that presented adequate sex and age data on ≥10 cases during 1986–2009.

and surveillance for trichinellosis that could yield better information about sex- and age-specific rates.

Although infections also occur in children and teenagers, the predominance of infection in adults probably results from culture-driven food behavior. Improperly cooked or prepared meat dishes may be more commonly eaten at adult-oriented events, particularly if alcohol is consumed. There are only a few published studies on the link between food behavior and trichinellosis (15,16), but this potential behavioral risk factor is similar to that that occurs in other foodborne parasites, such as fish-borne parasites (17).

### Clinical Signs and Sequelae

For 5,377 cases, the chief clinical signs of trichinellosis were compatible in type and frequency with the classical trichinellosis syndrome (7), i.e., myalgia, diarrhea, fever, facial edema, and headaches that, after treatment, disappeared within 2–8 weeks (Table 4). Their rapid recovery reflects improvements in diagnostic methods, drug therapy, and public health education. The more rapid diagnosis and treatment in recent decades may also account for the low death rate; 42 deaths occurred worldwide during the 24-year period. Determining the disease burden of trichinellosis, however, is hampered by lack of data on the long-term sequelae of infection; few clinical reports included posttreatment follow-up evaluations, particularly

beyond 1 month. The few studies that included follow-up over a longer time span indicate that myalgia and fatigue can persist for 4 months and, in a substantial proportion of cases, for up to 2 years (18–20). There is a need for internationally recognized epidemiologic and clinical protocols for trichinellosis outbreaks that include follow-up investigations that would facilitate reliable calculations of disease estimates.

### Sources of Infection

Domestic pigs and wild boars were the major sources of *Trichinella* spp. infection for humans, but in recent years new infection sources, particularly from exotic hosts, have emerged (Table 6). An example is the cause of outbreaks in France, where in addition to wild boar sources, most trichinellosis cases for the past 2 decades have resulted from consumption of raw horse meat, a strong food preference in French culture (21). In Italy, human infections from consumption of horse meat have also been documented in 2 areas (Emilia Romagna and Lombardy regions in northern Italy and the Apulia region in southern Italy), where the French fondness for raw horse meat was introduced centuries ago (16). In China and the Slovak Republic, dog meat was the source of infection in several outbreaks (22,23). Although Judaic and Muslim religions forbid the consumption of pork, in Israel, Lebanon, and

## SYNOPSIS

Table 6. Types of meat linked to trichinellosis cases and outbreaks in the world, by World Health Organization region and country, 1986–2009\*

Region/country	Meat source, % cases or outbreaks		
	Domestic pig	Wild game	Other
African Region: Ethiopia	0	100	0
Region of the Americas			
Argentina, Chile	100	0	0
Canada	0	100	0
United States	57	43	0
Mexico	86	0	14 (horse)
Eastern Mediterranean Region: Iran and Lebanon	0	100	0
European Region			
Belarus, Croatia, Georgia, Macedonia, Serbia, United Kingdom	100	0	0
Estonia, Turkey, Ukraine	50	50	0
France	0	65	35 (horse)
Germany	83	17	0
Greece, Israel	0	100	0
Hungary	52	48	0
Italy	38	38	24 (horse)
Lithuania	48	52	0
Poland	41	59	0
Romania	95	5	0
Slovakia	50	25	25 (dog)
Spain	60	40	0
South-East Asian Region: Thailand	50	50	0
Western Pacific Region			
People's Republic of China	86	13	1 (dog and others)
Japan	25	75	0
South Korea	0	100	0
India, Laos, Papua New Guinea	50	50	0
Singapore, Vietnam	100	0	0

\*Data for each country aggregated from our database of studies. Data are from reports cited in the online Technical Appendix ([wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0896-Techapp.pdf)), section A.

Syria human outbreaks of trichinellosis have occurred after consumption of meat from wild boars among the Christian Arab population and immigrant laborers (24–27). Muslim populations are not entirely protected from acquiring trichinellosis, however, as demonstrated by a large outbreak in Turkey from the consumption of minced beef illegally mixed with pork of unknown origin (Table 2) (28).

The demographic movements of humans with culturally unique food practices, the illegal importation of uncontrolled meat from trichinellosis-endemic to -nonendemic countries, and the introduction of risky new food practices have resulted in cases in Denmark, Germany, Italy, Spain, Sweden, and the United Kingdom (Tables 2, 3) (29–33). Many cases of trichinellosis have occurred among international travelers who acquired *Trichinella* spp. infections while visiting or hunting in disease-endemic areas and the disease developed after they returned to their home countries (Table 3) (34–38).

### Issues Affecting the Effective Control of Trichinellosis

Human behavior is the biggest determinant in the persistence of trichinellosis in the face of increasing regulations directed at ensuring the safety of meat and the

enhancement of good management practices in farming, especially in areas in which trichinellosis is highly endemic, such as the European and the Americas regions. Of particular concern is an increase in the association of wild animals with domestic pigs. For example, in the United States, the expansion of the range of feral pigs (wild boars) into major areas of pig production, including free-range systems, may increase the risk for incursion of *Trichinella* spp. into the human food chain (39). The increased frequency of outbreaks from eating pork from wild boars in Europe is believed to be related to the great increase in wild boar populations (40). As with other foodborne zoonoses, cultural traditions in food behavior and practices in the use of domestic and wild animals are not easily altered, and trichinellosis can be expected to remain a food-safety risk in many areas of the world for the foreseeable future.

### Acknowledgments

We thank the following persons for their willingness to provide articles, national data on human trichinellosis incidence, or translations for key articles and abstracts: N. Akkoc, H. Auer, R. Beck, S. Boutsini, J.Y. Chai, G. Deksne, Do Dung, P. Dubinsky, J. Dupouy-Camet, J. Epshtein, T. Garate, E. Golab,

M.A. Gómez Morales, B. Gottstein, L. Kolarova, T. Kortbeek, A. Malakauskas, Y. Nawa, Nguyen De, K. Nöckler, Ming-Bao Quiang, M. Ribicich, and J. Waikagul.

Dr Murrell is honorary professor at the Faculty of Life Sciences, University of Copenhagen, Denmark, and primarily associated with the WHO/Food and Agricultural Organization of the United Nations Collaborating Centre for Parasitic Zoonoses. His primary research interests are meat-borne and fish-borne parasitic zoonoses.

Dr Pozio is a parasitologist in the Department of Infectious, Parasitic and Immunomediated Diseases at Istituto Superiore di Sanità, Rome, Italy. His research interests include foodborne parasitic zoonoses, particularly trichinellosis, echinococcosis, opisthorchiasis, cryptosporidiosis, and giardiasis.

## References


- Campbell WC. Historical introduction. In: Campbell WC, editor. *Trichinella* and trichinosis. New York: Plenum Press; 1983. p. 1–30.
- Alban L, Pozio E, Boes J, Boireau P, Boué F, Claes M, et al. Towards a standardised surveillance for *Trichinella* in the European Union. *Prev Vet Med*. 2011;99:148–60. doi:10.1016/j.prevetmed.2011.02.008
- Kapel CM. Changes in the EU legislation on *Trichinella* inspection—new challenges in the epidemiology. *Vet Parasitol*. 2005;132:189–94. doi:10.1016/j.vetpar.2005.05.055
- Pyburn DG, Gamble HR, Wagstrom EA, Anderson LA, Miller LE. Trichinae certification in the United States pork industry. *Vet Parasitol*. 2005;132:179–83. doi:10.1016/j.vetpar.2005.05.051
- Pozio E, Hoberg E, La Rosa G, Zarlenga DS. Molecular taxonomy, phylogeny and biogeography of nematodes belonging to the *Trichinella* genus. *Infect Genet Evol*. 2009;9:606–16. doi:10.1016/j.meegid.2009.03.003
- Pozio E. World distribution of *Trichinella* spp. infections in animals and humans. *Vet Parasitol*. 2007;149:3–21. doi:10.1016/j.vetpar.2007.07.002
- Dupouy-Camet J, Bruschi F. Management and diagnosis of human trichinellosis. In: Dupouy-Camet J, Murrell KD, editors. *FAO/WHO/OIE guidelines for the surveillance, management, prevention and control of trichinellosis*. Paris: World Organisation for Animal Health; 2007. p. 37–68.
- Gómez-Morales MA, Ludovisi A, Amati M, Cherchi S, Pezzotti P, Pozio E. Validation of an ELISA for the diagnosis of human trichinellosis. *Clin Vaccine Immunol*. 2008;15:1723–9. doi:10.1128/CVI.00257-08
- Djordjevic M, Bacic M, Petricevic M, Cuperlovic K, Malakauskas A, Kapel CM, et al. Social, political, and economic factors responsible for the reemergence of trichinellosis in Serbia: a case study. *J Parasitol*. 2003;89:226–31. doi:10.1645/0022-3395(2003)089[0226:SPA EFR]2.0.CO;2
- Pozio E, Murrell KD. Systematics and epidemiology of *Trichinella*. *Adv Parasitol*. 2006;63:367–439. doi:10.1016/S0065-308X(06)63005-4
- Cui J, Wang ZQ, Xu BL. The epidemiology of human trichinellosis in China during 2004–2009. *Acta Trop*. 2011;118:1–5. doi:10.1016/j.actatropica.2011.02.005
- Barennes H, Sayasone S, Odermatt P, De Bruyne A, Hongskhone S, Newton PN, et al. A major trichinellosis outbreak suggesting a high endemicity of *Trichinella* infection in northern Laos. *Am J Trop Med Hyg*. 2008;78:40–4.
- Kaewpitoon N, Kaewpitoon SJ, Pengsaa P. Food-borne parasitic zoonosis: distribution of trichinosis in Thailand. *World J Gastroenterol*. 2008;14:3471–5. doi:10.3748/wjg.14.3471
- Vu Thi N, Dorny P, La Rosa G, To Long T, Nguyen Van C, Pozio E. High prevalence of anti-*Trichinella* IgG in domestic pigs of the Son La province, Vietnam. *Vet Parasitol*. 2010;168:136–40. doi:10.1016/j.vetpar.2009.10.024
- Blaga R, Durand B, Antoniu S, Gherman C, Cretu CM, Cozma V, et al. A dramatic increase in the incidence of human trichinellosis in Romania over the past 25 years: impact of political changes and regional food habits. *Am J Trop Med Hyg*. 2007;76:983–6.
- Gottstein B, Pozio E, Nöckler K. Epidemiology, diagnosis, treatment, and control of trichinellosis. *Clin Microbiol Rev*. 2009;22:127–45. doi:10.1128/CMR.00026-08
- Chai JY, Murrell KD, Lymbery AJ. Fish-borne parasitic zoonoses: status and issues. *Int J Parasitol*. 2005;35:1233–54. doi:10.1016/j.ijpara.2005.07.013
- Jongwutiwes S, Chantachum N, Kraivichian P, Siriyasatien P, Putapornpit C, Tamburrini A, et al. First outbreak of human trichinellosis caused by *Trichinella pseudospiralis*. *Clin Infect Dis*. 1998;26:111–5. doi:10.1086/516278
- Watt G, Saisorn S, Jongsakul K, Sakolvaree Y, Chaicumpa W. Blind-ed, placebo-controlled trial of antiparasitic drugs for trichinosis myositis. *J Infect Dis*. 2000;182:371–4. doi:10.1086/315645
- Calcagno MA, Teixeira C, Forastiero MA, Costantino SN, Venturriello SM. Clinical, serological and parasitological aspects of an outbreak of human trichinellosis in Villa Mercedes, San Luis, Argentina. The acute and chronic phases of the infection [in Spanish]. *Medicina (B Aires)*. 2005;65:302–6.
- Boireau P, Vallée I, Roman T, Perret C, Mingyuan L, Gamble HR, et al. *Trichinella* in horses: a low frequency infection with high human risk. *Vet Parasitol*. 2000;93:309–20. doi:10.1016/S0304-4017(00)00348-4
- Dubinský P, Stefániková A, Kinčeková J, Ondriska F, Reiterová K, Medvedová M. Trichinellosis in the Slovak Republic. *Parasite*. 2001;8(2 Suppl):S100–2.
- Liu M, Boireau P. Trichinellosis in China: epidemiology and control. *Trends Parasitol*. 2002;18:553–6. doi:10.1016/S1471-4922(02)02401-7
- Haim M, Efrat M, Wilson M, Schantz PM, Cohen D, Shemer J. An outbreak of *Trichinella spiralis* infection in southern Lebanon. *Epidemiol Infect*. 1997;119:357–62. doi:10.1017/S0950268897007875
- Hefer E, Rishpon S, Volovik I. Trichinosis outbreak among Thai migrant workers in the Hadera sub-district [in Hebrew]. *Harefuah*. 2004;143:656–60, 694.
- Marva E, Markovics A, Gdalevich M, Asor N, Sadik C, Leventhal A. Trichinellosis outbreak. *Emerg Infect Dis*. 2005;11:1979–81.
- Olaison L, Ljungström I. An outbreak of trichinosis in Lebanon. *Trans R Soc Trop Med Hyg*. 1992;86:658–60. doi:10.1016/0035-9203(92)90178-F
- Akkoc N, Kuruuzum Z, Akar S, Yuce A, Onen F, Yapar N, et al. A large-scale outbreak of trichinellosis caused by *Trichinella britovi* in Turkey. *Zoonoses Public Health*. 2009;56:65–70. doi:10.1111/j.1863-2378.2008.01158.x
- Milne LM, Bhagani S, Bannister BA, Laitner SM, Moore P, Eza D, et al. Trichinellosis acquired in the United Kingdom. *Epidemiol Infect*. 2001;127:359–63. doi:10.1017/S0950268801005994
- Pozio E, Marucci G. *Trichinella*-infected pork products: a dangerous gift. *Trends Parasitol*. 2003;19:338. doi:10.1016/S1471-4922(03)00138-7
- Gallardo MT, Mateos L, Artieda J, Wesslen L, Ruiz C, García MA, et al. Outbreak of trichinellosis in Spain and Sweden due to consumption of wild boar meat contaminated with *Trichinella britovi*. *Euro Surveill*. 2007;12:E070315.1.



## SYNOPSIS

32. Nöckler K, Wichmann-Schauer H, Hiller P, Müller A, Bogner K. Trichinellosis outbreak in Bavaria caused by cured sausage from Romania, January 2007. *Euro Surveill.* 2007;12:E070823.2.
33. Stensvold CR, Nielsen HV, Mølbak K. A case of trichinellosis in Denmark, imported from Poland, June 2007. *Euro Surveill.* 2007;12:E070809.3.
34. Dupouy-Camet J, Allegretti S, Truong TP. Enquete sur l'incidence de la trichinellose en France (1994–1995). *Bulletin Épidémiologique Hebdomadaire.* 1998;28:122–3.
35. McAuley JB, Michelson MK, Schantz PM. Trichinellosis surveillance, United States, 1987–1990. *MMWR CDC Surveill Summ.* 1991;40:35–42.
36. Shiota T, Arizono N, Yoshioka T, Ishikawa Y, Fujitake J, Fujii H, et al. Imported trichinellosis with severe myositis—report of a case [in Japanese]. *Kansenshogaku Zasshi.* 1999;73:76–82.
37. Ancelle T, De Bruyne A, Niang M, Poisson D-M, Prazuck T, Fur A, et al. Outbreak of trichinellosis caused by *Trichinella nativa* due to consumption of bear meat [in French]. *Bulletin Épidémiologique Hebdomadaire.* 2006;14:96–8.
38. Houzé S, Ancelle T, Matra R, Boceno C, Carlier Y, Gajadhar AA, et al. Trichinellosis acquired in Nunavut, Canada in September 2009: meat from grizzly bear suspected. *Euro Surveill.* 2009;14:pii=19383.
39. Burke R, Masuoka P, Murrell KD. Swine *Trichinella* infection and geographic information system tools. *Emerg Infect Dis.* 2008;14:1109–11. doi:10.3201/eid14.07.071538
40. Sáaez-Royuela C, Telleria JL. The increased population of the wild boar (*Sus scrofa* L.) in Europe. *Mammal Rev.* 1986;16:97–101. doi:10.1111/j.1365-2907.1986.tb00027.x

Address for correspondence: K. Darwin Murrell, 5126 Russett Rd, Rockville, MD 20853, USA; email: [kdmurrell@comcast.net](mailto:kdmurrell@comcast.net)



# Worldwide Occurrence and Impact of Human Trichinellosis, 1986–2009

## Technical Appendix

### Section A. Total cases and incidence of human *Trichinella* infections by country and World Health Organization region, 1986–2009\*

Region/Country	Period	No. cases	Average incidence (range/dates)†	References
<b>African Region</b>				
		28		
Ethiopia	1986	8	0.02	Kefenie et al., 1988
	1990	20	0.04	Kefenie and Bero, 1992
<b>Region of the Americas</b>				
		7,179		
Canada	1987–2009	257	21.3 for Kewatin region (1989–1995)	Heinzig, 1996 Schellenberg et al., 2003 Ancelle et al., 2006 Gaulin et al., 2006 Houzé et al., 2009 Serhir et al., 2001 Møller et al., 2005 McIntyre et al., 2007
USA	1987–2007	652	0.016	McAuley et al., 1991
			(0.01–0.04/1987–1990)	Moorhead et al., 1999
			0.014	Roy et al., 2003
			(0.001–0.03/1991–1996)	Kennedy et al., 2009
			0.005	Smith et al., 2004
			(0.004–0.005/1997–2001)	Nelson et al., 2003
0.004	Madariaga et al., 2007			
		(0.001–0.002/2002–2007)	Centers for Disease Control and Prevention, 1987	
			Landry et al., 1992	
			Petri et al., 1988	
			Graves et al., 1996	
			McAuley et al., 1992	
			Dworkin et al., 1996	
Mexico	1986–2001	351	0.25 for State of Mexico (1986–1997)	Ortega-Pierres et al., 2000
Argentina	1990–2005	5,221	1.48 (1990–1999)	Chapa-Ruiz et al., 2001 Bolpe and Boffi, 2001 Nunez et al., 2008
Chile	1991–2004	698	0.36 (1996–2000) 0.2 (2000)	Rossi et al., 2007 Schenone et al., 2002 Morales et al., 1999 García et al., 2005
<b>Eastern Mediterranean Region</b>				
		50		
Iran	2007	6	0.008	Kia et al., 2008
Lebanon	1995	44	1.25	Haim et al., 1997
<b>European Region</b>				
		56,912		
Belarus	1986–1989	16	0.08 0.55	Ivanov et al., 1990 Skripova and Kovchur, 1994
Bosnia-Herzegovina	1993–2003	1,600	4.1 (0.1–8.0)	Cuperlovic et al., 2005
Bulgaria	1990–2006	4,108	2.9 (0.3–7.4/1990–1999) 2.4 (0.7–4.0/2000–2006)	Kurdova-Mintcheva et al., 2009 Petkova et al., 2008

Region/Country	Period	No. cases	Average incidence (range/dates)†	References
Croatia	1994–1999	1,350	4.8 (2.1–12.3)	Aleraj, 2008
	2000–2009	760	1.7 (0.02–3.9)	Croatia National Institute of Public Health, 2010 Marinculic et al., 2001 Vojnikovic et al., 2001 Puljiz et al., 2005 Cvitovic et al., 2007 Venus et al., 2008
Czech Republic	1986–2009	31	0.01	National Reference Laboratory for Tissue Helminthosis, 2010 Epshtein, 2010
Estonia	1986–2009	91	0.2 (0.0–2.9)	Dupouy-Camet et al., 2010
France	1986–1989	11	0.005 (0.0–0.02)	Durant et al., 1991
	1990–1999	1,158	0.19 (0.0–0.95)	Roumier et al., 1992 Soulé, 1992
	2000–2009	34	0.006 (0.0–0.02)	Ancelle et al., 1994 Gari-Toussaint et al., 1994 Soulé, 1994 Basset et al., 1995 Bernard et al., 1995 Laurichesse et al., 1997 Maillot et al., 1997 Ancelle et al., 1998 Dupouy-Camet et al., 1998 Haeghebaert et al., 1998 Hemery and Haeghebaert, 1999 Ranque et al., 2000 Gari-Toussaint et al., 2004 Gari-Toussaint et al., 2005 Ancelle et al., 2006 De Bruyne et al., 2006 Gaillard et al., 2007 Ivanov et al., 1990 Anonymous, 2010 Robert Koch Institute, 2010 Nothdurft et al., 1995 Littmann et al., 2006 Nockler et al., 2007 Jansen et al., 2008 Boutsini, 2009 Glatz et al., 2010
Georgia	1988	3	0.05	
Germany	1986–1989	19	0.005 (0.003–0.007)	
	1990–1999	105	0.009 (0.0–0.01)	
	2000–2009	61	0.005 (0.0–0.01)	
Greece	2009	1	0.008	
Hungary	1986–1989	74	0.18 (0.08–0.32)	
	1990–1999	27	0.027	
	2000–2009	57	0.057 (0.0–0.13)	
Ireland	2007	2	0.04	McHugh et al., 2007
Israel	2002	30	0.5	Hefer et al., 2004
	2004	200	3.0	Marva et al., 2005
Italy	1986–1989	448	0.19 (0.0–0.7)	E. Pozio, unpub. data
	1990–1999	649	0.01 (0.0–0.9)	Pozio et al., 1986 Pozio et al., 1987 Di Bari et al., 1990 Frongillo et al., 1992 Pozio et al., 1993 Pagni et al., 1994 Pozio et al., 1998 Pozio et al., 2001 Pozio and Marucci, 2003 Pozio et al., 2006 Pozio et al., 2009
	2000–2009	84	0.01 (0.0–0.07)	Witschi, 2010
Kyrgyzstan	1996	10	0.2	
Latvia	1986–1989	22	0.20 (0.07–0.34)	NRL of Latvia Dr. G. Deksnis
	1990–1999	345	1.37 (0.60–3.40)	
	2000–2009	269	1.13 (0.20–3.80)	
Lithuania	1990–2004	3,705	6.6 (0.6–21.8)	Bartuliene et al., 2005
	1989	75	2.0	Rockiene, 1996
	2005–2009	199	1.2 (0.4–3.4)	EpiNorth, 2010 Malakauskas, 2002 Bartuliene et al., 2009 Hristovski et al., 1992
Macedonia	1992	6	0.3	

Region/Country	Period	No. cases	Average incidence (range/dates)†	References
Poland	1986–1991	955	0.4 (0.05–0.8)	Adonajlo, 1988–1993
	1992	219	0.6	Koncki, 1994
	1993–1999	1,085	0.4 (0.05–1.5)	Seroka, 1995–2001
	2000–2001	88	0.1	Przybylska, 2002–2003
	2002–2007	737	0.3 (0.1–0.7)	Sadkowska-Todys and Golab, 2005–2009
Romania	1986–1989	2,621	2.9 (1.7–3.9)	Neghina et al., 2010a
	1990–1999	20,059	8.57 (4.1–16.1)	Blaga et al., 2007
	2000–2007	5,884	3.3 (1.8–5.9)	Neghina et al., 2009a Neghina et al., 2009b Nemet et al., 2009 Neghina et al., 2010b
Russia	1996–2002	971	0.48 (0.3–0.6)	Ozeretskovskaya et al., 2005
Serbia	1994–2003	5,210	5.05 (1.8–7.8)	Cuperlovic et al., 2005 Cuperlovic et al., 2001 Djordjevic et al., 2003
Slovakia	1986–1992	83	0.22 (0.0–0.64)	Dubinský et al., 1993
	1998	336	6.2	Dubinský et al., 2001
	2001	11	0.2	Reiterova et al., 2007
	2008	10	0.2	Paraličová et al., 2009
Slovenia	1989	200	10.5	Brglez, 1996
	1995	2	0.1	Saletinger et al., 2007
	2006	1	0.05	
Spain	1986–1989	233	0.15 (0.0–0.3)	Anonymous, 2010
	1990–1999	607	0.13 (0.02–0.40)	Anonymous, 1986
	2000–2009	404	0.08 (0.0–0.25)	Anonymous, 1987 Benitez et al., 1987 Anonymous, 1988 de la Torre et al., 1989 Serrano et al., 1989 Anonymous, 1990 Cobo et al., 1991 Mangas Gallardo and Tello Anchueta, 1994 de la Cruz de Julián et al., 1994 Anonymous, 1995 Tiberio et al., 1995 Tiberio et al., 1997 Rodríguez-Osorio et al., 1999 Martínez Corral et al., 2000 Lopez Hernandez et al., 2001 Cortés Blanco et al., 2002 Gomez Garcia et al., 2003 Herraez Garcia et al., 2003 Gallardo et al., 2007 Arevalo et al., 2009 Lupinc et al., 2003 J.C. Lozano-Becerra, pers. comm. Heper et al., 2005 Akkoc et al., 2009 Milne et al., 2001
Switzerland	1994	3	0.04	Ukraine Ministry of Health, 2010
Turkey	2003	7	0.01	
	2004	418	0.59	
United Kingdom	1999	7	0.01	
Ukraine	1986–1989	147	0.06 (0.04–0.10)	
	1990–1999	955	0.18 (0.09–0.3)	
	2000–2009	108	0.018 (0.0–0.08)	
South-East Asia Region		219		
Thailand	1993–2007	216	0.35	Limsuwan and Sirprasert 1994 Charkrit, 1998 Jongwutiwes et al., 1998 Watt et al., 2000 Chotmongkul et al., 2005 Khumjui et al., 2008
India	1996–2002	3	0.0003	Alipuria et al., 1996 Handa et al., 2000 Mohan et al., 2002
Western Pacific Region		1,344		



Region/Country	Period	No. cases	Average incidence (range/dates)†	References
People's Republic of China	1995–2009	1,137	Yunan, 1.0 Henan, 0.001 Tibet, 1.74 Sichuan, 0.028 Hubei, 0.03 (2000–2003)	Cui et al., 1997 Lo et al., 2009 Wang and Cui, 2001 Xu et al., 1995 Gong et al., 2008 Ye et al., 2003 Ci et al., 2003
Japan	1999–2005	4	NA	Kusuhara et al., 1999 Shiota et al., 1999 Nakamura et al., 2003 Yoshikawa et al., 2005
South Korea	1999–2003	8	0.016	Sohn et al., 2003 Kim et al., 2003 Lee et al., 1999
Laos	2004–2006	123	2.09	Sayasone et al., 2006 Taybouavone et al., 2009 Suwansrinen et al., 2007 Barennes et al., 2008
Singapore	1998	25	0.64	Kurup et al., 2000
Vietnam	2001–2004	47	0.058	Taylor et al., 2009 De et al., 2006

\*Human trichinellosis infections acquired in countries different from those where the disease was developed and diagnosed were not included in this table. NA, insufficient data for incidence calculation.

†Incidence/10<sup>5</sup> person-years. The incidence period is reported because the data may have been obtained over a period briefer than the interval on which total cases are based. For some countries, incidence was not reported and was calculated from data available in the report referenced.

## References

### African Region

#### Ethiopia

Kefenie H, Bero G. Trichinosis from wild boar meat in Gojjam, north-west Ethiopia. *Trop Geogr Med.* 1992;44:278–80. [PubMed](#)

Kefenie H, Wolde H, Abuherpo M. Trichinosis from wild boar meat in Arsi, central Ethiopia. *Ethiop Med J.* 1988;26:97–100. [PubMed](#)

### Region of the Americas

#### Canada

Ancelle T, Bruyne A, Nuang M, Poisson DM, Prazuck T, Fur A, et al. Epidemic of *Trichinella nativa* trichinosis caused by consumption of bear meat, France, 2005. *Bulletin Épidémiologique Hebdomadaire.* 2006;14:96–8.

Public Health Agency of Canada. Outbreak of trichinellosis in French hunters who ate Canadian black bear meat. *Can Commun Dis Rep.* 2006;32:109–12. [PubMed](#)

Heinzig L. Case reviews—botulism and trichinosis. *EpiNorth.* 1996;8(4):7.

Houzé S, Ancelle T, Matra R, Boceno C, Carlier Y, Gajadhar AA, et al. Trichinellosis acquired in Nunavut, Canada in September 2009: meat from grizzly bear suspected. *Euro Surveill.* 2009;14:5. [PubMed](#)

McIntyre L, Pollack SL, Fyfe M, Gajadhar A, Issac-Renton J, Fung J, et al. Trichinosis from consumption of wild game meat. *Can Med Assoc J.* 2007;176:449–51. [doi:10.1503/cmaj.061530](https://doi.org/10.1503/cmaj.061530)

Møller LN, Petersen E, Kapel CMO, Melbye M, Koch A. Outbreak of trichinellosis associated with consumption of game meat in West Greenland. *Vet Parasitol.* 2005;132:131–6. [PubMed](#)  
[doi:10.1016/j.vetpar.2005.05.041](https://doi.org/10.1016/j.vetpar.2005.05.041)

Schellenberg RS, Tan B, Irvine JD, Stockdale DR, Gajadhar AA, Serhir B, et al. An outbreak of trichinellosis due to consumption of bear meat infected with *Trichinella nativa* in 2 northern Saskatchewan communities. *J Infect Dis.* 2003;188:835–43. [PubMed](#) [doi:10.1086/378094](https://doi.org/10.1086/378094)

Serhir B, MacLean JD, Healey S, Forbes L. Outbreak of trichinellosis associated with arctic walrus in northern Canada, 1999. *Can Commun Dis Rep.* 2001;27–04:31–36.

#### **United States**

Centers for Disease Control and Prevention. Trichinosis—Hawaii. *MMWR Morb Mortal Wkly Rep.* 1987;36:14–6. [PubMed](#)

Dworkin MS, Gamble HR, Zarlenga DS, Tennican PO. Outbreak of trichinellosis associated with eating cougar jerky. *J Infect Dis.* 1996;174:663–6. [PubMed](#) [doi:10.1093/infdis/174.3.663](https://doi.org/10.1093/infdis/174.3.663)

Graves T, Harkness J, Crutcher JM. Case report: locally acquired trichinosis in an immigrant from Southeast Asia. *J Okla State Med Assoc.* 1996;89:402–4. [PubMed](#)

Kennedy ED, Hall RL, Montgomery SP, Pyburn DG, Jones JJ. Trichinellosis surveillance—United States, 2002–2007. *MMWR Surveill Summ.* 2009;58(9):1–7. [PubMed](#)

Landry SM, Kiser D, Overby T, Mays B, Caplen CW. Trichinosis: common source outbreak related to commercial pork. *South Med J.* 1992;85:428–9. [PubMed](#) [doi:10.1097/00007611-199204000-00022](https://doi.org/10.1097/00007611-199204000-00022)

Madariaga MG, Cachay ER, Zarlenga DS. Case report: a probable case of human neurotrichinellosis in the United States. *Am J Trop Med Hyg.* 2007;77:347–9. [PubMed](#)

McAuley JB, Michelson MK, Schantz PM. Trichinellosis surveillance, United States, 1987–90. *MMWR Surveill Summ.* 1991;40(SS-3):35–42. [PubMed](#)

- McAuley JB, Michelson MK, Hightower AW, Engeran S, Wintermeyer LA, Schantz PM. A trichinosis outbreak among Southeast Asian refugees. *Am J Epidemiol.* 1992;135:1404–10. [PubMed](#)
- Moorhead A, Grunenwald PE, Dietz VJ, Schantz PM. Trichinellosis in the United States, 1991–1996: declining but not gone. *Am J Trop Med Hyg.* 1999;60:66–9. [PubMed](#)
- Nelson M, Wright TL, Pierce A, Krogwold RA. A common source outbreak of trichinosis from consumption of bear meat. *J Environ Health.* 2003;65:16–19, 24. [PubMed](#)
- Petri WA, Holsinger JR, Pearson RD. Common-source outbreak of trichinosis associated with eating raw home-butchered pork. *South Med J.* 1988;81:1056–8. [PubMed](#) [doi:10.1097/00007611-198808000-00029](https://doi.org/10.1097/00007611-198808000-00029)
- Roy SL, Lopez AS, Schantz PM. Trichinellosis surveillance—United States, 1997–2001. *MMWR Surveill Summ.* 2003;52(6):1–8. [PubMed](#)
- Centers for Disease Control and Prevention. Trichinellosis associated with bear meat—New York and Tennessee, 2003. *MMWR Morb Mortal Wkly Rep.* 2004;53:606–10. [PubMed](#)
- Mexico**
- Chapa-Ruiz MR, González-Pantaleón D, Morales-Galán A, Contreras-Ramos A, Salinas-Tobón MR, Martínez Y, et al. A follow-up study of the human class and subclass antibody response developed against the adult stage of *Trichinella spiralis*. *Parasite.* 2001;8(Suppl):S163–7. [PubMed](#)
- Ortega-Pierres MG, Arriga C, Yopez-Mulia L. Epidemiology of trichinellosis in Mexico, Central and South America. *Vet Parasitol.* 2000;93:201–25. [PubMed](#) [doi:10.1016/S0304-4017\(00\)00342-3](https://doi.org/10.1016/S0304-4017(00)00342-3)
- Argentina**
- Bolpe J, Boffi R. Human trichinellosis in Argentina. Review of the casuistry registered from 1990 to 1999. *Parasite.* 2001;8(2 Suppl):S78–80. [PubMed](#)
- Costantino SN, Gentile T, Venturiello SM. Immunoparasitological evaluation of *Trichinella spiralis* infection during human pregnancy: a small case series. *Trans R Soc Trop Hyg.* 2008;102:662–8. [PubMed](#) [doi:10.1016/j.trstmh.2008.03.009](https://doi.org/10.1016/j.trstmh.2008.03.009)
- Rossi L, Coca F, Cricelli C, Troncoso A. First case of trichinosis caused by consumption of undercooked horse meat. *J Infect Develop Countries.* 2007;1:217–9.

## Chile

García E, Mora L, Torres P, Jercic MI, Mercado R. First record of human trichinosis in Chile associated with consumption of wild boar (*Sus scrofa*). Mem Inst Oswaldo Cruz. 2005;100:17–8. [PubMed](#)  
[doi:10.1590/S0074-02762005000100003](https://doi.org/10.1590/S0074-02762005000100003)

Morales MA, Vasquez J, Luengo J. Distribution and prevalence of trichinosis in man and pigs in Chile (1989–1995). J Parasitol Dia. 1999;23:63–5.

Schenone H, Olea A, Schenone H, Contreras MC, Mercado R, Sandoval L, et al. Current epidemiological situation of trichinosis in Chile. 1991–2000. Rev Med Chil. 2002;130:281–5. [PubMed](#)

## Eastern Mediterranean Region

### Iran

Kia EB, Meamar AR, Zahabiun F, Soodbakhsh A, Kordbacheh P. An outbreak of human trichinellosis due to the consumption of boar meat infected with *Trichinella* sp. [in Iranian]. Iran J Infect Dis Trop Med. 2008;41:35–8.

### Lebanon

Haim M, Efrat M, Wilson M, Schantz PM, Cohen D, Shemer J. An outbreak of *Trichinella spiralis* infection in southern Lebanon. Epidemiol Infect. 1997;119:357–62. [PubMed](#)  
[doi:10.1017/S0950268897007875](https://doi.org/10.1017/S0950268897007875)

## European Region

### Belarus

Ivanov KS, Antonov VS, Knysh GG, Khadzhaeva AN, Antykova LP, Lavrova VP, et al. The clinical characteristics of 2 outbreaks of trichinellosis [in Russian]. Med Parazitol (Mosk). 1990;4:41–2. [PubMed](#)

Skripova LV, Kovchur VN. Trichinellosis in Byelorussia. Wiad Parazytol. 1994;40:389–91. [PubMed](#)

### Bosnia-Herzegovina

Cuperlovic K, Djordjevic M, Pavlovic S. Re-emergence of trichinellosis in southern Europe due to political and economic changes. Vet Parasitol. 2005;132:159–66. [PubMed](#)  
[doi:10.1016/j.vetpar.2005.05.047](https://doi.org/10.1016/j.vetpar.2005.05.047)



## **Bulgaria**

Kurdova-Mintcheva R, Jordanova D, Ivanova M. Human trichinellosis in Bulgaria—epidemiological situation and trends. *Vet Parasitol.* 2009;159:316–9. [PubMed](#)

Petkova S, Mihov L, Vutova K, Tsenov I, La Rosa G, Pozio E. Epidemiological and clinical patterns of trichinellosis in Bulgaria from 1995 to 2002. *Parasite.* 2008;15:86–8. [PubMed](#)

## **Croatia**

Aleraj B. Actual epidemiology of trichinellosis in Croatia [in Croatian]. *Croatian Journal of Public Health* 2008;4(14). [http://www.hczj.hr/sadrzaj.php?broj\\_casopisa=14](http://www.hczj.hr/sadrzaj.php?broj_casopisa=14)

Croatian National Institute of Public Health. 2010. <http://www.hzjz.hr/epocetna>

Cvitović A, Miletić-Medved M, Gjenaro-Margan I. An epidemic of trichinellosis in autumn 2004 in Slavonski Brod [in Croatian]. *Acta Med Croatica.* 2007;61:215–8. [PubMed](#)

Marinculić A, Gaspar A, Duraković E, Pozio E, La Rosa G. Epidemiology of swine trichinellosis in the Republic of Croatia. *Parasite.* 2001;8(2 Suppl):S92–4. [PubMed](#)

Puljiz I, Beus A, Kuzman I, Seiwerth S. Electrocardiographic changes and myocarditis in trichinellosis: a retrospective study of 154 patients. *Ann Trop Med Parasitol.* 2005;99:403–11. [PubMed](#)  
[doi:10.1179/136485905X36307](https://doi.org/10.1179/136485905X36307)

Venus M, Puntarić D, Grgić M, Gmajnić R, Miskulin M. The effect of pest control on the incidence of trichinosis in Virovitica-Podravina County, Croatia. *Vet Parasitol.* 2008;156:226–33. [PubMed](#)  
[doi:10.1016/j.vetpar.2008.06.010](https://doi.org/10.1016/j.vetpar.2008.06.010)

Vojniković B, Brncić N, Zamolo G, Budiselić R, Njirić S, Novak S. Severe trichinellosis cured with pulse doses of glucocorticoids. *Coll Antropol.* 2001;25 Suppl:131–5. [PubMed](#)

## **Czech Republic**

National Reference Laboratory for Tissue Helminthosis of the Czech Republic 1986–2009 [email: [Libuse.kolarova@lfl.cuni.cz](mailto:Libuse.kolarova@lfl.cuni.cz)]

## **Estonia**

Epshtein J. Estonian trichinellosis morbidity data 1985–2009. *Dep. Communicable Disease Surveillance and Control.* [email: [jevgenia.epstein@terviseamet.ee](mailto:jevgenia.epstein@terviseamet.ee)]

## France

- Ancelle T, Dupouy-Camet J, Desenclos JC, Maillot E, Charlet F, Gravelat-Desclaux C, et al. Epidémie de trichinellose (France, 1993) bilan des investigations. Bulletin Épidémiologique Hebdomadaire. 1994;29:127–9.
- Ancelle T, Dupouy-Camet J, Desenclos JC, Maillot E, Savage-Houze S, Drucker FCJ, et al. A multifocal outbreak of trichinellosis linked to horse meat imported from North America to France in 1993. Am J Trop Med Hyg. 1998;59:615–9. [PubMed](#)
- Ancelle T, De Bruyne A, Niang M, Poisson DM, Prazuck T, Fur A, et al. Epidémie de trichinellose à *Trichinella nativa* due à la consommation de viande d'ours, France 2005. Bulletin Épidémiologique Hebdomadaire. 2006;14:96–8.
- Basset D, Thiebaut MM, Pratlong F, Abraham B, Moryoussef A, Baldet P, et al. Epidémies familiales de trichinose dues a l'ingestion de sanglier sauvage dans la region Languedoc. Bulletin Épidémiologique Hebdomadaire. 1995;44:195.
- Bernard E, Ozouf N, Toussaint-Gari M, Marty P, Pozio E, Le Fichoux Y, et al. Two new outbreaks of trichinosis [in French]. Med Mal Infect. 1995;25:611–4. [doi:10.1016/S0399-077X\(05\)81038-8](https://doi.org/10.1016/S0399-077X(05)81038-8)
- De Bruyne A, Ancelle T, Vallée I, Boireau P, Dupouy-Camet J. Human trichinellosis from wild boar meat: a continuing parasitic risk in France. Euro Surveill. 2006;11: E060914.5. [PubMed](#)
- Dupouy-Camet J, Allegretti S, Truong TP. Enquete sur l'incidence de la trichinellose en France (1994–1995). Bulletin Épidémiologique Hebdomadaire. 1998;28:122–3.
- Dupouy-Camet J, Ancelle T, Talabani H. Surveillance de la trichinellose humaine en France. Rapport du CNR des Trichinella, 1 janvier 2009 - 31 decembre 2009. Rapport du CNR des Trichinella 2010.
- Durant J, Toussaint-Gari M, Bernard E, Marty P, Le Fichoux Y, Dellamonica P. Epidémie familiale de trichinose. Sem Hop Paris. 1991;67:1507–12.
- Gaillard T, Martinaud C, Kérébel S, Cellarier G, Muzellec Y, Brisou P. A propos de deux cas de trichinellose à *Trichinella britovi*. Ann Biol Clin (Paris). 2007;65:308–12. [PubMed](#)
- Gari-Toussaint M, Bernard E, Quaranta JF, Marty P, Soler C, Ozouf N, et al. First report in France of an outbreak of human trichinellosis due to *Trichinella britovi*. In: Campbell WC, Pozio E, Bruschi F, editors. Trichinellosis. Proceedings of the 8th International Conference on Trichinellosis, 1994. Rome: Istituto Superiore di Sanita Press; 1994. p. 465–8.

Gari-Toussaint M, Tieulié N, Baldin JL, Marty P, Dupouy-Camet J, Delaunay P, et al. Trichinellose à *Trichinella britovi* dans les Alpes-Maritimes après consommation de viande de sanglier congelée, automne 2003. Bulletin Épidémiologique Hebdomadaire. 2004;21:87–8.

Gari-Toussaint M, Tieulié N, Baldin JL, Dupouy-Camet J, Delaunay P, Fuzibet JG, et al. Human trichinellosis due to *Trichinella britovi* in southern France after consumption of frozen wild boar meat. Euro Surveill. 2005;10:117–8. [PubMed](#)

Haeghebaert S, Servat M, Duchon C, Minet JC, Agrech AE, Thiese I, et al. Outbreak of trichinellosis in the Midi-Pyrénées region of France January–March 1998. Euro Surveill. 1998;3:83–5. [PubMed](#)

Hemery C, Haeghebaert S. New outbreak of trichinellosis in the Midi-Pyrénées region of France, September–October 1998. Euro Surveill. 1999;4:13–4. [PubMed](#)

Laurichesse H, Cambon M, Perre D, Ancelle T, Mora M, Hubert B, et al. Outbreak of trichinosis in France associated with eating horse meat. Commun Dis Rep CDR Rev. 1997;7:R69–73.

Maillot E, Desenclos JC, Dupouy-Camet J, Aubert P. Bulletin Épidémiologique Hebdomadaire. 1997;49:217–8.

Ranque S, Faugère B, Pozio E, La Rosa G, Tamburrini A, Pelissier JF, et al. *Trichinella pseudospiralis* outbreak in France. Emerg Infect Dis. 2000;6:543–7. [PubMed](#) doi:10.3201/eid0605.000517

Roumier M, Milhe P, Hautefort B, Benoist B. Four cases of trichinosis in Camargue (France) by consumption of meat of boar [in French]. Med Mal Infect. 1992;22:947–8. doi:10.1016/S0399-077X(05)80639-0

Soulé C. Trichinellose in France (1988–1990). Wiad Parazytol. 1992;38:153. [PubMed](#)

Soulé C. Trichinellose in France (1991–1992). Wiad Parazytol. 1994;40:394. [PubMed](#)

### **Georgia**

Ivanov KS, Antonov VS, Knysh GG, Khadzhaeva AN, Antykova LP, Lavrova VP, et al. The clinical characteristics of 2 outbreaks of trichinelliasis [in Russian]. Med Parazitol (Mosk). 1990;Jul–Aug:41–2. [PubMed](#)

### **Germany**

Anonymous. Human cases of trichinellosis in Germany (1965–2008). <http://beck-online.beck.de/default.aspx?bcid=Y-100-G-BSeuchG>

Jansen A, Schöneberg I, Stark K, Nöckler K. Epidemiology of trichinellosis in Germany, 1996–2006. Vector Borne Zoonotic Dis. 2008;8:189–96. [PubMed](#) doi:10.1089/vbz.2007.0183

Littmann M, Nöckler K, Hallauer J. Cluster of trichinellosis cases in Mecklenburg-Vorpommern, Germany. *Euro Surveill.* 2006;11: E060518.1.

Nöckler K, Wichmann-Schauer H, Hiller P, Müller A, Bogner K. Trichinellosis outbreak in Bavaria caused by cured sausage from Romania, January 2007. *Euro Surveill.* 2007;12: E070823.2.

Nothdurft HD, Brommer M, Eichenlaub D, Löscher T. A small outbreak of trichinosis in Germany caused by imported smoked ham [in German]. *Dtsch Med Wochenschr.* 1995;120:173–6. [PubMed doi:10.1055/s-2008-1055330](#)

Robert Koch Institute. InstitutSurStat@RKI, 2010. Human cases of trichinellosis in Germany from 2001 to 12 May 2010.

### **Greece**

Boutsini S. Report on a trichinellosis outbreak in 2009. Pers. comm. [email: [sboutsini@yahoo.gr](mailto:sboutsini@yahoo.gr)].

### **Hungary**

Glatz K, Danka J, Kucsera I, Pozio E. Human trichinellosis in Hungary from 1995 to 2009. *Parasite.* 2010;17:193–8. [PubMed](#)

### **Ireland**

McHugh G, Kiely D, Low J, Healy ML, Hayes C, Clarke S. Importation of Polish trichinellosis cases to Ireland, June 2007. *Euro Surveill.* 2007;12: E070719.3. [PubMed](#)

### **Israel**

Hefer E, Rishpon S, Volovik I. Trichinosis outbreak among Thai immigrant workers in the Hadera sub-district [in Hebrew]. *Harefuah.* 2004;143:656–60. [PubMed](#)

Marva E, Markovics A, Gdalevich M, Asor N, Sadik C, Leventhal A. Trichinellosis outbreak. *Emerg Infect Dis.* 2005;11:1979–81. [PubMed](#)

### **Italy**

Di Bari C, Santagada G, Pozio E, Schiraldi O. Epidemiological research on trichinellosis in Apulia and Basilicata (southern Italy). *Eur J Epidemiol.* 1990;6:412–5. [PubMed doi:10.1007/BF00151717](#)

Frongillo RF, Baldelli B, Pozio E, Crapa G, Di Giuli C, Santirocchi M, et al. Report on an outbreak of trichinellosis in central Italy. *Eur J Epidemiol.* 1992;8:283–8. [PubMed doi:10.1007/BF00144815](#)

Pagni P, Bindi M, Burzagli L, Melchior M. An outbreak of trichinellosis in central Italy caused by pork meat ingestion. In: Campbell WC, Pozio E, Bruschi F, editors. *Trichinellosis. Proceedings of the*



8th International Conference on Trichinellosis, 1994. Rome: Istituto Superiore di Sanita Press; 1994. p. 487–90.

Pozio E, Santagada G, Di Bari C. Outbreak of trichinellosis in southern Italy. *Trans R Soc Trop Med Hyg.* 1986;80:997–8. [PubMed doi:10.1016/0035-9203\(86\)90291-9](#)

Pozio E, Cappelli O, Marchesi L, Valeri P, Rossi P. Third outbreak of trichinellosis caused by consumption of horse meat in Italy. *Ann Parasitol Hum Comp.* 1987;63:48–53.

Pozio E, Varese P, Gómez Morales MA, Croppo GP, Pelliccia D, Bruschi F. Comparison of human trichinellosis caused by *Trichinella spiralis* and *Trichinella britovi*. *Am J Trop Med Hyg.* 1993;48:568–75. [PubMed](#)

Pozio E, Sacchini D, Boni P, Tamburrini A, Alberici F, Paterlini F. Human outbreak of trichinellosis associated with the consumption of horsemeat in Italy. *Euro Surveill.* 1998;3:85–6.

Pozio E, La Rosa G, Gomez Morales MA. Epidemiology of human and animal trichinellosis in Italy since its discovery in 1887. *Parasite.* 2001;8:S106–8. [PubMed](#)

Pozio E, Marucci G. *Trichinella*-infected pork products: a dangerous gift. *Trends Parasitol.* 2003;19:338. [PubMed doi:10.1016/S1471-4922\(03\)00138-7](#)

Pozio E, Mesina P, Sechi F, Pira M, Liciardi M, Cossu P, et al. Human out break of trichinellosis in the Mediterranean island of Sardinia, Italy. *Vet Parasitol.* 2006;140:177–80. [PubMed doi:10.1016/j.vetpar.2006.03.012](#)

Pozio E, Cossu P, Marucci G, Amati M, Ludovisi A, Gomez Morales MA, et al. The birth of a *Trichinella britovi* focus on the Mediteranean island of Sardinia (Italy). *Vet Parasitol.* 2009;159:361–3. [PubMed](#)

### **Kyrgyzstan**

Witschi M. WHO Country Office of Kyrgyzstan 2010 [email: [wim@euro.who.int](mailto:wim@euro.who.int)].

### **Latvia**

Public Health Agency Annual Report of Latvia [email: [gunita.deksne@bior.gov.lv](mailto:gunita.deksne@bior.gov.lv)].

### **Lithuania**

Bartuliene A, Jasulaitiene V, Malakauskas A. Human trichinellosis in Lithuania, 1990–2004. *Euro Surveill.* 2005;10: E050714.6. [PubMed](#)

Bartuliene A, Liausediene R, Motiejuniene V. Trichinellosis outbreak in Lithuania, Ukmerge region, June 2009. *Euro Surveill.* 2009;14:24. [PubMed](#)

EpiNorth.

[http://www.epinorth.org/eway/default.aspx?pid=230&trg=Area\\_5279&MainArea\\_5260=5279:0:15,2937:1:0:0:0:0&Area\\_5279=5291:44530:1:5290:1:0:0&diseaseid=38](http://www.epinorth.org/eway/default.aspx?pid=230&trg=Area_5279&MainArea_5260=5279:0:15,2937:1:0:0:0:0&Area_5279=5291:44530:1:5290:1:0:0&diseaseid=38)

Malakauskas A. molecular epidemiology of trichinellosis in Lithuania and biological characteristics of *Trichinella* genotypes in rat. PhD Thesis, Copenhagen, Denmark: The Royal Veterinary and Agricultural University; 2002. p. 1–135.

Rockiene A. Trichinellosis in Lithuania. 2nd Nordic-Baltic congress on Infectious Diseases “Prevention in Focus.” Latvia 1996, p. 38–39.

### **Macedonia**

Hristovski M, Dzimrevski J, Bosnakovski J, Markovski G, Ancevski V, Krstic I. Trichinellosis in the Republic of Macedonia. Makedonski Veterinaren Pregled. 1992;21:85–90.

### **Poland**

Adonajlo A. Trichinellosis in Poland in 1986 [in Polish]. Przegł Epidemiol. 1988;42:113–5. [PubMed](#)

Adonajlo A. Trichinellosis in Poland in 1987 [in Polish]. Przegł Epidemiol. 1989;43:107–9. [PubMed](#)

Adonajlo A. Trichinellosis in Poland in 1988 [in Polish]. Przegł Epidemiol. 1990;44:119–23. [PubMed](#)

Adonajlo A. Trichinellosis in Poland in 1989 [in Polish]. Przegł Epidemiol. 1991;45:113–5. [PubMed](#)

Adonajlo A. Trichinellosis in Poland in 1990 [in Polish]. Przegł Epidemiol. 1992;46:127–30. [PubMed](#)

Adonajlo A. Trichinellosis in Poland in 1991 [in Polish]. Przegł Epidemiol. 1993;47:161–4. [PubMed](#)

Koncki A. Trichinellosis in 1992 [in Polish]. Przegł Epidemiol. 1994;48:145–7. [PubMed](#)

Przybylska A. Trichinellosis in Poland in 2000 [in Polish]. Przegł Epidemiol. 2002;56:353–6. [PubMed](#)

Przybylska A. Trichinellosis in Poland in 2001 [in Polish]. Przegł Epidemiol. 2003;57:153–8. [PubMed](#)

Przybylska A. Trichinellosis in Poland in 2002 [in Polish]. Przegł Epidemiol. 2004;58:157–64. [PubMed](#)

Sadkowska-Todys M, Gołab E. Trichinellosis in Poland in 2003 [in Polish]. Przegł Epidemiol. 2005;59:327–30. [PubMed](#)

Sadkowska-Todys M, Gołab E. Trichinellosis in Poland in 2004 [in Polish]. Przegł Epidemiol. 2006;60:503–7. [PubMed](#)

Sadkowska-Todys M, Gołab E. Trichinellosis in Poland in 2005 [in Polish]. Przegł Epidemiol. 2007;61:301–4. [PubMed](#)

- Sadkowska-Todys M, Gołab E. Trichinellosis in Poland in 2008 [in Polish]. *Przegl Epidemiol.* 2008;62:347–50. [PubMed](#)
- Sadkowska-Todys M, Gołab E. Trichinellosis in Poland in 2007 [in Polish]. *Przegl Epidemiol.* 2009;63:263–6. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1993 [in Polish]. *Przegl Epidemiol.* 1995;49:181–7. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1994 [in Polish]. *Przegl Epidemiol.* 1996;50:191–7. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1995 [in Polish]. *Przegl Epidemiol.* 1997;51:157–60. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1996 [in Polish]. *Przegl Epidemiol.* 1998;52:133–7. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1997 [in Polish]. *Przegl Epidemiol.* 1999;53:155–8. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1998 [in Polish]. *Przegl Epidemiol.* 2000;54:175–9. [PubMed](#)
- Seroka D. Trichinellosis in Poland in 1999 [in Polish]. *Przegl Epidemiol.* 2001;55:155–8. [PubMed](#)
- Romania**
- Blaga R, Durand B, Antoniu S, Gherman C, Cretu C, Cozma V, et al. A dramatic increase in the incidence of human trichinellosis in Romania over the past 25 years: impact of political changes and regional food habits. *Am J Trop Med Hyg.* 2007;76:983–6. [PubMed](#)
- Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Epidemiological and diagnostic findings during a 16-year-long trichinellosis surveillance in Timis County, Romania. *Vet Parasitol.* 2009a;159:328–31. [PubMed](#)
- Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Trichinellosis, a threatening and re-emerging disease in a Romanian western county. *Vector Borne Zoonotic Dis.* 2009b;9:717–21. [PubMed](#) [doi:10.1089/vbz.2008.0204](#)
- Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Epidemiology and epizootology of trichinellosis in Romania 1868–2007. *Vector Borne Zoonotic Dis.* 2010a;10:323–8. [PubMed](#) [doi:10.1089/vbz.2009.0084](#)
- Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Evidence of *Trichinella spiralis* in Timis county, Romania: a report of a winter trichinellosis outbreak in 2008 due to consumption of contaminated pork. *Vector Borne Zoonotic Dis.* 2010b;10:931–3. [PubMed](#) [doi:10.1089/vbz.2009.0057](#)

Nemet C, Rogozea L, Dejica R. Results of the follow-up of the former trichinosis patients from Brasov county—Romania. *Vet Parasitol.* 2009;159:320–3. [PubMed](#)

### **Russia**

Ozeretskovskaya NN, Mikhailova LG, Sabgaida TP, Dovgalev AS. New trends and clinical patterns of human trichinellosis in Russia at the beginning of the XXI century. *Vet Parasitol.* 2005;132:167–71. [PubMed](#) [doi:10.1016/j.vetpar.2005.05.056](https://doi.org/10.1016/j.vetpar.2005.05.056)

### **Serbia**

Cuperlovic K, Djorjevic M, Pavlovic S, Sofronic-Milosavljevic L. Present status of trichinellosis in Yugoslavia: Serbia. *Parasite.* 2001;8(2 Suppl):S95–7. [PubMed](#)

Cuperlovic K, Djordjevic M, Pavlovic S. Re-emergence of trichinellosis in southern Europe due to political and economic changes. *Vet Parasitol.* 2005;132:159–66. [PubMed](#)  
[doi:10.1016/j.vetpar.2005.05.047](https://doi.org/10.1016/j.vetpar.2005.05.047)

Djordjevic M, Bacic M, Petricevic M, Cuperlovic K, Malakauskas A, Kapel CMO, et al. Social, political, and economic factors responsible for the re-emergence of trichinellosis in Serbia: a case study. *J Parasitol.* 2003;89:226–31. [PubMed](#) [doi:10.1645/0022-3395\(2003\)089\[0226:SPAEFR\]2.0.CO;2](https://doi.org/10.1645/0022-3395(2003)089[0226:SPAEFR]2.0.CO;2)

### **Slovakia**

Dubinský P, Stefancikova A, Hovorka I, Mitterpark J. Epidemiology of trichinellosis in Slovakia. In: Campbell WC, Pozio E, Bruschi F, editors. *Trichinellosis. Proceedings of the 8th International Conference on Trichinellosis, 7–10 September, 1993. Orvieto, Italy*, p. 515–520.

Dubinský P, Stefanciková A, Kinčeková J, Ondriska F, Reiterová K, Medvedová M. Trichinellosis in the Slovak Republic. *Parasite.* 2001;8(2 Suppl):S100–2. [PubMed](#)

Paraličová Z, Kinčeková J, Schréter I, Jarčuska P, Dubinský P, Porubčín S, et al. Outbreak of trichinellosis in eastern Slovakia. *Helminthologia.* 2009;46:209–13. [doi:10.2478/s11687-009-0039-2](https://doi.org/10.2478/s11687-009-0039-2)

Reiterová K, Kinčeková J, Snábel V, Marucci G, Pozio E, Dubinský P. *Trichinella spiralis*—outbreak in the Slovak Republic. *Infection.* 2007;35:89–93. [PubMed](#) [doi:10.1007/s15010-007-6122-z](https://doi.org/10.1007/s15010-007-6122-z)

### **Slovenia**

Brglez J. A new epidemic focus on trichinellosis [in Slovenian]. *Veterinarske Novice.* 1996;22:135–6.

Saletinger R, Rejc-Marko J, Unuk S, Logar J. Trichinosis—case presentation [in Slovenian]. *Zdrav Vestn.* 2007;76:175–8.

## Spain

Anonymous. Brotes de triquinosis. Espana. Temporada 1985/1986. Boletin Epidemiologico Semanal. 1986; N. 1748.

Anonymous. Brotes de triquinosis. Espana. Temporada 1986–1987. Boletin Epidemiologico Semanal. 1987; N. 1779.

Anonymous. Brotes de triquinosis. Espana. Temporadas 1987–1988. Boletin Epidemiologico Semanal. 1988; N. 1823.

Anonymous. Brotes de triquinosis. Espana. Temporadas 88/89 Y 89/90. Boletin Epidemiologico Semanal. 1990; N. 1857.

Anonymous. Vigilancia de la triquinosis en Espana. Boletin Epidemiologico Semanal. 1995;3:105–12.

Anonymous. Number of trichinellosis cases in Spain from 1986 to 2009. [tgarate@isciii.es](mailto:tgarate@isciii.es); 2010.

Arévalo A, Bringas MJ, Rodriguez R, Menor A. Description of a trichinosis outbreak in the province of Salamanca [in Spanish]. Rev Esp Quimioter. 2009;22:115–6. [PubMed](#)

Benitez JA, Cruz JM, Adrados G, Gonzalez T. Triquinosis: a proposito de 17 casos aparecidos en un mismo brote. Enferm Infecc Microbiol Clin. 1987;5:122–3.

Cobo J, Gómez Cerezo J, Medraño JC, Zapico R, Cruz Martinez A, Molina F, et al. Trichinosis. A study of a large outbreak on the Iberian peninsula [in Spanish]. An Med Interna. 1991;8:441–4.

Cortés Blanco M, Garcia-Cabanias A, Guerra-Peguero F, Ramos-Aceitero JM, Herrera-Guibert D, Martinez-Navarro JF. Outbreak of trichinellosis in Caceres, Spain, December 2001–February 2002. Euro Surveill. 2002;7:136–8. [PubMed](#)

de la Cruz de Julián I, Díaz García JM, Alvarez Lana P, García Colmenero C. An outbreak of trichinosis in Huerta del Marquesado (the Cañete-Cuenca basic health area). December 1992 to January 1993 [in Spanish]. Rev Sanid Hig Publica (Madr). 1994;68:513–20.

de la Torre Cecilia C, Espino Aguilar R, Cárdenas Talaverón C, Cañuelo Ruiz O, Garrido Palomo R, Baena Sáez J, et al. Trichinosis: presentation of 2 cases [in Spanish]. An Esp Pediatr. 1989;30:227–8. [PubMed](#)

Gallardo MT, Mateos L, Artieda J, Wesslen L, Ruiz C, Garcia MA, et al. Outbreak of trichinellosis in Spain and Sweden due to the consumption of wild boar meat contaminated with *Trichinella britovi*. Euro Surveill. 2007;12:E070315.1. [PubMed](#)

- Gomez-Garcia V, Hernandez-Quero J, Rodriguez-Orsorio M. Short report: human infection with *Trichinella britovi* in Granada, Spain. *Am J Trop Med Hyg.* 2003;68:463–4. [PubMed](#)
- Herráez García J, Leon Garcia LA, Lanusse Senderos C, Cortés Blanco M, García Cabanas A. Outbreak of trichinellosis in the region of la Vera (Carceres, Spain) caused by *Trichinella britovi* [in Spanish]. *An Med Interna.* 2003;20:63–6. [PubMed doi:10.4321/S0212-71992003000200003](#)
- López Hernández B, Velázquez de Castro MT, Galicia García MD, Sabonet JC. Outbreak of *Trichinella britovi* infection in Granada in the spring of 2000 [in Spanish]. *Rev Esp Salud Publica.* 2001;75:467–73. [PubMed doi:10.1590/S1135-57272001000500007](#)
- Mangas Gallardo I, Tello Anchuela O. Vigilancia de la triquinosis en Espana. Temporadas 1989/90–1992/93. *Bol Epidemiol Microbiol.* 1994;1:233–4.
- Martinez Corral JJ, Esteban Niveiro MJ, Oliet Pala R. Triquinosis producida por el consumo de carne de porcino infestado con *Trichinella spiralis*. Investigation y estudio de un brote alimentario. *Salud Publica.* 2000;12:36–40.
- Rodriguez-Orsorio M, Abad JM, De Haro T, Villa-Real R, Gomez Garcia V. Human trichinellosis in southern Spain: serologic and epidemiologic study. *Am J Trop Med Hyg.* 1999;61:834–7. [PubMed](#)
- Serrano R, Lacasa J, Velazquez J, Ziad F, Aznar R. Triquinosis: nuevo brote epidémico por ingesta de embutido de jabali. *Enferm Infecc Microbiol Clin.* 1989;7:428–31. [PubMed](#)
- Tiberio G, Lanzas G, Galarza MI, Sanchez J, Quilez I, Martinez Artola V. Short report: an outbreak of trichinosis in Navarra, Spain. *Am J Trop Med Hyg.* 1995;53:241–2. [PubMed](#)
- Tiberio G, Rivero M, Lanzas G, Redin D, Ardanaz E, Fernandez C, et al. Trichinellosis: study of 2 outbreaks in Navarre [in Spanish]. *Enferm Infecc Microbiol Clin.* 1997;15:151–3. [PubMed](#)

#### **Switzerland**

- Lozano-Becerra J.C., pers. comm. [email: [JuanCarlos.Lozano-Becerra@eoc.ch](mailto:JuanCarlos.Lozano-Becerra@eoc.ch)].
- Lupinc L, Markwalder K, Nigg C. Eosinophilie bei einer familie aus Bosnien. *Praxis.* 2003;92:2212–6. [PubMed doi:10.1024/0369-8394.92.51.2212](#)

#### **Turkey**

- Akkoc N, Kuruuzum Z, Aker S, Yuce A, Onen F, Yapar N, et al. A large-scale out break of trichinellosis caused by *Trichinella britovi* in Turkey. *Zoonoses Public Health.* 2009;56:65–70. [PubMed doi:10.1111/j.1863-2378.2008.01158.x](#)



Heper Y, Yilmaztepe F, Komitova R, Akalin H, Vutova K, Helvaci S. A trichinosis outbreak caused by wild boar meat in Turkey. *Parasite*. 2005;12:191–2. [PubMed](#)

### **United Kingdom**

Milne LM, Bhagani S, Bannister BA, Laitner SM, Moore P, Eza D, et al. Trichinellosis acquired in the United Kingdom. *Epidemiol Infect*. 2001;127:359–63. [PubMed](#)  
[doi:10.1017/S0950268801005994](https://doi.org/10.1017/S0950268801005994)

### **Ukraine**

Ministry of Healthcare of Ukraine. Trichinellosis in Ukraine from 1986 to 2010. Central health-epidemiology station of Ministry of healthcare of Ukraine. 2010.

## **South-East Asia Region**

### **Thailand**

Charkrit S. Study on clinical manifestations of trichinosis in Payao province. *Communicable Diseases Journal*. 1998;24:242–7.

Chotmongkol V, Intapan PM, Koonmee S, Kularbkaew C, Aungaree T. Case report: acquired progressive muscular hypertrophy and trichinosis. *Am J Trop Med Hyg*. 2005;72:649–50. [PubMed](#)

Limsuwan S, Siriprasert V. A clinical study on trichinosis in Changwat Phayao, Thailand. *Southeast Asian J Trop Med Public Health*. 1994;25:305–8. [PubMed](#)

Jongwutiwes S, Chantachum N, Kraivichian P, Siriyasatien P, Putaporntip C, Tamburrini A, et al. First outbreak of human trichinellosis caused by *Trichinella pseudospiralis*. *Clin Infect Dis*. 1998;26:111–5. [PubMed](#) [doi:10.1086/516278](https://doi.org/10.1086/516278)

Khumjui C, Choomkasien P, Dekumyoy P, Kusolsuk T, Kongkaew W, Chalamaat M, et al. Outbreak of trichinellosis caused by *Trichinella papuae*, Thailand. 2006. *Emerg Infect Dis*. 2008;14:1913–5. [PubMed](#)

Watt G, Saisorn S, Jongsakul K, Sakolvaree Y, Chaicumpa W. Blinded, placebo-controlled trial of antiparasitic drugs for trichinosis myositis. *J Infect Dis*. 2000;182:371–4. [PubMed](#)  
[doi:10.1086/315645](https://doi.org/10.1086/315645)

### **India**

Alipuria S, Sangha HK, Singh G, Pandhi S. Trichinosis—a case report. *Indian J Pathol Microbiol*. 1996;39:231–2. [PubMed](#)

Handa R, Aggarwal P, Sarkar C, Vijayarghavan M, Mattewal A, Arya V, et al. A patient with muscle weakness. *J Indian Rheum Assoc.* 2000;8:85–7.

Mohan H, Aggarwal R, Nada R, Punia RPS, Ahluwalia M. Trichinosis of psoas muscle. *J Assoc Physicians India.* 2002;50:729–30. [PubMed](#)

## Western Pacific Region

### China

Ci R, Si T, Wang HJ. An outbreak of trichinellosis in Milin County of Tibet Autonomous Region [in Chinese]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi.* 2003;21:60.

Cui J, Wang ZQ, Wu F, Jin XX. Epidemiological and clinical studies on an outbreak of trichinosis in central China. *Ann Trop Med Parasitol.* 1997;91:481–8. [PubMed doi:10.1080/00034989760842](#)

Wang ZQ, Cui J. The epidemiology of human trichinellosis in China during 1964–1999. *Parasite.* 2001;8:S63. [PubMed](#)

Gong XH, Guo WM, Cirendunzhu, Long EK, Ma Y, Bianbazuoma, et al. Investigation on an outbreak of trichinosis and clinical analysis [in Chinese]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi.* 2008;26(1):79–80.

Lo YC, Hung CC, Lai CS, Wu Z, Nagano I, Maeda T, et al. Human trichinosis after consumption of soft-shelled turtles, Taiwan. *Emerg Infect Dis.* 2009;15:2056–8. [PubMed](#)

Xu B, Chang J, Gao G, Tang B, Hong J, Chen Y, et al. Four outbreaks of human trichinellosis in Henan Province. *Chin Med J (Engl).* 1995;108:872–4. [PubMed](#)

Ye P, Jianhua H, Zhengkun Z, Desong G, Hao W. The investigation over an outbreak of trichinellosis caused by eating *Sus scrofa*. *Chinese Journal of Zoonoses.* 2003;19:12.

### Japan

Kusuhara Y, Maeno Y, Nagase K, Taniguchi K, Torikai K, Takahashi Y. A case of mixed infection with *Schistosoma haematobium* and *Trichinella sp.* *Kansenshogaku Zasshi.* 1999;73(6):614–7.

Nakamura T, Miura T, Nakaoka T, Nagano I, Takahashi Y, Iwamoto A. A case of trichinellosis with spontaneous remission [in Japanese]. *Kansenshogaku Zasshi.* 2003;77:839–43.

Shiota T, Arizono N, Yoshioka T, Ishikawa Y, Fujitake J, Fujii H, et al. Imported trichinellosis with severe myositis—report of a case [in Japanese]. *Kansenshogaku Zasshi.* 1999;73(1):76–82.

Yoshikawa M, Oujii Y, Moriya K, Saito K, Ishizaka S, Sukurai M, et al. Immunopositivity for *Trichinella spiralis* presenting in a case of paragonimiasis wetermani-suspected infection by ingestion of raw bear meat. J Nara Med Assoc. 2005;56:253–9.

#### **Korea**

Lee HC, Kim HJ, Oh HY, Kim JH, Kim HG, Lee MS, et al. A case of trichinosis caused by eating a wild badger. Korean Journal of Medicine 1999;56:134–8.

Sohn WM, Huh S, Chung DI, Pozio E. Molecular identification of Korean *Trichinella* isolates. Korean J Parasitol. 2003;41:125–7. [PubMed doi:10.3347/kjp.2003.41.2.125](#)

Kim E, Pyun RH, Park JH, Kim KH, Choi I, Park HH, et al. Family outbreak of trichinosis after eating a raw meat of wild swine. Korean J Parasitol. 2003;35:180–4.

#### **Laos**

Barennes H, Sayasone S, Odermatt P, Bruyney AD, Sitthivone HS, Newton PN, et al. A major trichinellosis outbreak suggesting a high endemicity of *Trichinella* infection in northern Laos. Am J Trop Med Hyg. 2008;78:40–4. [PubMed](#)

Sayasone S, Odermatt P, Vongphrachanh P, Keoluangkhot N, Dupouy-Camet J, Newton PN, et al. Trichinellosis outbreak in Borikhamxay Province, Lao PDR. Trans R Soc Trop Med Hyg. 2006;100:1126–9. [PubMed doi:10.1016/j.trstmh.2006.01.010](#)

Suwansrinon K, Wilde H, Burford B, Hanvesakul R, Visith Sitprijia V. Human trichinellosis from Laos. J Travel Med. 2007;14:274–7. [PubMed doi:10.1111/j.1708-8305.2007.00136.x](#)

Taybouavone T, Odermatt P, Keoluangkhot V, Delanos-Gregoire N, Dupouy-Camet J, Strobel M, et al. Trichinellosis during pregnancy: a case control study in the Lao People's Democratic Republic. Vet Parasitol. 2009;159:332–6. [PubMed](#)

#### **Singapore**

Kurup A, Yew WS, San LM, Ang B, Lim S, Tai GK. Outbreak of suspected trichinosis among travelers returning from a neighboring island. J Travel Med. 2000;7:189–93. [PubMed doi:10.2310/7060.2000.00058](#)

#### **Vietnam**

De NV, Toan ND, Son DT, Van P. Report an epidemic area of trichinellosis in Tuan Giao district, Lai Chau province. Proc 5th Seminar on Food- and Water-Borne Parasitic Zoonoses. Bangkok, Thailand. 2006Nov 28–30. p. 48.

Taylor WRJ, Tran GV, Nguyen TQ, Dang DV, Nguyen VK, Nguyen CT, et al. Acute febrile myalgia in Vietnam due to trichinellosis following the consumption of raw pork. *Clin Infect Dis*. 2009;49:e79–83. [PubMed](#) [doi:10.1086/605533](https://doi.org/10.1086/605533)

## Section B. Trichinellosis infections acquired in countries different from those where the disease was developed and diagnosed, 1986–2009

Country where the infection was developed and diagnosed	Country where infection was acquired	Period	No. clinical cases	References
Austria	Yugoslavia	2000–2006	10	Austrian Ministry of Health Report, 1986–2009
Czech Republic	Poland	1992–2006	2	National Reference Laboratory for Tissue Helminthosis
	Ukraine		2	
	France		1	
Denmark	Poland	2007	12	Stensvold et al., 2007
France	Algeria	1986	5	Michel et al., 1986
	Turkey	1994–1995	3	Dupouy-Camet et al., 1998
	Kenya		2	
	Greenland	1995	2	Nozais et al., 1996
	Laos	1991	1	Ancelle et al., 2006
	Yugoslavia	1996	1	
	Croatia	1999	1	
	Cameroon	1999	1	
	Spain	2001	1	
	Canada	2004	1	
	Thailand	2004	1	
	Laos	2005	3	
	Canada	2005	8	
	Algeria	2004	1	Nezri et al., 2006
	Laos	2007	1	Dupouy-Camet et al., 2008
Senegal	2009	5	Dupouy-Camet et al., 2009	
Belgium	Canada	2009	1	Houzé et al., 2009
People's Republic of China (Hong Kong Special Administrative Region)			1	
France			4	
Germany			1	
Germany	Poland	2007	3	Schmiedel and Kramme, 2007
Italy	Romania	2008	4	Angheben et al., 2008
Netherlands	Yugoslavia	1992	3	T. Kortbeek, pers. comm.
	Montenegro	1999	5	Pinelli et al., 2004
Total			86	

## References

- Ancelle T, De Bruyne A, Niang M, Poisson DM, Prazuck T, Fur A, et al. Epidémie de trichinellose à *Trichinella nativa* due à la consommation de viande d'ours, France 2005. Bulletin Épidémiologique Hebdomadaire. 2006;14:96–8.
- Angheben A, Mascarello M, Zavarise G, Gobbi F, Monteiro G, Marocco S, et al. Outbreak of imported trichinellosis in Verona, Italy, January 2008. Euro Surveill. 2008;13:pii18891. [PubMed](#)
- Austrian Ministry of Health Report 1986–2009, received by email from Dr. Herbert Auer, Dep. Med. Parasitol., Institute of Specific Prophylaxis and Tropical Medicine Center of Pathophysiology, Infectiology and Immunology Kinderspitalgasse 15 1090 Wien.
- Dupouy-Camet J, Allegretti S, Truong TP. Enquete sur l'incidence de la trichinellose en France (1994–1995). Bulletin Épidémiologique Hebdomadaire. 1998;28:122–3.

- Dupouy-Camet et al. Surveillance de la trichinellose humaine en France. Rapport du CNR des *Trichinella*, 1<sup>er</sup> janvier 2007–31 décembre 2007. 2008.
- Dupouy-Camet J et al. Surveillance de la trichinellose humaine en France. Rapport du CNR des *Trichinella*, 1<sup>er</sup> janvier 2008–31 décembre 2008. 2009.
- Houzé S, Ancelle T, Matra R, Boceno C, Carlier Y, Gajadhar AA, et al. Trichinellosis acquired in Nunavut, Canada in September 2009: meat from grizzly bear suspected. *Euro Surveill.* 2009;14:pii: 19383. [PubMed](#)
- Koortbek T. [email: [Titia.Kortbeek@rivm.nl](mailto:Titia.Kortbeek@rivm.nl)].
- Michel PH, Zurlinden A, Charvillat L, Levenq P, Janin G, Bayle A, et al. Five new cases of trichinosis [in French]. *Presse Med.* 1986;15:2073–4. [PubMed](#)
- National Reference Laboratory for Tissue Helminthosis of the Czech Republic 1986–2009 [email: [Libuse.kolarova@lfl.cuni.cz](mailto:Libuse.kolarova@lfl.cuni.cz)]
- Nezri M, Ruer J, De Bruyne A, Cohen-Valensi R, Pozio E, Dupouy-Camat J. Première observation d'un cas de humain de trichinellose à *Trichinella britovi* en Algérie après consommation de viande de chacal (*Canis aureus*). *Bull Soc Pathol Exot.* 2006;99:94–5. [PubMed](#)
- Nozais JP, Mannevy V, Danis M. Two cases of trichinosis from polar bear (*Thalarctos maritimus*) meat [in French]. *Med Mal Infect.* 1996;26:732–3. [doi:10.1016/S0399-077X\(96\)80107-7](https://doi.org/10.1016/S0399-077X(96)80107-7)
- Pinelli E, Mommers M, Homan W, van Maanen T, Kortbeek LM. Imported human trichinellosis: sequential IgG4 antibody response to *Trichinella spiralis*. *Eur J Clin Microbiol Infect Dis.* 2004;23:57–60. [PubMed](#) [doi:10.1007/s10096-003-1039-7](https://doi.org/10.1007/s10096-003-1039-7)
- Schmiedel S, Kramme S. Cluster of trichinellosis cases in Germany, imported from Poland, June 2007. *Euro Surveill.* 2007;12:E070719.4. [PubMed](#)
- Stensvold CR, Nielsen HV, Mølbak K. A case of trichinellosis in Denmark, imported from Poland, June 2007. *Euro Surveill.* 2007;12:E070809.3. [PubMed](#)



## Section C. References for data on frequency of major clinical signs and symptoms in persons infected during 1986–2009

### African Region

#### Ethiopia

Kefenie H, Bero G. Trichinosis from wild boar meat in Gojjam, north-west Ethiopia. *Trop Geogr Med.* 1992;44:278–80. [PubMed](#)

Kefenie H, Wolde H, Abuhferpo M. Trichinosis from wild boar meat in Arsi, central Ethiopia. *Ethiop Med J.* 1988;26:97–100. [PubMed](#)

### Region of the Americas

#### Canada

Ancelle T, Bruyne A, Nuang M, Poisson DM, Prazuck T, Fur A, et al. Epidemic of *Trichinella nativa* trichinosis caused by consumption of bear meat, France, 2005. *Bull Epidemiol Hebd.* 2006;14:96–8.

Greenbloom SL, Martin-Smith P, Issacs S, Marshall B, Kittle DC, Kain KC, et al. Outbreak of trichinosis in Ontario secondary to the ingestion of wild boar meat. *Can J Public Health.* 1997;88:52–6. [PubMed](#)

Heinzig L. Case reviews—botulism and trichinosis. *EpiNorth.* 1996;8:7.

Houzé S, Ancelle T, Matra R, Boceno C, Carlier Y, Gajadhar AA, et al. Trichinellosis acquired in Nunavut, Canada in September 2009: meat from grizzly bear suspected. *Euro Surveill.* 2009;14: pii: 19383. [PubMed](#)

MacLean JD, Poirier L, Gyorkos TW, Proulx JF, Bourgeault L, Corriveau A, et al. Epidemiologic and serologic definition of primary and secondary trichinosis in the Arctic. *J Infect Dis.* 1992;165:908–12. [PubMed](#) [doi:10.1093/infdis/165.5.908](https://doi.org/10.1093/infdis/165.5.908)

Møller LN, Petersen E, Kapel CMO, Melbye M, Koch A. Outbreak of trichinellosis associated with consumption of game meat in West Greenland. *Vet Parasitol.* 2005;132:131–6. [PubMed](#) [doi:10.1016/j.vetpar.2005.05.041](https://doi.org/10.1016/j.vetpar.2005.05.041)

Nozias JP, Mannevey V, Danis M. Two cases of trichinosis from polar bear (*Thalarctos maritimus*) meat. *Med Mal Infect.* 1996;26:732–3.

Schellenberg RS, Tan B, Irvine JD, Stockdale DR, Gajadhar AA, Serhir B, et al. An outbreak of trichinellosis due to consumption of bear meat infected with *Trichinella nativa*, in 2 northern Saskatchewan communities. *J Infect Dis.* 2003;188:835–43. [PubMed](#) [doi:10.1086/378094](https://doi.org/10.1086/378094)

Serhir B, MacLean JD, Healey S, Forbes L. Outbreak of trichinellosis associated with arctic walrus in northern Canada, 1999. *Can Commun Dis Rep.* 2001; 27–04:31–6.

Viallet J, MacLean JD, Goresky CA, Staudt M, Routhier G, Law C. Arctic trichinosis presenting as prolonged diarrhea. *Gastroenterology.* 1986;91:938–46. [PubMed](#)

### **United States**

Centers for Disease Control and Prevention. Trichinosis—Hawaii. *MMWR Morb Mortal Wkly Rep.* 1987;36:14–6. [PubMed](#)

Centers for Disease Control and Prevention. Trichinellosis associated with bear meat—New York and Tennessee, 2003. *MMWR Morb Mortal Wkly Rep.* 2004;53:606–10. [PubMed](#)

Dworkin MS, Gamble HR, Zarlenga DS, Tennican PO. Outbreak of trichinellosis associated with eating cougar jerky. *J Infect Dis.* 1996;174:663–6. [PubMed](#) [doi:10.1093/infdis/174.3.663](https://doi.org/10.1093/infdis/174.3.663)

Graves T, Harkness J, Crutcher JM. Case report: locally acquired trichinosis in an immigrant from Southeast Asia. *J Okla State Med Assoc.* 1996;89:402–4. [PubMed](#)

Kennedy ED, Hall RL, Montgomery SP, Pyburn DG, Jones JJ. Trichinellosis surveillance—United States, 2002–2007. *MMWR Surveill Summ.* 2009;58:1–7. [PubMed](#)

Landry SM, Kiser D, Overby T, Mays B, Caplen CW. Trichinosis: common source outbreak related to commercial pork. *South Med J.* 1992;85:428–9. [PubMed](#) [doi:10.1097/00007611-199204000-00022](https://doi.org/10.1097/00007611-199204000-00022)

McAuley JB, Michelson MK, Hightower AW, Engeran S, Wintermeyer LA, Schantz PM. A trichinosis outbreak among Southeast Asian refugees. *Am J Epidemiol.* 1992;135:1404–10. [PubMed](#)

McAuley JB, Michelson MK, Schantz PM. Trichinellosis surveillance, United States, 1987–1990. *MMWR CDC Surveill Summ.* 1991;40:35–42. [PubMed](#)

Nelson M, Wright TL, Pierce A, Krogwold RA. A common-source outbreak of trichinosis from consumption of bear meat. *J Environ Health.* 2003;65:16–9, 24. [PubMed](#)

Petri WA, Holsinger JR, Pearson RD. Common-source outbreak of trichinosis associated with eating raw home-butchered pork. *South Med J*. 1988;81:1056–8. [PubMed doi:10.1097/00007611-198808000-00029](#)

Roy SL, Lopez AS, Schantz PM. Trichinellosis surveillance—United States, 1997–2001. *MMWR Surveill Summ*. 2003;52:1–8. [PubMed](#)

### **Mexico**

Alvarez-Chacón R, Riega-Carnero RE, Garcia-Rosales JJ, Wong-Chio M, Cob-Sosa CE. Triquinosis en el niño. Informe de 8 casos. *Bol Med Hosp Infant Mex*. 1992;49:286–90. [PubMed](#)

Hernández M, Ramos-Martinez E, Casco-Sanchez EJ, Morales-Gomez JM, Perez-Matos EM. Triquinos aguda. Epidemia de 166 casos en Ciudad Delicias, Chih. Diagnostico por compression tisular y tinicion. *Gac Med Mex*. 1992;128:45–50. [PubMed](#)

Santos Durán-Ortiz J, García-de la Torre I, Orozco-Barocio G, Martínez-Bonilla G, Rodríguez-Toledo A, Herrera-Zárate L. Trichinosis with severe myopathic involvement mimicking polymyositis. Report of a family outbreak. *J Rheumatol*. 1992;19:310–2. [PubMed](#)

Sartí Gutiérrez EJ, Gutierrez Ospina I, Kopman JS. Brote de triquinos ocurrido en una oficina de gobierno. Mexico, D.F. *Salud Publica Mex*. 1986;28:41–7. [PubMed](#)

Zamora-Chávez A. de la O-Cavazos ME, Bernal-Redondo RM, Berrones-Espericueta D, Vazquez-Antona C. Acute trichinosis in children. Intrafamilial epidemic outbreak in Mexico City [in Spanish]. *Bol Med Hosp Infant Mex*. 1990;47:395–400. [PubMed](#)

### **Argentina**

Ambrosioni J, Cecchini D, Castellaro P, Biscione F, Lloveras S, Orduna T. Trichinellosis: epidemiological, clinical and laboratory aspects. A retrospective study (1994–2003). *Enferm Infecc Microbiol Clin*. 2006;24:440–4. [PubMed doi:10.1157/13091782](#)

Calcagno MA, Teixeira C, Forastiero MA, Coastantino SN, Venturiello SM. Aspectos clinicos, serologicos y parasitologicos de un brote de triquinos humana en Villa Mercedes, San Luis, Argentina. *Medicina (B Aires)*. 2005;65:302–6. [PubMed](#)

Caminoa RA, Carbonell JA, Tapia N, Ledesma M. Brote de triquinos humana en Necochea, Buenos Aires, Argentina. *ACTA Bioquim Clin Latinamericana*. 1995;29:493–8.

Caminoa RA, Ledesma M, Gallicchio O, Sanchez G, Benitez M. Human trichinosis in Buenos Aires province, Argentina during 1998. *Vet-Investigacion Vet*. 2000; 2:101–4.

Nuñez GG, Susana N, Costantino SN, Gentile T, Venturiello SM. Immunoparasitological evaluation of *Trichinella spiralis* infection during human pregnancy: a small case series. *Trans R Soc Trop Med Hyg.* 2008;102:662–8. [PubMed doi:10.1016/j.trstmh.2008.03.009](#)

Siciliano C, Nieto Sosa L, Chalub E, Shictong G. Nuevo brote de triquinosis en Cordoba (Rep. Argentina). *Actual Infectologia (Montevideo).* 1988;4:14–20.

### **Chile**

García E, Mora L, Torres P, Jercic MI, Mercado R. First record of human trichinosis in Chile associated with consumption of wild boar (*Sus scrofa*). *Mem Inst Oswaldo Cruz.* 2005;100:17–8. [PubMed doi:10.1590/S0074-02762005000100003](#)

Zamorano CG, Contreras MC, Espinoza A, Paredes M, Sandoval L, Schulz E, et al. Brote de triquinosis en la Comuna de Purranque, X Region, Chile, Octubre–Noviembre, 1992. *Bol Chil Parasitol.* 1994;49:38–42. [PubMed](#)

## **Eastern Mediterranean Region**

### **Iran**

Kia EB, Meamar AR, Zahabiun F, Soodbakhsh A, Kordbacheh P. An outbreak of human trichinellosis due to the consumption of boar meat infected with *Trichinella* sp. [in Iranian]. *Iran J Infect Dis Trop Med.* 2008;41:35–48.

### **Lebanon**

Haim M, Efrat M, Wilson M, Schantz PM, Cohen D, Shemer J. An outbreak of *Trichinella spiralis* infection in southern Lebanon. *Epidemiol Infect.* 1997;119:357–62. [PubMed doi:10.1017/S0950268897007875](#)

## **European Region**

### **Belarus**

Ivanov KS, Antonov VS, Knysh GG, Khadzaeva AN, Antykova LP, Lavrova VP, et al. The clinical characteristics of 2 outbreaks of trichinellosis [in Russian]. *Med Parazitol (Mosk).* 1990;4:41–2. [PubMed](#)

### **Bulgaria**

Petkova S, Mihov L, Vutova K, Tsenov I, La Rosa G, Pozio E. Epidemiological and clinical patterns of trichinellosis in Bulgaria from 1995 to 2002. *Parasite.* 2008;15:86–8. [PubMed](#)

## Croatia

Cvitović A, Miletic-Medved M, Gjenaro-Margan I. An epidemic of trichinellosis in autumn 2004 in Slavonski Brod [in Croatian]. *Acta Med Croatica*. 2007;61:215–8. [PubMed](#)

Puljiz I, Beus A, Kuzman I, Seiwerth S. Electrocardiographic changes and myocarditis in trichinellosis: a retrospective study of 154 patients. *Ann Trop Med Parasitol*. 2005;99:403–11. [PubMed](#)  
[doi:10.1179/136485905X36307](https://doi.org/10.1179/136485905X36307)

## France

Ancelle T, De Bruyne A, Niang M, Poisson DM, Prazuck T, Fur A, et al. Epidémie de trichinellose à *Trichinella nativa* due à la consommation de viande d'ours, France 2005. *Bulletin Épidémiologique Hebdomadaire*. 2006;14:96–8.

Ancelle T, Dupouy-Camet J, Desenclos JC, Maillot E, Charlet F, Gravelat-Desclaux C, et al. Epidémie de trichinellose (France, 1993) bilan des investigations. *Bulletin Épidémiologique Hebdomadaire*. 1994;29:127–9.

Ancelle T, Dupouy-Camet J, Desenclos JC, Maillot E, Savage-Houze S, Drucker FCJ, et al. A multifocal outbreak of trichinellosis linked to horse meat imported from North America to France in 1993. *Am J Trop Med Hyg*. 1998;59:615–9. [PubMed](#)

Basset D, Thiebaut MM, Pralong F, Abraham B, Moryoussef A, Baldet P, et al. Epidémies familiales de trichinose dues à la consommation de sanglier sauvage dans la région Languedoc. *Bulletin Épidémiologique Hebdomadaire*. 1995;44:195.

Bernard E, Ozouf N, Toussaint-Gari M, Marty P, Pozio E, Le Fichoux Y, et al. Deux épidémies familiales de trichinose. *Med Mal Infect*. 1995;25:611–4. [doi:10.1016/S0399-077X\(05\)81038-8](https://doi.org/10.1016/S0399-077X(05)81038-8)

Dupouy-Camet J, Ancelle T. Surveillance de la trichinellose en France. Rapport du CNR des *Trichinella*, 1 janvier 1999–31 décembre 2002. Rapport du CNR des *Trichinella* 2003.

Dupouy-Camet J, Ancelle T, Talabani H. Surveillance de la trichinellose humaine en France. Rapport du CNR des *Trichinella*, 1 janvier 2009–31 décembre 2009. Rapport du CNR des *Trichinella* 2010.

Durant J, Toussaint-Gari M, Bernard E, Marty P, Le Fichoux Y, Dellamonica P. Epidémie familiale de trichinose. *Sem Hop Paris*. 1991;67:1507–12.

Gaillard T, Martinaud C, Kérébel S, Cellarier G, Muzellec Y, Brisou P. A propos de deux cas de trichinellose à *Trichinella britovi*. *Ann Biol Clin (Paris)*. 2007;65:308–12. [PubMed](#)

Gari-Toussaint M, Bernard E, Quaranta JF, Marty P, Soler C, Ozouf N, et al. First report in France of an outbreak of human trichinellosis due to *Trichinella britovi*. In: Campbell WC, Pozio E, Bruschi F, editors. Trichinellosis. Proceedings of the 8th International Conference on Trichinellosis, 1994. Rome: Instituto Superiore di Sanita Press; 1994. p. 465–8.

Gari-Toussaint M, Tieulié N, Baldin JL, Dupouy-Camet J, Delaunay P, Fuzibet JG, et al. Human trichinellosis due to *Trichinella britovi* in southern France after consumption of frozen wild boar meat. Euro Surveill. 2005;10:117–8. [PubMed](#)

Gari-Toussaint M, Tieulié N, Baldin JL, Marty P, Dupouy-Camet J, Delaunay P, et al. Trichinellose à *Trichinella britovi* dans les Alpes-Maritimes après consommation de viande de sanglier congelée, automne 2003. Bulletin Épidémiologique Hebdomadaire. 2004;21:87–8.

Laurichesse H, Cambon M, Perre D, Ancelle T, Mora M, Hubert B, et al. Outbreak of trichinosis in France associated with eating horse meat. Commun Dis Rep CDR Rev. 1997;7:R69–73.

Nozais JP, Mannevy V, Danis M. Deux cas de trichinose après ingestion de viande d'ours blanc (*Thalarchos maritimus*) au Groenland. Med Mal Infect. 1996;26:732–3. [doi:10.1016/S0399-077X\(96\)80107-7](https://doi.org/10.1016/S0399-077X(96)80107-7)

Ranque S, Faugère B, Pozio E, La Rosa G, Tamburrini A, Pelissier JF, et al. *Trichinella pseudospiralis* outbreak in France. Emerg Infect Dis. 2000;6:543–7. [PubMed](#) [doi:10.3201/eid0605.000517](https://doi.org/10.3201/eid0605.000517)

Roumier M, Milhe P, Hautefort B, Benoist B. Four cases of trichinosis in Camargue (France) by consumption of meat of boar [in French]. Med Mal Infect. 1992;22:947–8. [doi:10.1016/S0399-077X\(05\)80639-0](https://doi.org/10.1016/S0399-077X(05)80639-0)

## Germany

Noeckler K, Reiter-Owona I, Heidrich J, Protz D, Rehmet S, Sinn G, et al. Aspects of clinical features, diagnosis, notification and tracing back referring to *Trichinella* outbreaks in North Rhine-Westphalia, Germany, 1998. Parasite. 2001;8:S183–5. [PubMed](#)

Nothdurft HD, Brommer M, Eichenlaub D, Löscher T. A small outbreak of trichinosis in Germany caused by imported smoked ham [in German]. Dtsch Med Wochenschr. 1995;120:173–6. [PubMed](#) [doi:10.1055/s-2008-1055330](https://doi.org/10.1055/s-2008-1055330)

Rehmet S, Sinn G, Robstad O, David H, Lesser D, Noeckler K, et al. Two outbreaks of trichinellosis in the state of Northrhine-Westfalia, Germany, 1998. Euro Surveill. 1999;4:78–81. [PubMed](#)

Robert Koch Institute. InstitutSurStat@RKI, 2010. Human cases of trichinellosis in Germany from 2001 to 12 May 2010.



## Israel

Marva E, Markovics A, Gdalevich M, Asor N, Sadik C, Leventhal A. Trichinellosis outbreak. *Emerg Infect Dis.* 2005;11:1979–81. [PubMed](#)

## Italy

Frongillo RF, Baldelli B, Pozio E, Crapa G, Di Giuli C, Santirocchi M, et al. Report on an out break of trichinellosis in central Italy. *Eur J Epidemiol.* 1992;8:283–8. [PubMed doi:10.1007/BF00144815](#)

Pagni P, Bindi M, Burzagli L, Melchior M. An outbreak of trichinellosis in central Italy caused by pork meat ingestion. In: Campbell WC, Pozio E, Bruschi F, editors. *Trichinellosis. Proceedings of the 8th International Conference on Trichinellosis, 1994.* Rome: Istituto Superiore di Sanita Press; 1994. p. 487–90.

Pozio E, Cappelli O, Marchesi L, Valeri P, Rossi P. Third outbreak of trichinellosis caused by consumption of horse meat in Italy. *Ann Parasitol Hum Comp.* 1988;63:48–53.

Pozio E, Santagada G, Di Bari C. Outbreak of trichinellosis in southern Italy. *Trans R Soc Trop Med Hyg.* 1986;80:997–8. [PubMed doi:10.1016/0035-9203\(86\)90291-9](#)

Pozio E, Varese P, Gomez Morales MA, Croppo GP, Pelliccia D, Bruschi F. Comparison of human trichinellosis caused by *Trichinella spiralis* and *Trichinella britovi*. *Am J Trop Med Hyg.* 1993;48:568–75. [PubMed](#)

Tamburrini A, Sacchini D, Pozio E. An expected out break of human trichinellosis for the consumption of horsemeat. *Parasite.* 2001;8:S186–7. [PubMed](#)

## Lithuania

Bartuliene A, Liausediene R, Motiejuniene V. Trichinellosis outbreak in Lithuania, Ukmerge region, June 2009. *Euro Surveill.* 2009;14:pii: 19336. [PubMed](#)

## Romania

Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Epidemiological and diagnostic findings during a 16-year-long trichinellosis surveillance in Timis County, Romania. *Vet Parasitol.* 2009;159:328–31. [PubMed doi:10.1016/j.vetpar.2008.10.045](#)

Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Trichinellosis, a threatening and re-emerging disease in a Romanian western county. *Vector Borne Zoonotic Dis.* 2009;9:717–21. [PubMed doi:10.1089/vbz.2008.0204](#)

## Slovakia

Paraličová Z, Kincekova J, Schreter I, Jarcuska P, Dubinsky P, Porubcin S, et al. Outbreak of trichinellosis in eastern Slovakia. *Helminthologia*. 2009;46:209–13. [doi:10.2478/s11687-009-0039-2](https://doi.org/10.2478/s11687-009-0039-2)

Reiterová K, Kincekova J, Snabel V, Marucci G, Pozio E, Dubinsky P. *Trichinella spiralis*–outbreak in the Slovak Republic. *Infection*. 2007;35:89–93. [PubMed doi:10.1007/s15010-007-6122-z](https://pubmed.ncbi.nlm.nih.gov/doi/10.1007/s15010-007-6122-z)

## Spain

Arévalo Velasco A, Bringas MJ, Rodriguez R, Menor A. Description de un brote de triquinosis en la provincia de Salamanca. *Rev Esp Quimioter*. 2009;22:115–6. [PubMed](https://pubmed.ncbi.nlm.nih.gov/)

Benitez JA, Cruz JM, Adrados G, Gonzalez T. Triquinosis: a proposito de 17 casos aparecidos en un mismo brote. *Enferm Infecc Microbiol Clin*. 1987;5:122–3.

Cobo J, Gómez Cerezo J, Medraño JC, Zapico R, Cruz Martinez A, Molina F, et al. Trichinosis. A study of a large outbreak on the Iberian peninsula [in Spanish]. *An Med Interna*. 1991;8:441–4. [PubMed](https://pubmed.ncbi.nlm.nih.gov/)

de la Cruz de Julián I, Diaz Garcia JM, Alvarez Lana P, Garcia Colmenero C. An outbreak of trichinosis in Huerta del Marquesado (the Cañete-Cuenca basic health area). December 1992 to January 1993 [in Spanish]. *Rev Sanid Hig Publica (Madr)*. 1994;68:513–20. [PubMed](https://pubmed.ncbi.nlm.nih.gov/)

de la Torre Cecilia C, Espino Aguilar R, Cardenas Talaveron C, Cañuelo Ruiz O, Garrido Palomo R, Baena Saez J, et al. Triquinosis: asportacion de dos casos. *An Esp Pediatr*. 1989;30:227–8. [PubMed](https://pubmed.ncbi.nlm.nih.gov/)

Perucha González M, Lezaun Larumbe ME, Torres Baile JL, Campo Hernández JM, Bernal Martínez A. Outbreak of trichinosis in various localities of Rioja Baja [in Spanish]. *Rev Sanid Hig Publica (Madr)*. 1987;61:1035–47. [PubMed](https://pubmed.ncbi.nlm.nih.gov/)

Herráez García J, Leon Garcia LA, Lanusse Senderos C, Cortés Blanco M, García Cabanas A. Outbreak of trichinellosis in the region of la Vera (Carceres, Spain) caused by *Trichinella britovi* [in Spanish]. *An Med Interna*. 2003;20:63–6. [PubMed doi:10.4321/S0212-71992003000200003](https://pubmed.ncbi.nlm.nih.gov/doi/10.4321/S0212-71992003000200003)

López Hernández B, Gea Velazquez de Castro MT, Galicia Garcia MD, Sabonet JC. Brote epidemico por *Trichinella britovi* en Granada durante la primavera del 2000. *Rev Esp Salud Publica*. 2001;75:467–73. [PubMed doi:10.1590/S1135-57272001000500007](https://pubmed.ncbi.nlm.nih.gov/doi/10.1590/S1135-57272001000500007)

Mangas Gallardo I, Tello Anchuela O. Vigilancia de la triquinosis en Espana. Temporadas 1989/90–1992/93. *Bol. Epidemiol. Microbiol*. 1994;1:233–4.

Martinez Corral JJ, Esteban Niveiro MJ, Oliet Pala R. Triquinosis producida por el consumo de carne de porcino infestado con *Trichinella spiralis*. Investigation y estudio de un brote alimentario. Salud Publica. 2000;12:36–40.

Serrano R, Lacasa J, Velazquez J, Ziad F, Aznar R. Triquinosis: nuevo brote epidémico por ingesta de embutido de jabali. Enferm Infecc Microbiol Clin. 1989;7:428–31. [PubMed](#)

Tiberio G, Rivero M, Lanzas G, Redin D, Ardanaz E, Fernandez C, et al. Triquinelosis: estudio de dos brotes en Navarra. Enferm Infecc Microbiol Clin. 1997;15:151–3. [PubMed](#)

### **Turkey**

Akkoc N, Kuruuzum Z, Aker S, Yuce A, Onen F, Yapar N, et al. A large-scale outbreak of trichinellosis caused by *Trichinella britovi* in Turkey. Zoonoses Public Health. 2009;56:65–70. [PubMed](#)  
[doi:10.1111/j.1863-2378.2008.01158.x](https://doi.org/10.1111/j.1863-2378.2008.01158.x)

Heper Y, Yilmaztepe F, Komitova R, Akalin H, Vutova K, Helvaci S. A trichinosis outbreak caused by wild boar meat in Turkey. Parasite. 2005;12:191–2. [PubMed](#)

### **United Kingdom**

Milne LM, Bhagani S, Bannister BA, Laitner SM, Moore P, Eza D, et al. Trichinellosis acquired in the United Kingdom. Epidemiol Infect. 2001;127:359–63. [PubMed](#)  
[doi:10.1017/S0950268801005994](https://doi.org/10.1017/S0950268801005994)

## **Western Pacific Region**

### **China**

Cui J, Wang ZQ, Wu F, Jin XX. Epidemiological and clinical studies on an outbreak of trichinosis in central China. Ann Trop Med Parasitol. 1997;91:481–8. [PubMed](#) [doi:10.1080/00034989760842](https://doi.org/10.1080/00034989760842)

Gong XH, Guo WM, Cirendunzhu, Long EK, Ma Y, Bianbazuoma, et al. Investigation on an outbreak of trichinosis and clinical analysis [in Chinese]. Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi. 2008;26:79–80. [PubMed](#)

Lo YC, Hung CC, Lai CS, Wu Z, Nagano I, Maeda T, et al. Human trichinosis after consumption of soft-shelled turtles, Taiwan. Emerg Infect Dis. 2009;15:2056–8. [PubMed](#) [doi:10.3201/eid1512.090619](https://doi.org/10.3201/eid1512.090619)

Luo Z, Li W, Shen L. Report of an outbreak of trichinellosis. Chin J Zoonoses. 2004;20:359–60.

Wang Z, Cai J, Wa F, Jin X. Seven outbreaks of trichinosis in China (1992–1996). J Egypt Soc Parasitol. 1997;27:529–38. [PubMed](#)

Xu B, Chang J, Gao G, Tang B, Hong J, Chen Y, et al. Four outbreaks of human trichinellosis in Henan Province. *Chin Med J (Engl)*. 1995;108:872–4. [PubMed](#)

### **Japan**

Nakamura T, Miura T, Nakaoka T, Nagano I, Takahashi Y, Iwamoto A. A case of trichinellosis with spontaneous remission. *J J A Inf Dis*. 2003;77:839–43.

Shiota T, Arizono N, Yoshioka T, Ishikawa Y, Fujitake J, Fujii H, et al. Imported trichinellosis with severe myositis—report of a case [in Japanese]. *Kansenshogaku Zasshi*. 1999;73:76–82. [PubMed](#)

### **Korea**

Kim E, Pyun RH, Park JH, Kim KH, Choi I, Park HH, et al. Family outbreak of trichinosis after eating a raw meat of wild swine. *Korean J Parasitol*. 2003;35:180–4.

Lee HC, Kim HJ, Oh HY, Kim JH, Kim HG, Lee MS, et al. A case of trichinosis caused by eating a wild badger. *Korean J Med*. 1999;56:134–8.

Sohn WM, Huh S, Chung DI, Pozio E. Molecular identification of Korean *Trichinella* isolates. *Korean J Parasitol*. 2003;41:125–7. [PubMed doi:10.3347/kjp.2003.41.2.125](#)

### **Singapore**

Kurup A, Yew WS, San LM, Ang B, Lim S, Tai GK. Outbreak of suspected trichinosis among travelers returning from a neighboring island. *J Travel Med*. 2000;7:189–93. [PubMed doi:10.2310/7060.2000.00058](#)

### **Laos**

Barennes H, Sayasone S, Odermatt P, De Bruyne A, Hongsakhone S, Newton PN, et al. A major trichinellosis outbreak suggesting a high endemicity of *Trichinella* infection in northern Laos. *Am J Trop Med Hyg*. 2008;78:40–4. [PubMed](#)

Sayasone S, Odermatt P, Vongphrachanh P, Keoluangkot V, Dupouy-Camet J, Newton PN, et al. A trichinellosis outbreak in Borikhamxay Province, Lao PDR. *Trans R Soc Trop Med Hyg*. 2006;100:1126–9. [PubMed doi:10.1016/j.trstmh.2006.01.010](#)

### **Vietnam**

De NV, Toan ND, Son DT, Van P. Report an epidemic area of trichinellosis in Tuan Giao district, Lai Chau province. *Proc 5th Seminar on Food- and Water-Borne Parasitic Zoonoses*. Bangkok, Thailand. 2006 Nov 28–30; p. 48.

Taylor WRJ, Tran GV, Nguyen TQ, Dang DV, Nguyen VK, Nguyen CT, et al. Acute febrile myalgia in Vietnam due to trichinellosis following the consumption of raw pork. *Clin Infect Dis*. 2009;49:e79–83. [PubMed doi:10.1086/605533](#)

## South-East Asia Region

### Thailand

Jongwutiwes S, Chantachum N, Kraivichian P, Siriyasatien P, Putaporntip C, Tamburrini A, et al. First outbreak of human trichinellosis caused by *Trichinella pseudospiralis*. *Clin Infect Dis*. 1998;26:111–5. [PubMed doi:10.1086/516278](#)

Khumjui C, Choomkasien P, Dekumyoy P, Kusolsuk T, Kongkaew W, Chalamaat M, et al. Outbreak of trichinellosis caused by *Trichinella papuae*, Thailand. 2006. *Emerg Infect Dis*. 2008;14:1913–5. [PubMed doi:10.3201/eid1412.080800](#)

Kusolsuk T, Kamonrattanakun S, Wesanonthawech A, Dekumyoy P, Thaenkham U, Yoonuan T, et al. The second outbreak of trichinellosis caused by *Trichinella papuae* in Thailand. *Trans R Soc Trop Med Hyg*. 2010;104:433–7. [PubMed doi:10.1016/j.trstmh.2009.12.005](#)

Limsuwan S, Siriprasert V. A clinical study on trichinosis in Changwat Phayao, Thailand. *Southeast Asian J Trop Med Public Health*. 1994;25:305–8. [PubMed](#)

Watt G, Saisorn S, Jongsakul K, Sakolvaree Y, Chaicumpa W. Blinded, placebo-controlled trial of antiparasitic drugs for trichinosis myositis. *J Infect Dis*. 2000;182:371–4. [PubMed doi:10.1086/315645](#)

### India

Alipuria S, Sangha HK, Singh G, Pandhi S. Trichinosis—a case report. *Indian J Pathol Microbiol*. 1996;39:231–2. [PubMed](#)

Handa R, Aggarwal P, Sarkar C, Vijayarghavan M, Mattewal A, Arya V, et al. A patient with muscle weakness. *J Indian Rheum Assoc*. 2000;8:85–7.

## Section D. Summary of sex and age data on persons with trichinellosis cases, by World Health Organization region, 1986 to 2009

Region/Country	% Males (total no. cases)*	Age, y, of infected persons (no. cases)	References
<hr/>			
African Region			
Ethiopia	100 (28)	Range: 23–25 (3) Mean: 24 (3)	Kefenie et al., 1988 Kefenie and Bero, 1992
<hr/>			
Region of the Americas			
Canada	62.1 (150)	Range: 21–67 (85) Mean: 34.4 (65)	MacLean et al., 1992 Greenbloom et al., 1997 Schellenberger et al., 2003 Serhir et al., 2001 Ancelle et al., 2006
USA	57.5 (632)	Range: 1–87 (412) Mean: 42.0 (126) Median: 37.1 (232)	McAuley et al. 1991 McAuley et al., 1992 Dworkin et al., 1996 Moorhead et al., 1999 Roy et al., 2003 Kennedy et al., 2009
Mexico	35 (59)	Range: 25–44 (59)	Hernández et al., 1992
Chile	60 (667)	Range: 5–70 (667)	Zamorano et al., 1994 Schenone et al., 2002
<hr/>			
Eastern Mediterranean Region			
Lebanon	54 (44)	Range: 10–70 (44) Mean: 33 (44)	Haim et al., 1997
<hr/>			
European Region			
Bulgaria	49 (228)	Range: 1–70 (228)	Petkova et al., 2008
Croatia	57 (200)	Range: 3–67 (200) Mean: 35 (200)	Cvitovic et al., 2007 Venus et al., 2008
Czech Republic	41.9 (31)	Range: 9–68 (31) Mean: 35.9 (31)	National Reference Laboratory for Tissue Helminthosis
France	51.4 (586)	Range: 1–84 (581) Mean: 43.8 (581)	Durant et al., 1991 Roumier et al., 1992 Laurichesse et al., 1997 Bernard et al., 1995 Ancelle et al., 1998 Ranque et al., 2000 Gari-Toussaint et al., 2004 Ancelle et al., 2006 Gaillard et al., 2007 Dupouy-Camet et al., 2003 Dupouy-Camet et al., 2005 Dupouy-Camet et al., 2010
Germany	51.9 (104)	Range: 1–73 (101) Mean: 34.8 (101)	Jansen et al., 2008 Nothdurft et al., 1995 Schmiedel and Kramme, 2007 Nockler et al., 2007
Israel	100 (26)	Mean 32 (26)	Marva et al., 2005
Italy	50.3 (382)	Range: 1–90 (368) Mean: 36.7 (368)	Pozio et al., 1986 Pozio et al., 1988 Frongillo et al., 1992 Pozio et al., 1993 Tamburrini et al., 2001 Neghina et al., 2009
Romania	53.2 (521)	Range: 1– >60 (521) Mean: 31.4 (521)	
Slovakia	63.6 (11)	Range: 16–80 (21) Mean: 40.5 (21)	Reiterová et al., 2007 Paraličová et al., 2009



Spain	57.5 (237)	Range: 2–86 (140) Mean: 40.7 (177)	Benitez et al., 1987 Serrano et al., 1989 Cobo et al., 1991 de la Cruz de Julián et al., 1994 Rodríguez-Osorio et al., 1999 López Hernández et al., 2001 Cortés Blanco et al., 2002 Herráez Garcia et al., 2003 Akkoc et al., 2009
Turkey	52.6 (418)	Range: 1.5–73 (418) Mean: 31.1 (418)	
South-East Asia Region			
Thailand	71 (165)	Range: 7–70 ( 210) Mean: 35.6 ( 208) Median: 34.5 ( 140)	Limsuwan and Siripraesert, 1994 Jongwutiwes et al., 1998 Charkrit, 1998 Khumjui et al., 2008 Kusolsuk et al., 2010
Western Pacific Region			
China	58.2 (802)	Range: 1–90 (482)	Xu et al., 1995 Cui et al., 1997 Wang et al., 1997 Ye et al., 2003 Luo et al., 2004
Laos	47% (111)	Range: 5–69 (111) Median: 34 (21) Mean: 30.4 (90)	Sayasone et al., 2006 Barennes et al., 2008
Singapore	56% (25)	Mean: 22.5 (25)	Kurup et al., 2000
Vietnam	92% (42)	Range: 20–60 ( 42) Mean: 45.4 ( 42)	De et al., 2000

\*Data are from those reports which presented adequate sex and age data of no fewer than 10 cases during 1986–2009.

## References

### African Region

#### Ethiopia

Kefenie H, Bero G. Trichinosis from wild boar meat in Gojjam, north-west Ethiopia. *Trop Geogr Med.* 1992;44:278–80. [PubMed](#)

Kefenie H, Wolde H, Abuherpo M. Trichinosis from wild boar meat in Arsi, central Ethiopia. *Ethiop Med J.* 1988;26:97–100. [PubMed](#)

### Region of the Americas

#### Canada

Ancelle T, Bruyne A, Nuang M, Poisson DM, Prazuck T, Fur A, et al. Epidemic of *Trichinella nativa* trichinosis caused by consumption of bear meat, France, 2005. *Bulletin Épidémiologique Hebdomadaire.* 2006;14:96–8.

Greenbloom SL, Martin-Smith P, Issacs S, Marshall B, Kittle DC, Kain KC, et al. Outbreak of trichinosis in Ontario secondary to the ingestion of wild boar meat. *Can J Public Health.* 1997;88:52–6. [PubMed](#)

MacLean JD, Poirier L, Gyorkos TW, Proulx JF, Bourgeault L, Corriveau A, et al. Epidemiologic and serologic definition of primary and secondary trichinosis in the Arctic. *J Infect Dis.* 1992;165:908–12. [PubMed](#)

Schellenberg RS, Tan B, Irvine JD, Stockdale DR, Gajadhar AA, Serhir B, et al. An outbreak of trichinellosis due to consumption of bear meat infected with *Trichinella nativa* in two northern Saskatchewan communities. *J Infect Dis.* 2003;188:835–43. [PubMed](#)

Serhir B, MacLean JD, Healey S, Forbes L. Outbreak of trichinellosis associated with arctic walrus in northern Canada, 1999. *Can Commun Dis Rep.* 2001; 27–04:31–6.

### **United States**

Dworkin MS, Gamble HR, Zarlenga DS, Tennican PO. Outbreak of trichinellosis associated with eating cougar jerky. *J Infect Dis.* 1996;174:663–6. [PubMed](#)

Kennedy ED, Hall RL, Montgomery SP, Pyburn DG, Jones JJ. Trichinellosis surveillance—United States, 2002–2007. *MMWR Surveill Summ.* 2009;58(9):1–7. [PubMed](#)

McAuley JB, Michelson MK, Schantz PM. Trichinellosis surveillance, United States, 1987–1990. *MMWR CDC Surveill Summ.* 1991;40(3):35–42. [PubMed](#)

McAuley JB, Michelson MK, Hightower AW, Engeran S, Wintermeyer LA, Schantz PM. A trichinosis outbreak among Southeast Asian refugees. *Am J Epidemiol.* 1992;135:1404–10. [PubMed](#)

Moorhead A, Grunenwald PE, Dietz VJ, Schantz PM. Trichinellosis in the United States, 1991–1996: declining but not gone. *Am J Trop Med Hyg.* 1999;60:66–9. [PubMed](#)

Roy SL, Lopez AS, Schantz PM. Trichinellosis surveillance—United States, 1997–2001. *MMWR Surveill Summ.* 2003;52(6):1–8. [PubMed](#)

### **Mexico**

Hernández M, Ramos-Martinez E, Casco-Sánchez EJ, Morales-Gómez JM, Pérez-Matos EM. Acute trichinosis. Epidemics of 166 cases in Delicias City, Chih. Diagnosis with tissue compression and staining [in Spanish]. *Gac Med Mex.* 1992;128:45–50. [PubMed](#)

### **Chile**

Schenone H, Olea A, Schenone H, Contreceras MC, Mercado R, Sandoval L, et al. Epidemiology of trichinosis in Chile from 1991–2000. *Rev Med Chil.* 2002;130:281–5. [PubMed](#)

Zamorano CG, Contreras MC, Espinoza A, Paredes M, Sandoval L, Schulz E, et al. Brote de triquinosis en la Comuna de Purranque, X Region, Chile, Octubre–Noviembre, 1992. *Bol Chil Parasitol.* 1994;49:38–42. [PubMed](#)

### **Eastern Mediterranean Region**

#### **Lebanon**

Haim M, Efrat M, Wilson M, Schantz PM, Cohen D, Shemer J. An outbreak of *Trichinella spiralis* infection in southern Lebanon. *Epidemiol Infect.* 1997;119:357–62. [PubMed](#)

### **European Region**

#### **Bulgaria**

Petkova S, Mihov L, Vutova K, Tsenov I, La Rosa G, Pozio E. Epidemiological and clinical patterns of trichinellosis in Bulgaria from 1995 to 2002. *Parasite.* 2008;15:86–8. [PubMed](#)

#### **Croatia**

Cvitović A, Miletić-Medved M, Gjenero-Margan I. An epidemic of trichinellosis in autumn 2004 in Slavonski Brod [in Croatian]. *Acta Med Croatica.* 2007;61:215–8. [PubMed](#)

Venus M, Puntaric D, Grgic M, Gmajnic R, Miskulin M. The effect of pest control on the incidence of trichinosis in Virovitica-Podravina county, Croatia. *Vet Parasitol.* 2008;156:226–33.

#### **Czech Republic**

National Reference Laboratory for Tissue Helminthosis of the Czech Republic 1986–2009 [email: Libuse.kolarova@lfl.cuni.cz]

#### **France**

Ancelle T, Dupouy-Camet J, Desenclos JC, Maillot E, Savage-Houze S, Drucker FCJ, et al. A multifocal outbreak of trichinellosis linked to horse meat imported from North America to France in 1993. *Am J Trop Med Hyg.* 1998;59:615–9. [PubMed](#)

Ancelle T, De Bruyne A, Niang M, Poisson DM, Prazuck T, Fur A, et al. Epidémie de trichinellose à *Trichinella nativa* due à la consommation de viande d'ours, France 2005. *Bulletin Épidémiologique Hebdomadaire.* 2006;14:96–8.

Bernard E, Ozouf N, Toussaint-Gari M, Marty P, Pozio E, Le Fichoux Y, et al. Deux épidémies familiales de trichinose. *Med Mal Infect.* 1995;25:611–4.

Dupouy-Camet J, Ancelle T. Surveillance de la trichinellose en France. Rapport du CNR des *Trichinella*, 1 janvier 1999–31 décembre 2002. Rapport du CNR des *Trichinella* 2003.

- Dupouy-Camet J, Ancelle T, de Bruyne A. Surveillance de la trichinellose humaine en France. Rapport du CNR des Trichinella, 1 janvier 2002–31 décembre 2004. Rapport du CNR des Trichinella 2005.
- Dupouy-Camet J, Ancelle T, Talabani H. Surveillance de la trichinellose humaine en France. Rapport du CNR des Trichinella, 1 janvier 2009–31 décembre 2009. Rapport du CNR des Trichinella 2010.
- Durant J, Toussaint-Gari M, Bernard E, Marty P, Le Fichoux Y, Dellamonica P. Epidémie familiale de trichinose. *Sem Hop Paris*. 1991;67:1507–12.
- Gaillard T, Martinaud C, Kérébel S, Cellarier G, Muzellec Y, Brisou P. A propos de deux cas de trichinellose à *Trichinella britovi*. *Ann Biol Clin (Paris)*. 2007;65:308–12. [PubMed](#)
- Gari-Toussaint M, Tieulié N, Baldin JL, Marty P, Dupouy-Camet J, Delaunay P, et al. Trichinellose à *Trichinella britovi* dans les Alpes-Maritimes après consommation de viande de sanglier congelée, automne 2003. *Bulletin Épidémiologique Hebdomadaire*. 2004;21:87–8.
- Laurichesse H, Cambon M, Perre D, Ancelle T, Mora M, Hubert B, et al. Outbreak of trichinosis in France associated with eating horse meat. *Commun Dis Rep CDR Rev*. 1997;7:R69–73.
- Ranque S, Faugère B, Pozio E, La Rosa G, Tamburrini A, Pelissier JF, et al. *Trichinella pseudospiralis* outbreak in France. *Emerg Infect Dis*. 2000;6:543–7. [PubMed](#)
- Roumier M, Milhe P, Hautefort B, Benoist B. Four cases of trichinosis in Camargue (France) by consumption of meat of boar [in French]. *Med Mal Infect*. 1992;22:947–8. [doi:10.1016/S0399-077X\(05\)80639-0](https://doi.org/10.1016/S0399-077X(05)80639-0)
- Germany**
- Jansen A, Schöneberg I, Stark K, Nöckler K. Epidemiology of trichinellosis in Germany, 1996–2006. *Vector Borne Zoonotic Dis*. 2008;8:189–96. [PubMed](#)
- Noeckler K, Reiter-Owona I, Heidrich J, Protz D, Rehmet S, Sinn G, et al. Aspects of clinical features, diagnosis, notification and tracing back referring to *Trichinella* outbreaks in North Rhine-Westphalia, Germany, 1998. *Parasite*. 2001;8:S183–5. [PubMed](#)
- Nöckler K, Wichmann-Schauer H, Hiller P, Müller A, Bogner K. Trichinellosis outbreak in Bavaria caused by cured sausages from Romania, January 2007. *Euro Surveill*. 2007; 12:E070823.2. [PubMed](#)
- Nothdurft HD, Brommer M, Eichenlaub D, Löscher T. A small outbreak of trichinosis in Germany caused by imported smoked ham [in German]. *Dtsch Med Wochenschr*. 1995;120:173–6. [PubMed](#)

Schmiedel S, Kramme S. Cluster of trichinellosis cases in Germany, imported from Poland, June 2007. Euro Surveill. 2007;12:E070719.4. [PubMed](#)

### **Israel**

Marva E, Markovics A, Gdalevich M, Asor N, Sadik C, Leventhal A. Trichinellosis outbreak. Emerg Infect Dis. 2005;11:1979–81. [PubMed](#)

### **Italy**

Frongillo RF, Baldelli B, Pozio E, Crapa G, Di Giuli C, Santirocchi M, et al. Report on an outbreak of trichinellosis in central Italy. Eur J Epidemiol. 1992;8:283–8. [PubMed](#)

Pozio E, Santagada G, Di Bari C. Outbreak of trichinellosis in southern Italy. Trans R Soc Trop Med Hyg. 1986;80:997–8. [PubMed](#)

Pozio E, Cappelli O, Marchesi L, Valeri P, Rossi P. Third outbreak of trichinellosis caused by consumption of horse meat in Italy. Ann Parasitol Hum Comp. 1988;63:48–53.

Pozio E, Varese P, Gomez Morales MA, Croppo GP, Pelliccia D, Bruschi F. Comparison of human trichinellosis caused by *Trichinella spiralis* and *Trichinella britovi*. Am J Trop Med Hyg. 1993;48:568–75. [PubMed](#)

Tamburrini A, Sacchini D, Pozio E. An expected out break of human trichinellosis for the consumption of horsemeat. Parasite. 2001;8:S186–7. [PubMed](#)

### **Romania**

Neghina R, Neghina AM, Marincu I, Moldovan R, Iacobiciu I. Epidemiological and diagnostic findings during a 16-year-long trichinellosis surveillance in Timis County, Romania. Vet Parasitol. 2009;159:328–31. [PubMed](#)

### **Slovakia**

Paraličová Z, Kinceková J, Schreter I, Jarcuska P, Dubinsky P, Porubcin S, et al. Outbreak of trichinellosis in eastern Slovakia. Helminthologia. 2009;46:209–13.

Reiterová K, Kinceková J, Snábel V, Marucci G, Pozio E, Dubinsky P. *Trichinella spiralis*–outbreak in the Slovak Republic. Infection. 2007;35:89–93. [PubMed](#)

### **Spain**

Benitez JA, Cruz JM, Adrados G, Gonzalez T. Triquinosis: a proposito de 17 casos aparecidos en un mismo brote. Enferm Infecc Microbiol Clin. 1987;5:122–3.

Cobo J, Gómez Cerezo J, Medraño JC, Zapico R, Cruz Martínez A, Molina F, et al. Trichinosis. A study of a large outbreak on the Iberian peninsula [in Spanish]. *An Med Interna*. 1991;8:441–4.

Cortés Blanco M, García-Cabañas A, Guerra-Peguero F, Ramos-Aceitero JM, Herrera-Guibert D, Martínez-Navarro JF. Outbreak of trichinellosis in Cáceres, Spain, December 2001–February 2002. *Euro Surveill*. 2002;7:136–8 [PubMed](#)

de la Cruz de Julián I, Díaz García JM, Álvarez Lana P, García Colmenero C. An outbreak of trichinosis in Huerta del Marquesado (the Cañete-Cuenca basic health area). December 1992 to January 1993 [in Spanish]. *Rev Sanid Hig Publica (Madr)*. 1994;68:513–20.

Herráez García J, Leon García LA, Lanusse Senderos C, Cortés Blanco M, García Cabañas A. Brote de triquinosis en la comarca de la Vera (Caceres) causado por *Trichinella britovi*. *An Med Interna*. 2003;20:63–6. [PubMed](#)

López Hernández B, Velázquez de Castro MT, Galicia García MD, Sabonet JC. Brote epidemico por *Trichinella britovi* en Granada durante la primavera del 2000. *Rev Esp Salud Publica*. 2001;75:467–73. [PubMed](#)

Rodriguez-Osorio M, Abad JM, De Haro T, Villa-Real R, Gomez Garcia V. Human trichinellosis in southern Spain: serologic and epidemiologic study. *Am J Trop Med Hyg*. 1999;61:834–7. [PubMed](#)

Serrano R, Lacasa J, Velazquez J, Ziad F, Aznar R. Triquinosis: nuevo brote epidémico por ingesta de embutido de jabali. *Enferm Infecc Microbiol Clin*. 1989;7:428–31. [PubMed](#)

### **Turkey**

Akkoc N, Kuruuzum Z, Aker S, Yuce A, Onen F, Yapar N, et al. A large-scale out break of trichinellosis caused by *Trichinella britovi* in Turkey. *Zoonoses Public Health*. 2009;56:65–70. [PubMed](#)

### **South-East Asia Region**

#### **Thailand**

Charkrit S. Study on clinical manifestations of trichinosis in Payao province. *Communicable Diseases Journal*. 1998;24:242–7.

Limsuwan S, Siriprasert V. A clinical study on trichinosis in Changwat Phayao, Thailand. *Southeast Asian J Trop Med Public Health*. 1994;25:305–8. [PubMed](#)

Jongwutiwes S, Chantachum N, Kraivichian P, Siriyasatien P, Putaporntip C, Tamburrini A, et al. First outbreak of human trichinellosis caused by *Trichinella pseudospiralis*. Clin Infect Dis. 1998;26:111–5. [PubMed](#)

Khumjui C, Choomkasien P, Dekumyoy P, Kusolsuk T, Kongkaew W, Chalamaat M, et al. Outbreak of trichinellosis caused by *Trichinella papuae*, Thailand. 2006. Emerg Infect Dis. 2008;14:1913–5. [PubMed](#)

Kusolsuk T, Kamonrattanakun S, Wesanonthawech A, Dekumyoy P, Thaenkham U, Yoonuan T, et al. The second outbreak of trichinellosis caused by *Trichinella papuae* in Thailand. Trans R Soc Trop Med Hyg. 2010;104:433–7. [PubMed](#)

## **Western Pacific Region**

### **China**

Cui J, Wang ZQ, Wu F, Jin XX. Epidemiological and clinical studies on an outbreak of trichinosis in central China. Ann Trop Med Parasitol. 1997;91:481–8. [PubMed](#)

Luo Z, Li W, Shen L. Report of an outbreak of trichinellosis. Chinese Journal of Zoonoses. 2004;20:359–60.

Wang Z, Cui J, Wu F, Jin X. Seven outbreaks of trichinosis in China (1992–1996). J Egypt Soc Parasitol. 1997;27:529–38. [PubMed](#)

Xu B, Chang J, Gao G, Tang B, Hong J, Chen Y, et al. Four outbreaks of human trichinellosis in Henan Province. Chin Med J (Engl). 1995;108:872–4. [PubMed](#)

Ye P, Jianhua H, Zhengkun Z, Desong G, Hao W. The investigation over an outbreak of trichinellosis caused by eating *Sus scrofa*. Chinese Journal of Zoonoses. 2003;19:12.

### **Laos**

Barenes H, Sayasone S, Odermatt P, De Bruyne A, Hongsakhone S, Newton PN, et al. A major trichinellosis outbreak suggesting a high endemicity of *Trichinella* infection in northern Laos. Am J Trop Med Hyg. 2008;78:40–4. [PubMed](#)

Sayasone S, Odermatt P, Vongphrachanh P, Keoluangkot V, Dupouy-Camet J, Newton PN, et al. A trichinellosis outbreak in Borikhamxay Province, Lao PDR. Trans R Soc Trop Med Hyg. 2006;100:1126–9. [PubMed](#)

**Singapore**

Kurup A, Yew WS, San LM, Ang B, Lim S, Tai GK. Outbreak of suspected trichinosis among travelers returning from a neighboring island. *J Travel Med.* 2000;7:189–93. [PubMed](#)

**Vietnam**

De NV, Toan ND, Son DT, Van P. Report an epidemic area of trichinellosis in Tuan Giao district, Lai Chau province. *Proc 5th Seminar on Food- and Water-Borne Parasitic Zoonoses.* Bangkok, Thailand. 2006 Nov 28–30. p. 48.