

5. Vogel U, Claus H, von Müller L, Bunjes D, Elias J, Frosch M. Bacteremia in an immunocompromised patient caused by a commensal *Neisseria meningitidis* strain harboring the capsule null locus (cnl). *J Clin Microbiol.* 2004;42:2898–901. DOI: 10.1128/JCM.42.7.2898-2901.2004
6. Hoang LM, Thomas E, Tyler S, Pollard AJ, Stephens G, Gustafson L, et al. Rapid and fatal meningococcal disease due to a strain of *Neisseria meningitidis* containing the capsule null locus. *Clin Infect Dis.* 2005;40:e38–42. DOI: 10.1086/427875
7. Findlow H, Vogel U, Mueller JE, Curry A, Njanpop-Lafourcade BM, Claus H, et al. Three cases of invasive meningococcal disease caused by a capsule null locus strain circulating among healthy carriers in Burkina Faso. *J Infect Dis.* 2007;195:1071–7. DOI: 10.1086/512084
8. Gagneux SP, Hodgson A, Smith TA, Wirth T, Ehrhard I, Morelli G, et al. Prospective study of serogroup X *Neisseria meningitidis* outbreak in northern Ghana. *J Infect Dis.* 2002;185:618–26. DOI: 10.1086/339010
9. Shao Z, Zhou H, Gao Y, Ren H, Xu L, Kan B, et al. *Neisseria meningitidis* serogroup W135, China. *Emerg Infect Dis.* 2010;16:348–9.
10. Chen C, Zhang TG, He JG, Wu J, Chen LJ, Liu JF, et al. A first meningococcal meningitis case caused by serogroup X *Neisseria meningitidis* strains in China. *Chin Med J (Engl).* 2008;121:664–6.

Address for correspondence: Zhujun Shao, Department of Respiratory Infectious Diseases, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, PO Box 5, Changping, Beijing 102206, People's Republic of China; email: shaozhujun@icdc.cn

# etymologia

## *Pseudoterranova azarasi*

[sü-dō-’ter-ə-nō-və a-zär-a-sē]

From the Greek for false, Latin for earth and new, and Japanese for sea lion. First identified in 1878 as a parasite in pinnipeds by Danish scientist Harald Krabbe, who suggested the name *Ascaris decipiens*, the taxonomic designation for these nematodes changed as knowledge of the life cycles and morphologic features of members of the order Ascaridida expanded. In 1998, molecular examination found *Pseudoterranova decipiens*, long thought to be a monotype, consisted of genetically distinct sibling species. Mattiucci et al. proposed *Pseudoterranova azarasi* for 1 of the 5 sibling species, incorporating part of the name *Porrocaecum azarasi*, previously considered a synonym for *Pseudoterranova decipiens*.

Sources: Berland B. *Anisakis* spp. In: Akuffo H, Linder E, Ljungström I, Wahlgren M, editors. Parasites of the colder climates. London: Taylor & Francis; 2003. p. 160–8; Krabbe H. On the ascarides of the seals and toothed whales. *Annals and Magazine of Natural History.* 1878; 5-S, 2(11):430–2; Mattiucci S, Paggi L, Nascetti G, Ishikura H, Kikuchi K, Sato N, et al. Allozyme and morphological identification of *Anisakis*, *Contraaecum* and *Pseudoterranova* from Japanese waters (Nematoda, Ascaridoidea). *Syst Parasitol.* 1998;40:81–92. DOI: 10.1023/A:1005914926720

Manage your email to focus on content of interest to you.



GovDelivery

[www.cdc.eid.ncidod/eid/subscrib.htm](http://www.cdc.eid.ncidod/eid/subscrib.htm)