

### Acknowledgments

We thank Irina Suomalainen, Antonella Rossati, Diego Brustia, Maura Bandi, Letizia Krizmancic, and Vittorio Quaglia for assistance and the Hospital District of Helsinki and Uusimaa and Academy of Finland for financial support.

**Paolo Ravanini, Eili Huhtamo,  
Essi Hasu, Felicita Rosa,  
Stefano Costantino,  
Maria G. Crobu, Valentina Ilaria,  
Anna M. Nicosia,  
Pietro L. Garavelli,  
and Olli Vapalahti**

Author affiliations: Azienda Ospedaliero-Universitaria Maggiore della Carità, Novara, Italy (P. Ravanini, F. Rosa, S. Costantino, M.G. Crobu, V. Ilaria, A.M. Nicosia, P.L. Garavelli); and University of Helsinki, Helsinki, Finland (E. Huhtamo, E. Hasu, O. Vapalahti)

DOI: 10.3201/eid1705.101626

### References

- Jelinek T. Trends in the epidemiology of dengue fever and their relevance for importation to Europe. *Euro Surveill.* 2009;14:pii:19250.
- Scaramozzino N, Crance JM, Jouan A, DeBriel DA, Stoll F, Garin D. Comparison of flavivirus universal primer pairs and development of a rapid, highly sensitive heminested reverse transcription-PCR assay for detection of flaviviruses targeted to a conserved region of the NS5 gene sequences. *J Clin Microbiol.* 2001;39:1922–7. doi:10.1128/JCM.39.5.1922-1927.2001
- Huhtamo E, Uzcátegui NY, Siikamäki H, Saarinen A, Piiparinen H, Vaheeri A, et al. Molecular epidemiology of dengue virus strains from Finnish travelers. *Emerg Infect Dis.* 2008;14:80–3. doi:10.3201/eid1401.070865
- Lanciotti RS, Lewis JG, Gubler DJ, Trent DW. Molecular evolution and epidemiology of dengue-3 viruses. *J Gen Virol.* 1994;75:65–75. doi:10.1099/0022-1317-75-1-65
- Zaki A, Perera D, Jahan SS, Cardoso MJ. Phylogeny of dengue viruses circulating in Jeddah, Saudi Arabia: 1994 to 2006. *Trop Med Int Health.* 2008;13:584–92. doi:10.1111/j.1365-3156.2008.02037.x
- Jimenez-Lucho VE, Fisher EJ, Saravolatz LD. Dengue with hemorrhagic manifestations: an imported case from the Middle East. *Am J Trop Med Hyg.* 1984;33:650–3.
- ProMED-mail. Dengue/DHF update 2010 [cited 2010 Dec 19]. <http://www.promed-mail.org>, archive nos. 20100510.1528, 20100720.2435, 20100810.2726, and 20100826.3010
- Messer WB, Gubler DJ, Harris E, Sivananthan K, de Silva AM. Emergence and global spread of a dengue serotype 3, subtype III virus. *Emerg Infect Dis.* 2003;9:800–9.
- La Ruche G, Souares Y, Armengaud A, Peloux-Petiot F, Delaunay P, Despres P, et al. First two autochthonous dengue virus infections in metropolitan France, September 2010. *Euro Surveill.* 2010;15:pii:19676.
- Schmidt-Chanasit J, Haditsch M, Schoneberg I, Gunther S, Stark K, Frank C. Dengue virus infection in a traveller returning from Croatia to Germany. *Euro Surveill.* 2010;15:pii:19677.

Address for correspondence: Paolo Ravanini, AOU Maggiore della Carità–Laboratorio di Virologia Molecolare, Via Cottolengo, 2 Galliate 28069, Italy; email: paolo.ravanini@gmail.com

## Strongyloidiasis in Man 75 Years after Initial Exposure

**To the Editor:** Strongyloidiasis, caused by the roundworm *Strongyloides stercoralis*, affects 100–200 million persons worldwide (1) and is endemic to Southeast Asia, sub-Saharan Africa, Latin America, and the southeastern United States (2). Endogenous autoinfection enables this nematode to develop into its host, which leads to the persistence of chronic infection several decades after a person has left a disease-endemic area (3). We encountered a case of prolonged strongyloidiasis with an infection going back >75 years.

An 83-year-old man who lived in Paris and had no medical history sought treatment for fatigue and

weight loss. Results of a clinical examination were normal. Laboratory investigations showed mild hyperleukocytosis ( $12 \times 10^9$  cells/L) with hyper eosinophilia ( $2.4 \times 10^9$  cells/L), which was observed for >3 months. Results of stool examinations were negative for parasites. Because of the patient's poor condition, cancer or hematologic malignancy were suspected, but results of various examinations, including a computed tomography scan of the body and bone marrow aspiration, were normal.

After several weeks, the patient disclosed that he had spent a few years in Vietnam >75 years ago. The only other travel abroad reported by the patient was a 10-day stay in a tourist hotel in the Canary Islands 15 years before this illness. At this point in his assessment, results of serologic testing were positive for *Strongyloides* spp., and a new stool examination showed *Strongyloides* larvae. Serologic test results were negative for human T-cell lymphotropic virus type 1. The patient received 2 doses of 12 mg of ivermectin within 15 days and fully recovered. Hyper eosinophilia and *Strongyloides* larvae in feces disappeared.

*S. stercoralis* roundworms are ubiquitous intestinal parasites, endemic to tropical and subtropical regions. The larvae can develop into filariform larvae, which can penetrate the human skin and migrate through circulation to the lungs before settling in the intestine. In the human host, adult parasites may be generated by parthenogenesis in the mucosa of the small intestine. The resultant larvae can also penetrate the skin or the intestinal mucosa to establish a cycle of repeated endogenous reinfection. The parasite may then cause a long-lived autoinfection in the host, leading to chronic infection that can last for several decades (3,4).

Immunocompetent persons are usually asymptomatic and periodically exhibit eosinophilia. In

immunocompromised patients, the endogenous autoinfection cycle may result in the overproduction and dissemination of larvae into intestinal and extraintestinal tissues, including the central nervous system, leading to the hyperinfection syndrome which can be lethal (5). Most cases (96%) occur in immigrants, but some have been described in patients with a history of travel, sometimes many years previously. *S. stercoralis* infections have been reported up to 65 years after initial exposure in veterans who served in Asia during World War II (4,6).

Although our patient exhibited poor general condition, he likely did not experience hyperinfection syndrome because he was not immunosuppressed, and he completely recovered after receiving standard ivermectin treatment. That the patient was originally infected in the Canary Islands seems improbable, although a low level of transmission exists in rural and disadvantaged areas in continental Spain, Portugal, and Italy (7). We did not find evidence of *Strongyloides* spp. transmission in the Canary Islands. In particular, the patient stayed in a high-status tourist hotel for a short period, and he never walked in bare feet. He was probably infected when he lived in Vietnam.

This case highlights the importance of systematically considering chronic strongyloidiasis when seeking a diagnosis for persistent hypereosinophilia, even in patients with no underlying disease, and the value of systematically obtaining any history of travel in disease-endemic areas even if it occurred many years previously. The endogenous autoinfection cycle can possibly persist for a lifetime. In addition, systematic examination of stool samples should be carried out, and ivermectin should be given when an immunosuppressive drug is required in a patient who has a history of travel to, or residence in, an area to which strongyloidiasis is endemic.

**Virginie Prendki, Pierre Fenaux,  
Rémy Durand, Marc Thellier,  
and Olivier Bouchaud**

Author affiliations: Hôpital Jean Verdier, Bondy, France (V. Prendki); Hôpital Avicenne, Bobigny, France (P. Fenaux, R. Durand, O. Bouchaud); Université Léonard de Vinci, Bobigny (R. Durand, O. Bouchaud); and Hôpital La Pitié Salpêtrière, Paris, France (M. Thellier)

DOI: 10.3201/eid1705.100490

**References**

1. Genta RM. Global prevalence of strongyloidiasis: critical review with epidemiologic insights into the prevention of disseminated disease. *Rev Infect Dis.* 1989;11:755–67. doi:10.1093/clinids/11.5.755
2. Berk SL, Verghese A, Alvarez S, Hall K, Smith B. Clinical and epidemiologic features of strongyloidiasis. A prospective study in rural Tennessee. *Arch Intern Med.* 1987;147:1257–61. doi:10.1001/archinte.147.7.1257
3. Grove DI. Human strongyloidiasis. *Adv Parasitol.* 1996;38:251–309. doi:10.1016/S0065-308X(08)60036-6
4. Gill GV, Beeching NJ, Khoo S, Bailey JW, Partridge S, Blundell JW, et al. A British Second World War veteran with disseminated strongyloidiasis. *Trans R Soc Trop Med Hyg.* 2004;98:382–6. doi:10.1016/j.trstmh.2003.11.002
5. Marcos LA, Terashima A, Dupont HL, Gotuzzo E. *Strongyloides* hyperinfection syndrome: an emerging global infectious disease. *Trans R Soc Trop Med Hyg.* 2008;102:314–8. doi:10.1016/j.trstmh.2008.01.020
6. Robson D, Beeching NJ, Gill GV. *Strongyloides* hyperinfection syndrome in British veterans. *Ann Trop Med Parasitol.* 2009;103:145–8. doi:10.1179/136485909X385009
7. Román-Sánchez P, Pastor-Guzman A, Moreno-Guillen S, Igual-Adell R, Suner-Generoso S, Tornero-Estebanez C. High prevalence of *Strongyloides stercoralis* among farm workers on the Mediterranean coast of Spain: analysis of the predictive factors of infection in developed countries. *Am J Trop Med Hyg.* 2003;69:336–40.

Address for correspondence: Virginie Prendki, Service de Médecine Interne, Hôpital Jean Verdier, Ave du 14 juillet, 93140 Bondy, France; email: virginie.prendki@jvr.aphp.fr

*Anaplasma phagocytophilum*  
Infection in Ticks,  
China–Russia  
Border

**To the Editor:** *Anaplasma phagocytophilum*, an emerging human pathogen of public health importance, is transmitted to humans most commonly by tick bites (1). The agent has been detected in various species of *Ixodes* ticks around the world (2) and in *Dermacentor silvarum* ticks in northeastern People's Republic of China (3), where 3 *A. phagocytophilum* strains were isolated from wild and domestic animals (4). In the Asiatic region of Russia adjacent to China, *A. phagocytophilum* was identified in *Ixodes persulcatus* ticks, and *A. bovis* in *Haemaphysalis concinna* ticks (5). Human granulocytic anaplasmosis was reported in the southern area of the Russian Far East that borders China (6). The objectives of this study were to investigate the prevalence of *A. phagocytophilum* in ticks collected from the China–Russia border and to characterize the agent by molecular biology techniques.

During May–June 2009, host-seeking ticks were collected by flagging vegetation of grassland or woodland along the China–Russia border. Attached ticks were collected from sheep and goats in Hunchun, and from dogs in Suifenhe (Table). All ticks were identified by morphologic features to the species level and the developmental stage by 2 entomologists (Y. Sun and R.-M. Xu). DNA was extracted from tick samples by using Tissue DNA Extract kit (Tiangen Biotechnology Inc., Beijing, China), following the instructions of the manufacturer. Nested PCR was performed to amplify partial citrate synthase gene (*gltA*) of *A. phagocytophilum* as previously described (7). To avoid possible contamination, DNA extraction, the