

Seroepidemiology of Saffold Cardiovirus Type 2

To the Editor: Saffold virus (SAFV) is a new human virus belonging to the genus *Cardiovirus* of the family *Picornaviridae* (1–6). The virus has also been named human Theiler's-like cardiovirus (4). To date, 8 SAFV genotypes are known, based on molecular variation in the P1 region, which codes for the viral capsid (5). SAFVs are detected in respiratory and fecal samples from infants and children <6 years of age. The virus is generally detected at low frequency (0.5%–3%) in the general population, but in Afghanistan and Pakistan higher incidences have been reported (10%–12%) (5). Detection is mainly based on molecular techniques because virus isolation is cumbersome. Other than gastroenteritis, no clear disease manifestations are known (1–8), although recently, researchers in Japan have suggested that for a select group of children, SAFV infection may cause pharyngitis and tonsillitis (7). In rodents, however, cardioviruses are serious pathogens that can cause myocarditis, meningoencephalitis, and pancreatitis.

Previously, we have shown that SAFV-3 infections are ubiquitous and are transmitted early in life (6). This conclusion was based on antibody testing by virus neutralization, a highly specific test that can discriminate

serotypes within a single species, but formal proof for existence of serotypes was lacking. Extrapolated to other genotypes, the finding would indicate a high infection rate for SAFVs, which is at odds with the low detection frequency reported in most studies. Most SAFV-2 isolates, from our laboratory and others, grow poorly in cell culture, which hampers reading of neutralization. Recently, however, a SAFV-2 strain was isolated in Finland that grows well in HeLa cells and shows clear cytopathic effects (SAFV-2-FIN2008, GenBank accession no. FR682076; S. Blomqvist et al., unpub. data). This strain enabled us to set up a virus neutralization test similar to that described for SAFV-3 (6). For comparison, we used strain SAFV-3(NL2007) (GenBank accession no. FM207487), which was isolated in Nijmegen (6). The virus neutralization test was performed on HeLa cells with 100 TCID₅₀ of virus per serum dilution. Virus–serum mixtures were incubated for 1 hour at 37°C and overnight at 7°C to stabilize virus–antibody complexes (6). Human serum samples submitted previously were tested by using 3-fold dilution steps and duplicate testing (6). The results are presented in the Table. A low seroprevalence was found for SAFV-2 and SAFV-3 at 9 months of age, which is at the nadir of immunoglobulin G levels in infants. At the age of 24 months, a high seroprevalence of antibodies was found in the Netherlands for SAFV-2 and -3, pointing to early acquisition of infection with both strains. In Finland,

the seroprevalence for SAFV-2 and -3 was lower in young children, which suggests somewhat lower infection rate similar to what has been reported for enteroviruses (9). A high seroprevalence of 97%–100% was found in persons >4 years of age in Cameroon, Indonesia, and the Netherlands (Table). A high prevalence of SAFV-2 antibodies was recently also reported in blood donors from the United States (8). Thus, the seroepidemiology of SAFV-2 is similar to that of SAFV-3. As depicted in the last 3 columns of the Table, on several occasions serum samples independently neutralized either SAFV-2 or SAFV-3, which suggests that the viruses behave as different serotypes.

In conclusion, SAFV-2 and SAFV-3 show an almost identical epidemiologic pattern with infection acquired early in life and with a high seroprevalence in different continents. The outcome is concordant with universal occurrence of infection by both genotypes. Because several times there was a clear discrepancy between antibody titers against one or the other of the 2 genotypes, we conclude that they behave as separate serotypes, although weak cross-reactivity (below the detection limit of 1:15) cannot be excluded. Accepting that SAFV genotypes correspond with existence of different serotypes, the infection rate in the first years of life must be quite high, similar to that for human parechoviruses and enteroviruses, which are found in 15%–18% of stool samples from children <5 years of age (10). The

Table. SAFV-neutralizing antibodies in blood samples from humans of different ages and from different geographic regions*

Country	No. samples	Patient age†	Years collected	No. (%) samples				
				SAFV-2 pos‡	SAFV-3 pos‡	SAFV pos§	SAFV-2 pos + SAFV-3 neg	SAFV-2 neg + SAFV-3 pos
Netherlands	29	9 mo	2006–2007	15 (52)	4 (14)	15 (52)	11	0
Netherlands	26	24 mo	2006–2007	21 (81)	20 (77)	25 (96)	5	4
Netherlands	30	18–39 y	2004	30 (100)	29 (97)	30 (100)	1	0
Finland	30	2–2.5 y	1997–1998	10 (33)	21 (70)	24 (80)	3	14
Cameroon	29	5–15 y	1997	28 (97)	28 (97)	29 (100)	1	1
Indonesia	30	4–40 y	1997–1998	30 (100)	30 (100)	30 (100)	0	0

*SAFV, Saffold virus; pos, positive; neg, negative.

†Samples were collected from patients at this age or within this age range.

‡Titer ≥15.

§Cumulative positive for antibodies against SAFV-2 and/or SAFV-3.

low SAFV detection rate remains thereby difficult to explain.

The outcome can be explained by a short duration of virus excretion in stool, which, however, is unlikely for an infection spreading by the fecal–oral route. Alternatively, it may be that the virus is unstable in stool and rapidly degrades, such that fecal samples are inadequate for diagnosis of the infection. Other specimens, however, such as respiratory samples, yielded also low numbers of positive findings (4). Remarkably, the study with a high prevalence of positive stool samples made use of primers selected in a conserved region of 2C helicase (5), whereas other studies used primers in the 5' noncoding region (3,4). Hence, a difference in sensitivity between the different PCRs may be responsible for the discrepancy between seroepidemiology and the low diagnostic yield by PCR. This discrepancy, however, awaits further investigation.

**Jochem Galama,
Kjerstin Lanke, Jan Zoll,
Merja Roivainen,
and Frank van Kuppeveld**

Author affiliations: Radboud University Nijmegen Medical Center, Nijmegen, the Netherlands (J. Galama, K. Lanke, J. Zoll, F. van Kuppeveld); Nijmegen Center for Molecular Life Sciences, Nijmegen (J. Galama, K. Lanke, J. Zoll, F. van Kuppeveld); Nijmegen Institute for Infection, Inflammation and Immunity, Nijmegen (J. Galama, K. Lanke, J. Zoll, F. van Kuppeveld); and National Institute for Health and Welfare, Helsinki, Finland (M. Roivainen)

DOI: 10.3201/eid1708.101953

References

1. Jones MS, Lukashov VV, Ganac RD, Schnurr DP. Discovery of a novel human picornavirus in a stool sample from a pediatric patient presenting with fever of unknown origin. *J Clin Microbiol.* 2007;45:2144–50. doi:10.1128/JCM.00174-07
2. Abed Y, Boivin G. New Saffold cardioviruses in 3 children, Canada. *Emerg Infect Dis.* 2008;14:834–6.
3. Drexler JF, de Souza Luna LK, Stöcker A, Silva Almeida P, Medrado Ribeiro TC, Petersen N, et al. Circulation of three lineages of a novel Saffold cardiovirus in humans. *Emerg Infect Dis.* 2008;14:1398–405. doi:10.3201/eid1409.080570
4. Chiu CY, Greninger AL, Kanada K, Kwok T, Fischer KF, Runckel C, et al. Identification of cardioviruses related to Theiler's murine encephalomyelitis virus in human infections. *Proc Natl Acad Sci U S A.* 2008;105:14124–9. doi:10.1073/pnas.0805968105
5. Blinkova O, Kapoor A, Victoria J, Naeem A, Shaikat S, Sharif S, et al. Cardioviruses are genetically diverse and common enteric infections in South Asian children. *J Virol.* 2009;83:4631–41. doi:10.1128/JVI.02085-08
6. Zoll J, Erkens Hulshof S, Lanke K, Verduyn Lunel F, Melchers WJ, Schoondermark-van de Ven E, et al. Saffold virus, a human Theiler's-like cardiovirus, is ubiquitous and causes infection early in life. *PLoS Pathog.* 2009;5:e1000416. doi:10.1371/journal.ppat.1000416
7. Itagaki T, Abiko C, Aoki Y, Ikeda T, Mizuta K, Noda M, et al. Saffold cardiovirus infection in children associated with respiratory disease and its similarity to Coxsackievirus infection. *Ped Infect Dis J.* 2011 Mar 7; [Epub ahead of print].
8. Chiu CY, Greninger AL, Chen EC, Haggerty TD, Personnet J, Delwart E, et al. Cultivation and serological characterization of the human Theiler's-like cardiovirus associated with diarrheal disease. *J Virol.* 2010;84:4407–14. doi:10.1128/JVI.02536-09
9. Viskari H, Ludvigsson J, Uibo R, Salur L, Marciulionyte D, Hermann R, et al. Relationship between the incidence of type 1 diabetes and maternal enterovirus antibodies: time trends and geographical variation. *Diabetologia.* 2005;48:1280–7. doi:10.1007/s00125-005-1780-9
10. Benschop K, Thomas X, Serpenti C, Molenkamp R, Wolthers K. High prevalence of human parechovirus (HPeV) genotypes in the Amsterdam region and identification of specific HPeV variants by direct genotyping of stool samples. *J Clin Microbiol.* 2008;46:3965–70. doi:10.1128/JCM.01379-08

Address for correspondence: Jochem Galama, Department of Medical Microbiology, Radboud University Nijmegen Medical Center, PO Box 9101, 6500HB, Nijmegen, the Netherlands; email: j.galama@mmb.umcn.nl

Alkhurma Hemorrhagic Fever in Travelers Returning from Egypt, 2010

To the Editor: The report of 2 visitors from Italy being infected by Alkhurma hemorrhagic fever virus (AHFV) in southeastern Egypt near the border with Sudan (1) provides useful data to help clarify the evolutionary origin of these tick-borne flaviviruses. AHFV was first isolated in Saudi Arabia and is associated with camel ticks (2). It is a genetically close relative of Kyasanur Forest disease virus, which was first isolated in India in 1957. Following the original isolation of Kyasanur Forest disease virus, there was no clear explanation for its apparent isolation in the Indian forests. Indeed, its subsequent discovery in southern China (3) suggested that migratory birds might carry the infected ticks to or from that region.

The most likely explanation for these outbreaks of hemorrhagic disease now begins to fit a pattern that can be interpreted in terms of the diseases' evolutionary origin in Africa. Thousands of animals are annually transported from Africa and other countries to Mecca, Saudi Arabia, to meet the human demand for food and transport during the Hajj. Many of these animals, including camels, are infested with ticks that may carry AHFV and thus provide the source of this human infectious agent. Phylogenetic evidence had previously suggested that the tick-borne encephalitic flavivirus serocomplex originated in Africa and gradually evolved and dispersed across the Northern Hemisphere of the Old World (4,5). This concept is totally consistent with the discoveries of AHFV in Saudi Arabia and now in southeastern Egypt. Thus, Africa is a likely source of infected ticks that are regularly moved between Africa