

Human Herpesvirus 8 Seroprevalence, China

Tiejun Zhang, Xiaodan Shao, Yue Chen,
Tao Zhang, Veenu Minhas, Charles Wood,
and Na He

To summarize the seroprevalence of human herpesvirus 8 (HHV-8) in mainland China, we conducted a systematic review and meta-analysis based on available literature. Data show that differences in HHV-8 prevalence vary considerably among different ethnic groups and geographic regions. Blood-borne transmission could be a potential route for HHV-8 infection in China.

Human herpesvirus 8 (HHV-8) is the infectious etiologic agent associated with Kaposi sarcoma, primary effusion lymphoma, and multicentric Castleman disease. Worldwide seroprevalence of HHV-8 varies: generally low to moderate for populations in Western countries and Asia (1–4) but as high as 50% for the general population in sub-Saharan Africa and higher for HIV-positive populations (5–7). The transmission modes of HHV-8 may also differ in different geographic areas and subpopulations; sexual and nonsexual transmission have been described (8–10). Blood-borne transmission may exist, especially among intravenous drug users (IVDUs) and blood recipients (11).

The Ministry of Health of China, the United Nations Program on HIV/AIDS, and the World Health Organization estimate that ≈320,000 HIV/AIDS cases have been reported in China (12). However, the epidemiologic characteristics of HHV-8 infection, a severe HIV/AIDS opportunistic infection, have not been well described for China. Therefore, we conducted a systematic review and meta-analysis on the basis of available data for HHV-8 epidemiology from mainland China to have a better understanding of the prevalence, variation, and factors associated with its transmission.

Author affiliations: Fudan University, Shanghai, People's Republic of China (T. Zhang, X. Shao, Tao Zhang, V. Minhas, C. Wood, N. He); University of Nebraska-Lincoln, Lincoln, Nebraska, USA (T. Zhang, V. Minhas, C. Wood); and University of Ottawa, Ottawa, Ontario, Canada (Y. Chen)

DOI: <http://dx.doi.org/10.3201/eid1801.102070>

The Study

A comprehensive literature search of published studies indexed in global and databases in China during 1995–2010 was conducted. Initially, 125 reports published in English and 223 in Chinese concerning the seroprevalence in mainland China were identified. Among them, 85 articles published in England and 178 articles published in China were excluded after title and abstract screening. After reading the full text, we excluded another 33 English and 26 Chinese articles. Finally, 26 publications were included in this systematic review and have been summarized in online Technical Appendix Table 1 (wwwnc.cdc.gov/EID/pdfs/10-2070-Techapp.pdf). These studies were cross-sectional and were conducted in 8 of the 34 provinces. A substantial number (35.5%) of these studies were conducted in the Xinjiang Uygur Autonomous Region. Most samples tested were serum or plasma with few exceptions (1 whole blood, 1 peripheral blood mononuclear cells); sample sizes ranged from 37 to 4,461 (median 242, interquartile range 199–520). Overall, 18,547 participants were involved in the present analysis, and among them 15,913 were from the general population, 1,970 were immunocompromised patients, and 664 were IVDUs. Laboratory methods for all included studies were reported (19 detected HHV-8 by ELISA, 3 by PCR, and 4 by immunofluorescent assay).

The prevalence of HHV-8 pooled from reviewed studies was 11.3% (95% CI 7.2–15.5) for the general population, 22.2% (95% CI 12.7–31.8) for immunocompromised patients, and 31.2% (95% CI 27.7–34.7) for IVDUs. The prevalence among the general population was found to be the lowest in Guangdong Province and the highest in Xinjiang Province. A similar regional variation was found for immunocompromised persons. Among IVDUs, the prevalence was 34.3% (95% CI 28.3–40.3) in Zhejiang and 29.6% (95% CI 25.3–33.9) in Xinjiang Uygur Autonomous Region (online Technical Appendix Table 2; Figure).

Five studies, including 4,637 persons of Han ethnicity and 4,011 persons of ethnic minorities (2,040 Uygur, 1,169 Kazak, 200 Khalkas, 173 Hue, and 429 other) conducted in the Xinjiang Uygur Autonomous Region were analyzed for association of ethnicity with HHV-8 prevalence (online Technical Appendix Figure, panel A). The risk was significantly lower for the Han group than for other ethnic groups (odds ratio [OR] = 0.59, 95% CI 0.55–0.76). For the Han group, the pooled prevalence of HHV-8 in Xinjiang Uygur Autonomous Region was significantly higher when compared with that for other regions, 14.4% (95% CI 9.0–19.8) versus 6.4% (95% CI 4.1–8.6). Ten combined studies, with 5,716 male and 4,708 female participants, respectively, were included in meta-analysis of association between sex and HHV-8 infection (online Technical Appendix Figure, panel B).

There was no significant difference between the sexes: pooled OR 0.94 (95% CI 0.84–1.04).

Seven studies, with 863 HIV-positive patients and 3,438 negative controls, were included in the analysis. All studies yielded a significant difference in HHV-8 infection between HIV-positive and HIV-negative participants; ORs for individual studies ranged from 1.50 to 4.27, and the pooled OR was 2.97 (95% CI 2.22–3.97) (online Technical Appendix Figure, panel C). However, a significant publication bias was detected (Egger test $p = 0.013$; Begg test $p = 0.016$). A visual inspection of the funnel plot suggested that some large or small studies with negative or null results were not published (data not shown).

Few studies were designed to address the issue of possible transmission routes among the population of China. Six studies had information on possible blood transmission. Two blood transfusion studies and 4 studies of IVDUs included 837 persons who reported having been exposed to blood contact i.e., needle sharing and 1,397 who were never exposed (online Technical Appendix Figure, panel D). Substantial heterogeneity ($I^2 87%$, $p < 0.001$, by test for heterogeneity) was detected among those studies; therefore, a random-effects model was used to estimate the OR. No publication bias was detected (Begg test $p = 0.707$; Egger test $p = 0.363$). OR showed a marginal association of HHV-8 prevalence with blood transfusion (OR 2.01, 95% CI 0.89–4.56) for possible blood transmission.

Conclusions

This systematic review indicated that HHV-8 prevalence in China varies in different regions. Pooling of data from 26 studies provided us with a large sample size, which is one of the strengths of the study. Also, we included studies that were published in the Chinese language and were not accessible to the international community. The results of this meta-analysis show that HHV-8 prevalence was higher in the Xinjiang Uygur Autonomous Region than other areas in general and among high-risk populations. Historically, Xinjiang Uygur Autonomous Region has been regarded as an area in which Kaposi sarcoma is endemic (13). Notably, geographic variations of HHV-8 infection within China are not well known and need to be investigated as well.

It has been well documented that HHV-8 prevalence is higher among HIV-infected persons (14,15). In mainland China, we found a 3-fold increase in HHV-8 infection among persons with HIV compared with HIV-noninfected persons. Given the rapid increase of HIV/AIDS cases in China, HHV-8 could become a severe public health issue in the future.

According to data from the Xinjiang Uygur Autonomous Region, minority groups were at higher risk for HHV-8 infection than the Han ethnic group. Although there was




Figure. Regional distribution of pooled human herpesvirus 8 (HHV-8) prevalence in A) the general population and B) immunocompromised patients, China.

evidence for considerable heterogeneity among the studies, the association between ethnicity and HHV-8 risk showed that minorities were at higher risk for HHV-8 infection when compared with the Han ethnic group. The reasons behind this association are not well elucidated. Because all of the comparisons of HHV-8 difference between minority groups and the Han ethnic group are from the Xinjiang Uygur Autonomous Region, epidemiologic confirmation of this observation would require data from other regions, which is currently unavailable. Our analysis showed a marginally significant association between blood contact and HHV-8 infection; heterogeneity among studies was substantial. These data indicate that blood-borne transmission could occur among the Chinese population, a finding that is consistent with previous reports from other countries (11).

This study has some limitations. The studies included in this meta-analysis were not evenly distributed throughout China because information was not available from all the regions. Also, all of the studies might have used different methods for HHV-8 detection because of the lack of a standard assay; prevalence estimates may have been underestimated.

In summary, this meta-analysis clearly shows that the distribution of HHV-8 seroprevalence varies in China. The available information is still too limited to fully understand HHV-8 prevalence and the risk factors associated with transmission. Further studies are urgently needed to explore the epidemiology of HHV-8 infection in different subpopulations in China.

This study was supported by the National Institutes of Health (PHS grant RO1 CA75903; Fogarty International Training Grant D43 TW01492 and T32 AI060547) and a National Center for Research Resources Centers of Biomedical Research Excellence grant (grant P30 RR031151) to C.W. The study was also funded, in part, by the Fundamental Research Funds for the Central Universities (10FX058) and Chinese National Natural Science Foundation (81072345).

Dr Tiejun Zhang is a postdoctorate research fellow at the University of Nebraska-Lincoln and a lecturer at the School of Public Health, Fudan University. His research focuses on HHV-8 and HIV epidemiology in China.

References

- Hoffman LJ, Bunker CH, Pellett PE, Trump DL, Patrick AL, Dollard SC, et al. Elevated seroprevalence of human herpesvirus 8 among men with prostate cancer. *J Infect Dis*. 2004;189:15–20. doi:10.1086/380568
- Laney AS, Peters JS, Manzi SM, Kingsley LA, Chang Y, Moore PS. Use of a multiantigen detection algorithm for diagnosis of Kaposi's sarcoma-associated herpesvirus infection. *J Clin Microbiol*. 2006;44:3734–41. doi:10.1128/JCM.00191-06
- Pellett PE, Wright DJ, Engels EA, Ablashi DV, Dollard SC, Forghani B, et al. Multicenter comparison of serologic assays and estimation of human herpesvirus 8 seroprevalence among US blood donors. *Transfusion*. 2003;43:1260–8. doi:10.1046/j.1537-2995.2003.00490.x
- Huang LM, Huang SY, Chen MY, Chao MF, Lu CY, Tien HF, et al. Geographical differences in human herpesvirus 8 seroepidemiology: a survey of 1,201 individuals in Asia. *J Med Virol*. 2000;60:290–3. doi:10.1002/(SICI)1096-9071(200003)60:3<290::AID-JMV7>3.0.CO;2-G
- Baeten JM, Chohan BH, Lavreys L, Rakwar JP, Ashley R, Richardson BA, et al. Correlates of human herpesvirus 8 seropositivity among heterosexual men in Kenya. *AIDS*. 2002;16:2073–8. doi:10.1097/00002030-200210180-00013
- Engels EA, Sinclair MD, Biggar RJ, Whitby D, Ebbesen P, Goedert JJ, et al. Latent class analysis of human herpesvirus 8 assay performance and infection prevalence in sub-saharan Africa and Malta. *Int J Cancer*. 2000;88:1003–8. doi:10.1002/1097-0215(20001215)88:6<1003::AID-IJC26>3.0.CO;2-9
- Rezza G, Tchchangmena OB, Andreoni M, Bugarini R, Toma L, Bakary DK, et al. Prevalence and risk factors for human herpesvirus 8 infection in northern Cameroon. *Sex Transm Dis*. 2000;27:159–64. doi:10.1097/00007435-200003000-00008
- Butler LM, Dorsey G, Hladik W, Rosenthal PJ, Brander C, Neilands TB, et al. Kaposi sarcoma-associated herpesvirus (KSHV) seroprevalence in population-based samples of African children: evidence for at least 2 patterns of KSHV transmission. *J Infect Dis*. 2009;200:430–8. doi:10.1086/600103
- Mbulaiteye S, Marshall V, Bagni RK, Wang CD, Mbisa G, Bakaki PM, et al. Molecular evidence for mother-to-child transmission of Kaposi sarcoma-associated herpesvirus in Uganda and K1 gene evolution within the host. *J Infect Dis*. 2006;193:1250–7. doi:10.1086/503052
- Dukers NH, Renwick N, Prins M, Geskus RB, Schulz TF, Weverling GJ, et al. Risk factors for human herpesvirus 8 seropositivity and seroconversion in a cohort of homosexual men. *Am J Epidemiol*. 2000;151:213–24.
- Hladik W, Dollard SC, Mermin J, Fowlkes AL, Downing R, Amin MM, et al. Transmission of human herpesvirus 8 by blood transfusion. *N Engl J Med*. 2006;355:1331–8. doi:10.1056/NEJMoa055009
- Ministry of Health of China. Joint United Nations Programme on HIV/AIDS, World Health Organization. The Estimation of HIV/AIDS in China in 2009. Beijing: Ministry of Health; 2010. p. 10.
- Dilnur P, Katano H, Wang ZH, Osakabe Y, Kudo M, Sata T, et al. Classic type of Kaposi's sarcoma and human herpesvirus 8 infection in Xinjiang, China. *Pathol Int*. 2001;51:845–52. doi:10.1046/j.1440-1827.2001.01293.x
- Parisi SG, Sarmati L, Pappagallo M, Mazzi R, Carolo G, Farchi F, et al. Prevalence trend and correlates of HHV-8 infection in HIV-infected patients. *J Acquir Immune Defic Syndr*. 2002;29:295–9.
- Chakraborty R, Rees G, Bourboulia D, Cross AM, Dixon JR, D'Agostino A, et al. Viral coinfections among African children infected with human immunodeficiency virus type 1. *Clin Infect Dis*. 2003;36:922–4. doi:10.1086/368207

Address for correspondence: Tiejun Zhang or Na He, Department of Epidemiology, School of Public Health, Fudan University, 138# Yi Xue Yuan RD, Shanghai 200032, China; email: tjzhangsh@gmail.com

The opinions expressed by authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institutions with which the authors are affiliated.

