

# Role of Birds in Dispersal of Etiologic Agents of Tick-borne Zoonoses, Spain, 2009

Ana M. Palomar, Paula Santibáñez,  
David Mazuelas, Lidia Roncero,  
Sonia Santibáñez, Aránzazu Portillo,  
and José A. Oteo

We amplified gene sequences from *Anaplasma phagocytophilum*, *Borrelia garinii*, *B. valaisiana*, *B. turdi*, *Rickettsia monacensis*, *R. helvetica*, *R. sibirica sibirica*, and *Rickettsia* spp. (including *Candidatus Rickettsia vini*) in ticks removed from birds in Spain. The findings support the role of passerine birds as possible dispersers of these tick-borne pathogens.

Hard ticks are a major vector of infectious diseases in industrialized countries. Several tick-borne bacterial diseases, such as Lyme disease, Mediterranean spotted fever, and tick-borne lymphadenopathy (also called *Dermacentor*-borne necrosis erythema and lymphadenopathy), are endemic to Spain. Furthermore, a few cases of human anaplasmosis and *Rickettsia monacensis* infection in humans have been diagnosed in Spain (1–3).

Birds are the preferred host for some tick species. As carriers of infected ticks, birds could be responsible for the spread of tick-borne bacteria that cause human anaplasmosis, Lyme disease, rickettsioses, and other diseases (4). Multiple studies support the conclusion or propose the hypothesis that birds play a role as reservoirs of *Anaplasma phagocytophilum*, *Borrelia burgdorferi*, and *Rickettsia* spp. (4–6). Because the Iberian Peninsula plays a major role in the migratory routes of birds, we aimed to determine the presence and prevalence of *A. phagocytophilum*, *B. burgdorferi* sensu lato, and *Rickettsia* spp. in ticks removed from birds captured in northern Spain.

## The Study

During April–October 2009, bird bandings were conducted in the protected area of Finca Ribavellosa in La Rioja, Spain (42°14'N, 2°54'W). Ticks were collected from birds and classified through taxonomic keys (7) and molecular methods (8). DNA was individually extracted by using 2 incubations of 20 minutes each with ammonium hydroxide (1 mL of 25% ammonia and 19 mL of Milli-Q water that had been autoclaved) at 100°C and 90°C.

DNA extracts were used as templates for PCRs targeting fragment genes for tick classification and for bacteria detection (Table 1). Two negative controls, 1 containing water instead of template DNA and the other with template DNA but without primers, and a positive control (a tick extract, *A. phagocytophilum*, *B. burgdorferi* sensu stricto, or *R. slovaca*) were included in all PCRs. Amplification products were sequenced, and nucleotide sequences were compared with those available in GenBank by using a BLAST search ([www.ncbi.nlm.nih.gov/blast/Blast.cgi](http://www.ncbi.nlm.nih.gov/blast/Blast.cgi)). Phylogenetic and molecular evolutionary analyses were conducted by using MEGA4 (16 in online Technical Appendix, [wwwnc.cdc.gov/EID/pdfs/11-1777-Techapp1.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-1777-Techapp1.pdf)).

A total of 222 ticks belonging to the species *Haemaphysalis punctata* (n = 1), *Ixodes frontalis* (n = 7), *I. arboricola* (n = 26), *I. ricinus* (n = 181), and other *Ixodes* spp. (n = 7) were collected from 97 passerine birds. Two nucleotide sequences for the 16S rRNA fragment gene of *I. arboricola* ticks were recorded (GenBank accession nos. JF791812 and JF791813) (Table 2).

*A. phagocytophilum* was detected only in 1 larva of an *I. ricinus* tick (0.5%). Twenty-nine (13.1%) samples tested positive for *B. burgdorferi* s.l. The most prevalent genospecies was *B. garinii* (n = 19), which was detected in *I. ricinus* (n = 16), *H. punctata* (n = 1), *I. frontalis* (n = 1), and *Ixodes* sp. (n = 1) ticks. *B. valaisiana* was amplified in 9 samples (8 *I. ricinus* and 1 *Ixodes* sp. ticks). *B. turdi* was found in 1 *I. frontalis* tick. *Rickettsia* infection was detected in 39 (17.6%) ticks. *R. monacensis* (n = 1), *R. helvetica* (n = 1), *R. sibirica sibirica* (n = 1), and *Rickettsia* spp. (n = 9) were detected in 12 *I. ricinus* ticks. Furthermore, according to *gltA*, *ompA*, and *ompB* sequence analysis, a possible new *Rickettsia* sp. was found in 25 *I. arboricola* ticks and 2 *I. ricinus* ticks. For these 27 samples, highest identities with *R. heilongjiangensis* (97.1%) and *R. japonica* (99.1%) were found for *ompA* (GenBank accession no. JF758828) and *ompB* (GenBank accession no. JF758826) nucleotide sequences, respectively, whereas *gltA* nucleotide sequences were identical to those from both *Rickettsia* spp. According to multilocus sequence typing (data not shown) and genetic criteria agreed on by experts, a *Candidatus* status could be assigned. We named it *Candidatus Rickettsia vini* (17 in online Technical Appendix) (Table 2). The phylogenetic

Author affiliations: Hospital San Pedro—Centro de Investigación Biomédica de La Rioja, Logroño, Spain (A.M. Palomar, P. Santibáñez, S. Santibáñez, A. Portillo, J.A. Oteo); and Environment Resources Inc., Logroño (D. Mazuelas, L. Roncero)

DOI: <http://dx.doi.org/10.3201/eid1807.111777>

Table 1. PCR primer pairs used in study of the role of birds in dispersal of etiologic agents of tick-borne zoonoses, Spain, 2009\*

Bacteria	Gene target	Primer name	Primer sequence, 5' → 3'	Amplified fragment, bp	Annealing temp., °C	Ref.
<i>Anaplasma</i> spp.	16S rRNA, nested	ge3a	CACATGCAAGTCGAACGGATTATTC	932	55	(9)
		ge10r	TTCCGTTAAGAAGGAT CTAATCTCC			
		ge9f	AACGGATTATCTTTATAGCTTGCT	546	55	(9)
	<i>msp</i>	ge2	GGCAGTATTAAGAAGCAGCTCCAGG			
		msp3F	CCAGCGTTTTAGCAAGATAAGAG	334	56	(10)
<i>Borrelia</i> spp.	<i>flaB</i> , nested†	Outer 1	AARGAATTGGCAGTTCAATC	497	52	(11)
		Outer 2	GCATTTTCWATTTTAGCAAGTGATG			
		Inner 1	ACATATTCAGATGCAGACAGAGGTTCTA	389	55	(11)
	5S-23S intergenic spacer, nested	Inner 2	GAAGGTGCTGTAGCAGGTGCTGGCTGT			
		23SC1	TAAGCTGACTAATACTAATTACCC	380	52	(12)
		23SN1	ACCATAGACTCTTATTACTTTGAC			
		5SCB	GAGAGTAGGTTATTGCCAGGG	226	55	(12)
23SN2	ACCATAGACTCTTATTACTTTGACCA					
<i>Rickettsia</i> spp.	<i>ompA</i> , seminested	Rr190.70p	ATGGCGAATATTTCTCCAAAA	631	46	(13,14)
		Rr190.701n	GTTCCGTTAATGGCAGCATCT			
	<i>ompB</i> , nested	Rr190.70p	ATGGCGAATATTTCTCCAAAA	532	48	(14)
		Rr190.602n	AGTGCAGCATTGCTCCCCCT			
		rompB OF	GTAACCGGAAGTAATCGTTTCGTAA	511	54	(15)
		rompB OR	GCTTTATAACCAGCTAAACCACC			
	<i>gltA</i> central region, nested	rompB SFG IF	GTTTAATACGTGCTGCTAACCAA	420	56	(15)
		rompB SFG/TG IR	GGTTTGGCCCATATACCATAAG			
		RpCS.877p	GGGGGCCTGCTCACGGCGG	381	48	(14)
		RpCS.1258n	ATTGCAAAAAGTACAGTGAACA			
RpCS.896p	GGCTAATGAAGCAGTGATAA	337	54	(15)		
RpCS.1233n	GCGACGGTATACCCATAGC					

\*Temp., temperature; ref., reference; *msp*, p44 major surface protein gene; *flaB*, flagellin gene; *ompB*, 120-kDa genus common antigen gene; *ompA*, 190-kDa protein antigen gene; *gltA*, citrate synthase gene.

†R = A/G; W = A/T.

Table 2. *Anaplasma phagocytophilum*, *Borrelia burgdorferi* s.l., and *Rickettsia* spp. detected in ticks removed from birds, Spain, 2009\*

Bacteria	Tick		Bird species (no. specimens)	Gene targets	
	Species	Stage			
<i>A. phagocytophilum</i>	<i>Ixodes ricinus</i>	1 L	<i>Turdus merula</i> (1)	<i>msp</i>	
<i>B. garinii</i>	<i>I. ricinus</i>	4 L, 2 N	<i>T. merula</i> (9)	<i>flaB</i> , 5–23S is	
		3 L, 4 N		<i>flaB</i> or 5–23S is	
		1 L	<i>Erithacus rubecula</i> (1)	<i>flaB</i>	
		1 L	<i>T. philomelos</i> (1)	<i>flaB</i> , 5–23S is	
	<i>I. frontalis</i>	1 L	<i>Troglodytes troglodytes</i> (1)	<i>flaB</i> , 5–23S is	
		1 F	<i>T. philomelos</i> (1)	<i>flaB</i> , 5–23S is	
	<i>Ixodes</i> spp.	1 L	<i>E. rubecula</i> (1)	5–23S is	
		<i>Haemaphysalis punctata</i>	1 L	<i>T. merula</i> (1)	<i>flaB</i> , 5–23S is
	<i>B. valaisiana</i>	<i>Ixodes</i> spp.	1 L	<i>T. merula</i> (1)	<i>flaB</i> , 5–23S is
			<i>I. ricinus</i>	1 L, 1 N	<i>T. merula</i> (3)
<i>I. ricinus</i>		2 L		<i>flaB</i>	
		1 L, 1 N	<i>T. philomelos</i> (2)	<i>flaB</i> , 5–23S is	
		1 L	<i>E. rubecula</i> (1)	<i>flaB</i> , 5–23S is	
1 L	<i>Garrulus glandarius</i> (1)	<i>flaB</i>			
<i>B. turdi</i>	<i>I. frontalis</i>	1 F	<i>T. merula</i> (1)	<i>flaB</i> , 5–23S is	
<i>R. monacensis</i>	<i>I. ricinus</i>	1 N	<i>Sylvia atricapilla</i> (1)	<i>ompA</i>	
<i>R. helvetica</i>	<i>I. ricinus</i>	1 N	<i>G. glandarius</i> (1)	<i>gltA</i>	
<i>R. sibirica sibirica</i>	<i>I. ricinus</i>	1 L	<i>S. atricapilla</i> (1)	<i>ompA</i>	
<i>Rickettsia</i> spp.†	<i>I. ricinus</i>	1 N, 1 L	<i>T. philomelos</i> (1)	<i>ompB</i> or <i>gltA</i>	
		4 L	<i>E. rubecula</i> (4)	<i>ompB</i> or <i>gltA</i>	
	<i>I. ricinus</i>	2 N	<i>T. merula</i> (2)	<i>gltA</i>	
		1 L	<i>Tr. troglodytes</i> (1)	<i>gltA</i>	
<i>Candidatus Rickettsia vini</i>	<i>I. arboricola</i>	20 N	<i>Cyanistes caeruleus</i> (1)	<i>ompA</i> , <i>ompB</i> , <i>gltA</i>	
		5 L	<i>Parus major</i> (1)	<i>ompA</i> , <i>ompB</i> , <i>gltA</i>	
	<i>I. ricinus</i>	2 L	<i>E. rubecula</i> (2)	<i>ompA</i> , <i>ompB</i> , <i>gltA</i>	

\*L, larva; *msp*, p44 major surface protein gene; N, nymph; *flaB*, flagellin gene; 5S-23S is, 5S-23S rRNA intergenic spacer; *ompB*, 120-kDa genus common antigen gene; *ompA*, 190-kDa protein antigen gene; *gltA*, citrate synthase gene.

†Same identity with >1 validly published *Rickettsia* species.

tree based on *ompA* gene shows the nearest relationships among *Rickettsia* spp. (Figure).

Two *I. ricinus* larvae showed co-infection with *B. garinii* and *Rickettsia* sp. One nymph was co-infected with *B. valaisiana* and *Rickettsia* sp.

## Conclusions

The presence of *Anaplasma*, *Borrelia*, and *Rickettsia* species in ticks removed from passerine birds corroborates the role of these vertebrates in the epidemiology and dispersion of tick-borne pathogens in Spain and in other zones of the planet. Some of the parasitized birds in our study, such as the European robin (*Erithacus rubecula*) or Eurasian blackcap (*Sylvia atricapilla*), are considered migratory or partial migratory birds. In addition, these species share an ecologic niche and ectoparasites

(horizontal transmission) with other migratory birds that cover long distances from Africa to the Eurasian region.

Except for *I. arboricola*, the tick species captured in this study previously had been found on birds in Spain (18 in online Technical Appendix). Nevertheless, *I. arboricola* ticks are commonly hosted by birds. The high prevalence of *I. ricinus* ticks was expected because it is the most frequent tick in this area, and the immature stages of this tick frequently parasitize birds.

*I. ricinus* ticks are the main vectors of *A. phagocytophilum* in Europe, and this microorganism has been detected on vegetation in the studied area (1). However, the low prevalence (0.5%) of *A. phagocytophilum* in the ticks in our study corroborates data from other studies (19,20 in online Technical Appendix). The presence of *A. phagocytophilum* in a larva in our study supports the role of birds as reservoirs of *A. phagocytophilum*.

The prevalence (13.1%) of *B. burgdorferi* in our samples is similar to prevalences reported in other studies in Europe in which *I. ricinus* is the main species of tick captured from birds (19 in online Technical Appendix). In Spain, *B. garinii*, *B. valaisiana*, and *B. afzelii* have been detected in ticks from birds (18 in online Technical Appendix). According to our data, the human pathogen *B. garinii* was the most prevalent species, as reported in birds from Europe (21 in online Technical Appendix). *B. turdi* was discovered in Asia. Although it has been recently detected in ticks from birds in Norway (22 in online Technical Appendix), its finding in Spain was unexpected.

Regarding *Rickettsia* species, *R. monacensis* and *R. helvetica* are among the human pathogens detected in our study. Both species have been identified in ticks from birds in Europe (19,20,23 in online Technical Appendix). On the contrary, *Candidatus Rickettsia vini*, a potential new *Rickettsia* species, also detected in our study, has not been related to human disease (17 in online Technical Appendix). Several genospecies closely related to *R. heilongjiangensis* and *R. japonica* have been identified in *Ixodes* spp. ticks removed from birds (23 in online Technical Appendix). *R. sibirica sibirica*, responsible for Siberian tick typhus in western People's Republic of China and in Siberia, was also amplified in an *I. ricinus* larva in this study.

Our data confirm the involvement of birds in the cycle of human tick-borne diseases. The findings confirm that birds can disperse vectors and microorganisms.

## Acknowledgments

We thank Agustín Estrada-Peña for classifying the *I. arboricola* ticks and Óscar Gutiérrez for providing tick samples.

This study was presented in part at the XIV Congreso SEIMC (Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica) (Spain) in May 2010, the GEPE Scientific




Figure. The phylogenetic position of *Candidatus Rickettsia vini* based on the *ompA* nucleotide sequences in a study of the role of birds in dispersal of etiologic agents of tick-borne zoonoses, Spain, 2009. The evolutionary history was inferred by using the neighbor-joining method. The optimal tree with the sum of branch length = 1.09961140 is shown. The percentage of replicate trees in which the associated taxa clustered in the bootstrap test (1,000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed by using the Kimura 2-parameter method and are in the units of the number of base substitutions per site. Codon positions included were 1st+2nd+3rd+Noncoding. All positions containing gaps and missing data were eliminated from the dataset. A total of 563 positions were in the final dataset. Phylogenetic analyses were conducted in MEGA4 (16 in online Technical Appendix, [wwwnc.cdc.gov/EID/pdfs/11-1777-Techapp1.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-1777-Techapp1.pdf)).

Meeting from the XV Congreso SEIMC (Spain) in June 2011, and the 6th International Meeting on Rickettsiae and Rickettsial Diseases (Greece) in June 2011.

Fundación Rioja Salud awarded a grant (FRS/PIF-01/10) to A.M.P. Financial support was provided in part by a grant from "Instituto de Salud Carlos III" (EMER 07/033), Ministerio de Ciencia e Innovación (Spain).

Dr Palomar has worked in the Center of Rickettsiosis and Arthropod-borne Diseases at the Infectious Diseases Area, Hospital San Pedro—Center for Biomedical Research of La Rioja since March 2009. Her research interests are the taxonomy of ticks and their associated pathogens.

## References

- Blanco JR, Oteo JA. Human granulocytic ehrlichiosis in Europe. *Clin Microbiol Infect*. 2002;8:763–72. <http://dx.doi.org/10.1046/j.1469-0691.2002.00557.x>
- Oteo JA, Backenson PB, del Mar Vitutia M, García Moncó JC, Rodríguez I, Escudero R, et al. Use of the C3H/He Lyme disease mouse model for the recovery of a Spanish isolate of *Borrelia garinii* from erythema migrans lesions. *Res Microbiol*. 1998;149:39–46. [http://dx.doi.org/10.1016/S0923-2508\(97\)83622-4](http://dx.doi.org/10.1016/S0923-2508(97)83622-4)
- Oteo JA, Portillo A. Tick-borne rickettsioses in Europe. *Ticks Tick Borne Dis*. 2012. In press.
- Hubálek Z. An annotated checklist of pathogenic microorganisms associated with migratory birds. *J Wildl Dis*. 2004;40:639–59.
- Hulínska D, Votyčka J, Plch J, Vlček E, Valesová M, Bojar M, et al. Molecular and microscopical evidence of *Ehrlichia* spp. and *Borrelia burgdorferi* sensu lato in patients, animals and ticks in the Czech Republic. *New Microbiol*. 2002;25:437–48.
- Humair PF. Birds and *Borrelia*. *Int J Med Microbiol*. 2002;291:70–4. [http://dx.doi.org/10.1016/S1438-4221\(02\)80015-7](http://dx.doi.org/10.1016/S1438-4221(02)80015-7)
- Manilla G. Fauna D'Italia Ixodida. Bologna (Italy): Calderini; 1998.
- Black WC, Piesman J. Phylogeny of hard and soft tick taxa (*Acari:Ixodida*) based on mitochondrial 16S rDNA sequences. *Proc Natl Acad Sci U S A*. 1994;91:10034–8. <http://dx.doi.org/10.1073/pnas.91.21.10034>
- Massung RF, Slater K, Owens JH, Nicholson WL, Mather TN, Solberg VB, et al. Nested PCR assay for detection of granulocytic ehrlichiae. *J Clin Microbiol*. 1998;36:1090–5.
- Zeidner NS, Burkot TR, Massung R. Transmission of the agent of human granulocytic ehrlichiosis by *Ixodes spinipalpis* ticks: evidence of an enzootic cycle of dual infection with *Borrelia burgdorferi* in northern Colorado. *J Infect Dis*. 2000;182:616–9. <http://dx.doi.org/10.1086/315715>
- Clark K, Hendricks A, Burge D. Molecular identification and analysis of *Borrelia burgdorferi* sensu lato in lizards in the southeastern United States. *Appl Environ Microbiol*. 2005;71:2616–25. <http://dx.doi.org/10.1128/AEM.71.5.2616-2625.2005>
- Rijpkema SG, Molkenboer MJ, Schouls LM, Jongejan F, Schellekens JF. Simultaneous detection and genotyping of three genomic groups of *Borrelia burgdorferi* sensu lato in Dutch *Ixodes ricinus* ticks by characterization of the amplified intergenic spacer region between 5S and 23S rRNA genes. *J Clin Microbiol*. 1995;33:3091–5.
- Roux V, Fournier PE, Raoult D. Differentiation of spotted fever group rickettsiae by sequencing and analysis of restriction fragment length polymorphism of PCR-amplified DNA of the gene encoding the protein *rOmpA*. *J Clin Microbiol*. 1996;34:2058–65.
- Regnery RL, Spruill CL, Plikaytis BD. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol*. 1991;173:1576–89.
- Choi YJ, Jang WJ, Kim JY, Lee SH, Park KH, Paik HS, et al. Spotted fever group and typhus group rickettsioses in humans, South Korea. *Emerg Infect Dis*. 2005;11:237–44. <http://dx.doi.org/10.3201/eid1102.040603>

Address for correspondence: José A. Oteo, Área de Enfermedades Infecciosas, Hospital San Pedro-CIBIR, C/Piqueras 98-7ª NE, 26006 Logroño (La Rioja), Spain; email: jaoteo@riojasalud.es

Get the content you want delivered to your inbox.



Table of Contents  
Podcasts  
Ahead of Print Articles  
Medscape CME™  
Specialized Content

Online subscription: [www.cdc.gov/ncidod/eid/subscrib.htm](http://www.cdc.gov/ncidod/eid/subscrib.htm)

# Role of Birds in Dispersal of Etiologic Agents of Tick-borne Zoonoses, Spain, 2009

## Technical Appendix

### References

16. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol.* 2007;24:1596–9. [PubMed](#)  
<http://dx.doi.org/10.1093/molbev/msm092>
17. Palomar AM, Portillo A, Santibáñez P, Santibáñez S, García-Álvarez L, Oteo JA. Genetic characterization of *Candidatus Rickettsia vini*. A new rickettsia amplified in ticks from La Rioja, Spain. *Ticks Tick Borne Dis.* 2012. In press.
18. Estrada-Peña A, Osácar JJ, Pichon B, Gray JS. Hosts and pathogen detection for immature stages of *Ixodes ricinus* (Acari: Ixodidae) in north-central Spain. *Exp Appl Acarol.* 2005;37:257–68. [PubMed](#) <http://dx.doi.org/10.1007/s10493-005-3271-6>
19. Franke J, Fritsch J, Tomaso H, Straube E, Dorn W, Hildebrandt A. Coexistence of pathogens in host-seeking and feeding ticks within a single natural habitat in central Germany. *Appl Environ Microbiol.* 2010;76:6829–36. [PubMed](#) <http://dx.doi.org/10.1128/AEM.01630-10>
20. Santos-Silva MM, Sousa R, Santos AS, Melo P, Encarnação V, Bacellar F. Ticks parasitizing wild birds in Portugal: detection of *Rickettsia aeschlimannii*, *R. helvetica* and *R. massiliae*. *Exp Appl Acarol.* 2006;39:331–8. [PubMed](#) <http://dx.doi.org/10.1007/s10493-006-9008-3>
21. Comstedt P, Bergström S, Olsen B, Garpmo U, Marjavaara L, Mejlom H, et al. Migratory passerine birds as reservoirs of Lyme borreliosis in Europe. *Emerg Infect Dis.* 2006;12:1087–95. [PubMed](#)  
<http://dx.doi.org/10.3201/eid1207.060127>
22. Hasle G, Bjune GA, Midthjell L, Røed KH, Leinaas HP. Transport of *Ixodes ricinus* infected with *Borrelia* species to Norway by northward-migrating passerine birds. *Ticks Tick Borne Dis.* 2011;2:37–43. [PubMed](#) <http://dx.doi.org/10.1016/j.ttbdis.2010.10.004>

23. Elfving K, Olsen B, Bergström S, Waldenström J, Lundkvist A, Sjöstedt A, et al. Dissemination of spotted fever rickettsia agents in Europe by migrating birds. PLoS ONE. 2010;5:e8572. [PubMed](https://pubmed.ncbi.nlm.nih.gov/19711111/)  
<http://dx.doi.org/10.1371/journal.pone.0008572>