

# Vertical Transmission of *Babesia microti*, United States

Julie T. Joseph, Kerry Purtill, Susan J. Wong, Jose Munoz, Allen Teal, Susan Madison-Antenucci, Harold W. Horowitz,<sup>1</sup> Maria E. Aguero-Rosenfeld,<sup>1</sup> Julie M. Moore, Carlos Abramowsky, and Gary P. Wormser

Babesiosis is usually acquired from a tick bite or through a blood transfusion. We report a case of babesiosis in an infant for whom vertical transmission was suggested by evidence of *Babesia* spp. antibodies in the heel-stick blood sample and confirmed by detection of *Babesia* spp. DNA in placenta tissue.

**B**abesiosis is an emerging infection in the United States, principally caused by *Babesia microti* (1). The most common route of infection is the bite of an *Ixodes scapularis* tick; transmission can also occur by transfusion of infected blood products, and vertical transmission in animals has been documented (2,3) and is a potential route of transmission for humans. We report a case of babesiosis in an infant for whom vertical transmission was suggested by *Babesia* spp. antibodies in a heel spot blood sample and confirmed by detection of *Babesia* DNA in placenta tissue.

## The Case-Patient

A 6-week-old girl from Yorktown Heights, New York, was admitted to the hospital on September 16, 2002, with a 2-day history of fever, irritability, and decreased oral intake. The mother was asymptomatic during and after her pregnancy. The infant was delivered vaginally and full term at 3,430 g without complications. The infant's mother had visited parks in Westchester and Dutchess Counties in New York during the pregnancy but was unaware of any tick bites. The infant had no known tick exposure, and neither mother nor infant had a history of blood transfusion.

Author affiliations: New York Medical College, Valhalla, New York, USA (J.T. Joseph, K. Purtill, J. Munoz, H.W. Horowitz, M.E. Aguero-Rosenfeld, G.P. Wormser); New York State Department of Health, Albany, New York, USA (S.J. Wong, A. Teal, S. Madison-Antenucci); University of Georgia, Athens, Georgia, USA (J.M. Moore); and Emory University School of Medicine, Atlanta, Georgia, USA (C. Abramowsky)

DOI: <http://dx.doi.org/10.3201/eid1808.110988>




Figure. Peripheral blood smear of 6-week-old infant with suspected congenital babesiosis. Thin arrows indicate *Babesia* spp. parasites; thick arrow shows the classic tetrad formation or Maltese cross.

During examination, the infant was alert but irritable and pale. Axillary temperature was initially 36.8°C but increased to 38.1°C on the same day. Her conjunctivae were icteric, she had a palpable spleen tip, and her liver was palpable 3 cm below the costal margin. Initial laboratory findings included hemoglobin 7.1 g/dL, platelet count  $100 \times 10^3/\mu\text{L}$ , and leukocyte count  $19.7 \times 10^3$  cells/ $\mu\text{L}$  with a differential of 4% segmented neutrophils, 80% lymphocytes, and 16% monocytes. Reticulocyte count was 5.5%. Total bilirubin concentration was 2 mg/dL with a direct fraction of 0.4 mg/dL; aspartate aminotransferase level was 66 U/L, alanine aminotransferase level was 50 U/L, and alkaline phosphatase level was 339 U/L. Cultures of blood, urine, and cerebrospinal fluid samples yielded negative results. Lyme disease serologic test result was negative.

Routine examination of a peripheral blood smear showed *B. microti* in 4% of erythrocytes (Figure); a blood sample from the infant was positive by PCR for *B. microti* DNA. Total *B. microti* antibody titer was >256 by indirect immunofluorescence assay, with a polyvalent secondary antibody (anti-IgG+IgA+IgM) (4) that was presumed to be principally IgG because test results for IgM were negative (online Technical Appendix, [wwwnc.cdc.gov/EID/pdfs/11-0988-Techapp.pdf](http://wwwnc.cdc.gov/EID/pdfs/11-0988-Techapp.pdf)). The heel-stick blood sample obtained on the infant's third day of life as part of newborn screening was tested and found to be negative for *B. microti* by PCR (5) and for IgM but total antibody positive (>128) (online Technical Appendix).

Examination of the placenta showed focal basal decidual inflammation, mild chorangiosis, and villus dysmaturity. *Babesia* spp. piroplasms were not detected in

<sup>1</sup>Current affiliation: New York University School of Medicine, New York, New York, USA.

maternal or fetal blood by histologic examination of hematoxylin and eosin–stained sections of formalin-fixed, paraffin-embedded tissue of the placenta disk, amnion/chorion, and umbilical cord. *Babesia* DNA was detected by real-time PCR testing of paraffin-embedded placenta tissue (online Technical Appendix) (6). Cycle threshold values were relatively high (37.1–38.2), indicating that the amount of parasite DNA in the sample was close to the limit of detection; results were reproducible on duplicate

testing of DNA samples extracted from separate paraffin blocks. The real-time PCR product was of the correct size, and the melting curve demonstrated melting temperatures within 1°C from the placenta, the positive control, and a positive sample from an unrelated patient, confirming that the correct product was amplified. At time of the illness in the infant, the mother was negative for *Babesia* spp. according to PCR and smear but positive for total antibodies (>256).

Table. Comparison of selected clinical and laboratory data from reported cases of congenital babesiosis in 5 infants\*

Clinical data	Reference				This study
	(7)	(8)	(9)	(10)	
Year of diagnosis/ location	Not given/Long Island, New York	Not given/Long Island, New York	Not given/New Jersey	Not given/Long Island, New York	2002/Westchester County, New York
Infant age at time of symptom onset, d	30	32	19	27	41
Clinical findings	Fever, irritability, pallor, hepatosplenomegaly	Fever, lethargy, poor feeding, pallor, scleral icterus, hepatomegaly	Fever, poor feeding, gagging, irritability, pallor, scleral icterus, hepato- splenomegaly	Fever, pallor	Fever, decreased oral intake, irritability, scleral icterus, pallor, hepatosplenomegaly
Initial babesia parasitemia level, %	5	4.4	15	2	4
Hospitalization, d	6	5	8	NA	5
Maternal tick bite	1 wk before delivery	7 wk before delivery	4 wk before delivery	None known	None known
<i>Babesia</i> spp. serologic and PCR results for infant	30 d after birth: IgM+/IgG+ (128/128) by IFA; 32 d after birth: IgM+/IgG+ (256/512) by IFA; PCR ND	At illness onset: IgG IFA 160; IgM/IgG immunoblot +; PCR ND	At illness onset: IgM+/IgG+ (40/256) by IFA; PCR ND	NA	Newborn screening (heel stick): IgM– (<16); total antibody + (>128) by IFA; PCR–; 6 wks after birth: IgM– (<16); total antibody + (>256) by IFA; PCR+; Birth: placenta PCR+;
<i>Babesia</i> spp. evaluation results for mother	30 d after birth: IgM+/IgG+ (2,048/1,024); 32 d after birth: IgM+/ IgG+ (4,096/1,024); peripheral smear – at time of delivery and at 30 and 32 d after birth	7 wk before birth: IgG IFA <40; IgM/IgG immunoblot –; 2 mo after birth: IgG IFA 640; IgM/IgG immunoblot +; peripheral smear – at delivery and at infant illness onset	At infant illness onset: IgM+/IgG+ (80/>1,024) by IFA; peripheral smear negative at time of infant illness onset	At infant illness onset: PCR+	6 wk after birth: IgM ND; total antibody + (>256) by IFA; PCR–; peripheral smear –
HGB, g/dL	9.3	10.8	8.8	NA; HCT 24.3%	7.1
Platelets, x 10 <sup>3</sup> /μL	38	87	34	101	100
Leukocytes/PMN leukocytes, cells/μL	6,500/1,170	NA	9,000/1,890	NA	19,700/788
LDH, U/L	894	NA	2535	NA	NA
Bilirubin indirect, mg/dL	3.6	9.7	5.9	NA	1.6
AST, U/L	90	NA	53	NA	66
ALT, U/L	90	NA	18	NA	50
Treatment	CLI and quinine for 10 d	CLI and quinine with AZT added on day 3; on day 5 changed to AZT plus quinine for additional 7 d	AZT and ATO for 10 d	AZT and ATO, duration not given	AZT and ATO for 9 d
Follow-up	Well at 6 mo posttreatment	Improved at 2 wk	Lost to follow-up	NA	22 mo
Blood transfusion for anemia	Yes, for HCT of 18%	Yes, for HGB of 7.3 g/dL	Yes, for HGB of 7.0 g/dL	Yes, for HCT of 17.3%	Yes, for HGB of 5.2 g/dL with HCT of 15.8%

\*No mothers became ill. NA, not available; +, positive; IFA, indirect immunofluorescence assay; ND, not done; –, negative; HGB, hemoglobin; HCT, hematocrit; PMN, polymorphonuclear; LDH, lactate dehydrogenase level; AST, aspartate aminotransferase; ALT, alanine aminotransferase; CLI, clindamycin; AZT, azithromycin; ATO, atovaquone.

The infant was treated with a 9-day course of azithromycin plus atovaquone. A blood transfusion was administered when her hemoglobin concentration fell to 5.2 g/dL. The infant became afebrile by 72 hours and was discharged after a 5-day hospitalization. Repeat blood smears revealed a parasite load of 0.3% at discharge. On final evaluation at 22 months of age, physical examination revealed no abnormalities; hemoglobin level was 11.7 g/dL, *Babesia* PCR was negative, and total *Babesia* antibody level was positive at 128.

### Conclusions

Congenital babesiosis has been rarely reported (Table 7–10). This case provided convincing evidence for congenital babesiosis because of prepartum infection involving the placenta in the mother. On the basis of experience with congenital malaria, we assume that *Babesia* spp. parasites cross the placenta during pregnancy or at the time of delivery (11,12). In congenital malaria, increasing evidence suggests that the malaria parasites are most often acquired antenatally by transplacental transmission of infected erythrocytes (12).

Reported cases of congenital babesiosis share many similarities, including asymptomatic maternal infection and development of fever, hemolytic anemia, and thrombocytopenia in the infant detected between 19 and 41 days after birth. All of the infants responded to antimicrobial drug therapy; 3 were treated with azithromycin plus atovaquone (9,10), the preferred treatment regimen for mild babesiosis (1). All infants required a blood transfusion because of severe anemia. The clinical signs and symptoms for these cases of congenital babesiosis are similar to those of congenital malaria in non-disease endemic areas (11,13).

We found *Babesia* spp. antibodies on day 3 of life by analyzing the patient's heel-stick blood sample, which likely represented maternal transfer of IgG. Passive transfer of maternal antibodies is regarded as a protective factor against congenital malaria, and some newborns with malaria who are parasitemic at birth spontaneously clear the infection without ever becoming ill (11,14). The temporary presence of maternal IgG in infants has been suggested as an explanation for the typical 3–6 week incubation period of congenital malaria in non-disease endemic areas (14).

The real-time PCR used to find *B. microti* DNA in placenta tissue is  $\approx 20\times$  more sensitive than microscopic examination of Giemsa-stained blood smears (6). Assuming a blood sample with a parasitemia equivalent to that detected in the placental tissue, a blood smear would contain  $\leq 10$  infected cells per slide. Given the low level of *Babesia* DNA in the placenta tissue, it is not surprising that histologic examination did not reveal piroplasms. Nonetheless, limited evidence of placental abnormalities suggests a pathologic process.

In summary, babesiosis is an emerging infectious disease (15) that can rarely cause congenital infection. This diagnosis should be considered in the differential diagnosis of fever and hemolytic anemia in infants from disease-endemic areas.

### Acknowledgments

The authors thank Steven Smith, Jennifer Calder, Lisa Giaratano, Lenise Banwarie, Ewa Bajor-Dattilo, and Karen Kulas for their assistance.

Dr Joseph is an assistant professor of medicine in the Division of Infectious Diseases at New York Medical College. Her research interests are tick-borne illnesses, particularly babesiosis.


### References

- Vannier E, Gewurz BE, Krause PJ. Human babesiosis. *Infect Dis Clin North Am.* 2008;22:469–88. <http://dx.doi.org/10.1016/j.idc.2008.03.010>
- de Vos AJ, Imes GD, Cullen JSC. Cerebral babesiosis in a new-born calf. *Onderstepoort J Vet Res.* 1976;43:75–8.
- Fukumoto S, Suzuki H, Igarashi I, Xuan X. Fatal experimental transplacental *Babesia gibsoni* infections in dogs. *Int J Parasitol.* 2005;35:1031–5. <http://dx.doi.org/10.1016/j.ijpara.2005.03.018>
- Chisholm ES, Ruebush TK II, Sulzer AJ, Healy GR. *Babesia microti* infection in man: evaluation of an indirect immunofluorescent antibody test. *Am J Trop Med Hyg.* 1978;27:14–9.
- Persing DH, Mathiesen D, Marshall WF, Telford SR, Spielman A, Thomfod JW, et al. Detection of *Babesia microti* by polymerase chain reaction. *J Clin Microbiol.* 1992;30:2097–103.
- Teal AE, Habura A, Ennis J, Keithly J, Madison-Antenucci S. A new real-time PCR assay for improved detection of the parasite *Babesia microti*. *J Clin Microbiol.* 2012;50:903–8. <http://dx.doi.org/10.1128/JCM.05848-11>
- Esernio-Jenssen D, Scimeca PG, Benach JL, Tenenbaum MJ. Transplacental/perinatal babesiosis. *J Pediatr.* 1987;110:570–2. [http://dx.doi.org/10.1016/S0022-3476\(87\)80552-8](http://dx.doi.org/10.1016/S0022-3476(87)80552-8)
- New DL, Quinn J, Qureshi MZ, Sigler S. Vertically transmitted babesiosis. *J Pediatr.* 1997;131:163–4. [http://dx.doi.org/10.1016/S0022-3476\(97\)70143-4](http://dx.doi.org/10.1016/S0022-3476(97)70143-4)
- Sethi S, Alcid D, Kesarwala H, Tolan RW Jr. Probable congenital babesiosis in infant, New Jersey, USA. *Emerg Infect Dis.* 2009;15:788–91. <http://dx.doi.org/10.3201/eid1505.070808>
- Aderinboye O, Syed S. Congenital babesiosis in a four-week old female infant. *Pediatr Infect Dis J.* 2010;29:188. <http://dx.doi.org/10.1097/INF.0b013e3181c3c971>
- Vottier G, Arsac M, Farnoux C, Mariani-Kurddjian P, Baud O, Aujard Y. Congenital malaria in neonates: two case reports and review of the literature. *Acta Paediatr.* 2008;97:505–8. <http://dx.doi.org/10.1111/j.1651-2227.2008.00690.x>
- Malhotra I, Mungai P, Muchiri E, Kwiek JJ, Meshnick SR, King CL. Umbilical cord-blood infections with *Plasmodium falciparum* malaria are acquired antenatally in Kenya. *J Infect Dis.* 2006;194:176–83. <http://dx.doi.org/10.1086/505150>
- Lesko CR, Arguin PM, Newman RD. Congenital malaria in the United States. A review of cases from 1966 to 2005. *Arch Pediatr Adolesc Med.* 2007;161:1062–7. <http://dx.doi.org/10.1001/archpedi.161.11.1062>

14. Hagmann S, Khanna K, Niazi M, Purswani M, Robins EB. Congenital malaria, an important differential diagnosis to consider when evaluating febrile infants of immigrant mothers. *Pediatr Emerg Care*. 2007;23:326–9. <http://dx.doi.org/10.1097/01.pcc.0000270164.78238.7d>
15. Joseph JT, Roy SS, Shams N, Visintainer P, Nadelman RB, Hosur S, et al. Babesiosis in Lower Hudson Valley, New York, USA. *Emerg Infect Dis*. 2011;17:843–7.

Address for correspondence: Julie T. Joseph, New York Medical College, Division of Infectious Diseases, Munger Pavilion Room 245, Valhalla, NY 10595, USA; email: [julie\\_joseph@nyc.edu](mailto:julie_joseph@nyc.edu)

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or by the US Department of Health and Human Services.



# Vertical Transmission of *Babesia microti*, United States

## Technical Appendix

### Antibody testing of the infant's blood spot

Approximately one-fourth of the blood spot (three 1/8 inch diameter punches) was eluted in 140  $\mu$ L of 10 g/L goat serum (Sigma) in PBS overnight at 4°C. The eluate was filtered through a 0.45- $\mu$ m syringe filter and then tested in a polyvalent (IgM, IgA, IgG) indirect immunofluorescence assay (IFA) for *Babesia* spp. antibodies. Controls were made by spotting both *Babesia* antibody negative whole blood and the same blood spiked with a positive control serum. The spots were allowed to dry overnight and eluted as above. The IFA used *Babesia* spp.-infected golden hamster erythrocytes fixed to glass slides. The conjugate was polyvalent FITC-conjugated anti-human antibody (Kirkegaard & Perry Laboratories, Gaithersburg, MD) in 0.1% Evans blue dye (1). For detection of IgM antibody specifically, the eluates were treated with GullSorb (Meridian Diagnostics, now Meridian Bioscience, Cincinnati, OH) to remove IgG and then retested using the same IFA procedure with an IgM-specific FITC-conjugate (KPL). The controls achieved the expected results.

### DNA extraction and conventional PCR

DNA was extracted from EDTA-preserved whole blood, 3 mm diameter punches of the dried blood spot, and from freshly cut 10- $\mu$ m sections of paraffin-embedded tissue, using protocols for blood, dried blood spots, and FFPE Tissue respectively (QIAGEN). Conventional PCR was performed on the extracts from the blood spot and whole blood specimen essentially as previously described targeting the 18S rDNA of *B. microti* (2), except the primer sets were used independently to produce two amplification products. Amplification of the human  $\beta$ -globin gene was performed to test for inhibition. The positive control was a plasmid containing the rDNA target, spiked into a negative blood sample and extracted in parallel with the patient specimen.

Cycling conditions were: 95°C 10 min, 35 cycles (94°C 1 min, 58°C 1 min, 72°C 1 min), with a final extension at 72°C for 4 min.

## Real-Time PCR

DNA extracted from the paraffin embedded tissue was assayed for the presence of *B. microti* using a recently developed real-time PCR assay targeting the 18S rRNA gene (6). Extracted DNA (15 µl) was amplified using FastStart DNA Master mix without ROX (Roche) in a 25 µl reaction mixture with 5.5 mM MgCl<sub>2</sub>, 500 nM primers (forward-CAGGGAGGTAGTGACAAGAAATAACA and reverse-GGTTTAGATTCCCATTCCAAT, and 250 nM probe (6FAM-TACAGGGCTTAAAGTCT-MGBNFQ). A 2-step PCR cycling protocol (95°C 10 min, and 45 cycles of 95°C 15 s, 60°C 1 min) was used to amplify a 72-bp product. An internal control was used to evaluate the efficiency of extraction and test for inhibition of the sample, as described elsewhere (3).

## References

1. Chisholm ES, Ruebush TK II, Sulzer AJ, Healy GR. *Babesia microti* infection in man: evaluation of an indirect immunofluorescent antibody test. Am J Trop Med Hyg. 1978;27:14–9. [PubMed](#)
2. Persing DH, Mathiesen D, Marshall WF, Telford SR, Spielman A, Thomfod JW, et al. Detection of *Babesia microti* by polymerase chain reaction. J Clin Microbiol. 1992;30:2097–103. [PubMed](#)
3. Teal AE, Habura A, Ennis J, Keithly J, Madison-Antenucci S. A new real-time PCR assay for improved detection of the parasite *Babesia microti*. J Clin Microbiol. 2012;50:903–8. [PubMed](#)  
<http://dx.doi.org/10.1128/JCM.05848-11>