

are needed to determine the role of nasal carriage in *B. holmesii* bacteremia. That no *B. holmesii* infections occurred after rituximab was stopped suggests that rituximab played a role in the recurrent infections. In cases of recurrent infection or bacteremia, nasal carriage should be assessed, and the interruption of rituximab should be considered by physicians.

### Acknowledgments

We thank Alain Le Coustumier for his advice concerning antimicrobial treatment and Institut Pasteur Fondation, Institut National de Veille Sanitaire, and Le Centre National de la Recherche Scientifique for financial support.

**Liem Binh Luong Nguyen,  
Loïc Epelboin, Jean Gabarre,  
Marylin Lecso, Sophie Guillot,  
François Bricaire, Eric Caumes,  
and Nicole Guiso**

Author affiliations: Groupe Hospitalier Pitié-Salpêtrière, Paris, France (L.B. Luong Nguyen, L. Epelboin, J. Gabarre, M. Lecso, F. Bricaire, E. Caumes); Université Paris, Paris (L. Epelboin, F. Bricaire, E. Caumes); and Institut Pasteur, Paris (S. Guillot, N. Guiso)

DOI: <http://dx.doi.org/10.3201/eid1910.130345>

### References

- Njamkepo E, Bonacorsi S, Debruyne M, Gibaud SA, Guillot S, Guiso N. Significant finding of *Bordetella holmesii* DNA in nasopharyngeal samples from French patients with suspected pertussis. *J Clin Microbiol*. 2011;49:4347–8. <http://dx.doi.org/10.1128/JCM.01272-11>
- Weyant RS, Hollis DG, Weaver RE, Amin MFM, Steigerwalt AG, O'Connor SP, et al. *Bordetella holmesii* sp. nov., a new gram-negative species associated with septicemia. *J Clin Microbiol*. 1995; 33:1–7.
- Rodgers L, Martin SW, Cohn A, Budd J, Marcon M, Terranella A, et al. Epidemiologic and laboratory features of a large outbreak of pertussis-like illnesses associated with cocirculating *Bordetella holmesii* and *Bordetella pertussis*—Ohio, 2010–2011. *Clin Infect Dis*. 2013;56:322–31. <http://dx.doi.org/10.1093/cid/cis888>
- Mooi FR, Bruisten S, Linde I, Reubsæet F, Heuvelman K, van der Lee S, et al. Characterization of *Bordetella holmesii* isolates from patients with pertussis-like illness in the Netherlands. *FEMS Immunol Med Microbiol*. 2012;64:289–91. <http://dx.doi.org/10.1111/j.1574-695X.2011.00911.x>
- Kamiya H, Otsuka N, Ando Y, Odaira F, Yoshino S, Kawano K, et al. Transmission of *Bordetella holmesii* during pertussis outbreak, Japan. *Emerg Infect Dis*. 2012;18:1166–9. <http://dx.doi.org/10.3201/eid1807.120130>
- Yih WK, Silva EA, Ida J, Harrington N, Lett SM, George H. *Bordetella holmesii*-like organisms isolated from Massachusetts patients with pertussis-like symptoms. *Emerg Infect Dis*. 1999;5:441–3. <http://dx.doi.org/10.3201/eid0503.990317>
- Chambaraud T, Dickson Z, Ensergueix G, Barraud O, Essig M, Lacour C, et al. *Bordetella holmesii* bacteremia in a renal transplant recipient: emergence of a new pathogen. *Transpl Infect Dis*. 2012;14:E134–6. <http://dx.doi.org/10.1111/tid.12009>
- Vidal L, Gafter-Gvili A, Salles G, Dreyling MH, Ghilmini M, Hsu Schmitz SF, et al. Rituximab maintenance for the treatment of patients with follicular lymphoma: an updated systematic review and meta-analysis of randomized trials. *J Natl Cancer Inst*. 2011;103:1799–806. <http://dx.doi.org/10.1093/jnci/djr418>
- Fry NK, Duncan J, Pike R, Harrison TG. Emergence of *Bordetella holmesii* infections in the United Kingdom 2010. In: Abstracts of the 9th International *Bordetella* Symposium, Baltimore, Maryland, USA; 2010 Sep 30–Oct 3. Abstract 98.
- Abouanaser SF, Srigley JA, Nguyen T, Dale SE, Johnstone J, Wilcox L, et al. *Bordetella holmesii*, an emerging cause of septic arthritis. *J Clin Microbiol*. 2013;51:1313–5. <http://dx.doi.org/10.1128/JCM.06437-11>

Address for correspondence: Nicole Guiso, Institut Pasteur, Molecular Prevention and Therapy of Human Diseases Unit, 25 Rue du Dr Roux, 75724 Paris Cedex 15, France; email: [nicole.guiso@pasteur.fr](mailto:nicole.guiso@pasteur.fr)

### Another Dimension

Thoughtful essays, short stories, or poems on philosophical issues related to science, medical practice, and human health. Topics may include science and the human condition, the unanticipated side of epidemic investigations, or how people perceive and cope with infection and illness. This section is intended to evoke compassion for human suffering and to expand the science reader's literary scope. Manuscripts are selected for publication as much for their content (the experiences they describe) as for their literary merit.

## ***Rickettsia africae* in *Amblyomma* *variegatum* Ticks, Uganda and Nigeria**

**To the Editor:** *Rickettsia africae* is the most widespread spotted fever group (SFG) rickettsia in sub-Saharan Africa, where it causes African tick-bite fever (1), an acute, influenza-like syndrome. The number of cases in tourists returning from safari in sub-Saharan Africa is increasing (1). In western, central, and eastern sub-Saharan Africa, *R. africae* is carried by *Amblyomma variegatum* (Fabricius, 1794) ticks (2); usually associated with cattle, this 3-host tick also can feed on a variety of hosts, including humans (2). *R. africae* has not been reported in Uganda and rarely reported in Nigeria (3,4). Our objective was to determine the potential risk for human infection by screening for rickettsial DNA in *A. variegatum* ticks from cattle in Uganda and Nigeria.

In February 2010, ticks were collected from zebu cattle (*Bos indicus*) from 8 villages in the districts of Kaberamaido (Adektar [1°81'N–33°22' E], Awimon [1°66'N–33°04' E], Kalo-bo [1°88' N–33°25' E], Odidip [1°90' N–33°30' E], Odikara [1°91' N–33°30' E], and Olilimo [1°75' N–33°38' E], and Dokolo (Alela [2°09' N–33°16' E], and Angeta [1°87' N–33°10' E]) in Uganda and, in June 2010, in 3 villages (Mangar [9°14' N–8°93' E], Ruff [9°43' N–9°10' E], and Tambes [9°38' N–9°38' E]) in the Plateau State in Nigeria (Figure). This convenience sample was obtained as part of other ongoing research projects in both countries. Ticks were preserved in 70% ethanol and identified morphologically to the species level by using taxonomic keys (5). Because the anatomic features do not enable an objective assessment of the feeding status of adult male ticks, engorgement level was determined only in female tick specimens and nymphs.




Figure. Location of areas studied for *Rickettsia africae* in *Amblyomma variegatum* ticks in Nigeria (A) and Uganda (B), 2010.

After tick identification, DNA was extracted from ticks by using QIAmp DNeasy kits (QIAGEN, Hilden, Germany). Two PCR targets were assessed within each sample; the primer pair Rp.CS.877p and Rp.CS.1258n was selective for a 396-bp fragment of a highly conserved gene encoding the citrate synthase (*gltA*) shared by all *Rickettsia* spp. (6); the Rr190–70p and Rr190–701n primer pair amplified a 629–632-bp fragment of the gene encoding the 190-kD antigenic outer membrane protein A (*ompA*), common to all SFG rickettsiae (6,7). DNA extracted from 2 *A. variegatum* tick cell lines (AVL/CTVM13 and AVL/CTVM17), previously amplified and sequenced by using primers for *Rickettsia* 16S rRNA, *ompB*, and *sca4* genes revealing >98% similarity with *R. africae* (8), was used as a positive control. Negative controls consisted of DNA from 2 male and female laboratory-reared *Rhipicephalus appendiculatus* ticks and distilled water. DNA of positive samples was recovered, and confirmation of amplicon authenticity was obtained through sequence analysis by using

nucleotide BLAST ([www.ncbi.nlm.nih.gov/BLAST](http://www.ncbi.nlm.nih.gov/BLAST)).

A total of 39 ticks were collected in Uganda (32 adult males, 5 females, and 2 nymphs), and 141 were collected in Nigeria (80 males, 59 females, and 2 nymphs); all were identified as *A. variegatum* (online Technical Appendix Table, [wwwnc.cdc.gov/EID/articlepdfs/19/10/13-0389-Techapp1.pdf](http://wwwnc.cdc.gov/EID/articlepdfs/19/10/13-0389-Techapp1.pdf)). SFG rickettsiae DNA was amplified in 26 (67%) of 39 ticks from Uganda and 88 (62%) of 141 ticks from Nigeria by using the *ompA* gene primers; amplicons of the *gltA* genes were obtained in 16 (41%) of 39 ticks and 84 (60%) of 141 ticks, respectively (online Technical Appendix Table). Overall, 81 (45%) of 180 ticks were positive by *gltA* and *ompA* PCRs (online Technical Appendix Table). DNA sequences of the 22 *gltA* and *ompA* products from Uganda and the 22 from Nigeria showed 100% similarity with published sequences of *R. africae* (GenBank accession nos. U59733 and RAU43790, respectively). For both countries, ticks positive for *Rickettsia* spp. and SFG rickettsiae DNA were

male and female specimens (online Technical Appendix Table). Among females, both unengorged and engorged specimens contained DNA from rickettsiae and SFG rickettsiae (online Technical Appendix Table).

These findings represent a novelty for Uganda. With reference to Nigeria, our results contrast with the prevalence of 8% recorded in a similarly sized sample ( $n = 153$ ) of *A. variegatum* ticks collected from cattle in the same part of the country (3); this discrepancy might be the result of previous targeting of the rickettsial 16S rDNA gene. In the study reported here, the SFG-specific *ompA* PCR proved to be more sensitive than *gltA* for detecting rickettsiae DNA, as has also been reported in previous work (9). Although finding *R. africae* DNA in engorged female and nymphal tick specimens might be attributable to prolonged rickettsemia in cattle (10), the presence of *R. africae* in distinctly unengorged female ticks indicates the potential for *A. variegatum* ticks to act as a reservoir of this SFG rickettsia (2).

This study extends the known geographic range of *R. africae* in *A. variegatum* ticks in sub-Saharan Africa. The number of potentially infective ticks recorded in Uganda and Nigeria suggests that persons in rural areas of northern Uganda and central Nigeria might be at risk for African tick-bite fever. Awareness of this rickettsiosis should be raised, particularly among persons who handle cattle (e.g., herders and paraveterinary and veterinary personnel). Physicians in these areas as well as those who care for returning travelers, should consider African tick-bite fever in their differential diagnosis for patients with malaria and influenza-like illnesses.

#### Acknowledgments

We thank Abraham Goni Dogo for helping with tick collections in Nigeria, Lesley Bell-Sakyi and Pilar Alberdi for providing positive controls, Tim Connelley for providing laboratory-reared ticks to be used as negative controls, Cristina Socolovschi for her valuable suggestions on the molecular proceedings, and Albert Mugenyi for his indispensable assistance with map design.

The research leading to these results received funding from the UK Department for International Development under the umbrella of the Stamp Out Sleeping Sickness Programme, the UK Biotechnology and Biological Sciences Research Council under the Combating Infectious Diseases in Livestock for International Development scheme, and the European Union's Seventh Framework Program (FP7/2007-2013) under grant agreement no. 221948, Integrated Control of Neglected Zoonoses.

**Vincenzo Lorusso,  
Karolina Anna Gruszka,  
Ayodele Majekodunmi,  
Augustine Igweh,  
Susan C. Welburn,  
and Kim Picozzi**

Author affiliations: University of Edinburgh, Edinburgh, Scotland, UK (V. Lorusso, K.A. Gruszka, A. Majekodunmi, S. Welburn, K. Picozzi); and Nigerian Institute for Trypanosomiasis Research, Jos, Nigeria (A. Igweh)

DOI: <http://dx.doi.org/10.3201/eid1910.130389>

#### References

1. Parola P, Paddock CD, Raoult D. Tick-borne rickettsioses around the world: emerging diseases challenging old concepts. *Clin Microbiol Rev*. 2005;18:719–56. <http://dx.doi.org/10.1128/CMR.18.4.719-756.2005>
2. Socolovschi C, Huynh TP, Davoust B, Gomez J, Raoult D, Parola P. Transovarial and trans-stadial transmission of *Rickettsia africae* in *Amblyomma variegatum* ticks. *Clin Microbiol Infect*. 2009;15(Suppl. 2):317–8. <http://dx.doi.org/10.1111/j.1469-0691.2008.02278.x>
3. Ogo NI, de Mera IG, Galindo RC, Okubanjo OO, Inuwa HM, Agbede RI, et al. Molecular identification of tick-borne pathogens in Nigerian ticks. *Vet Parasitol*. 2012;187:572–7. <http://dx.doi.org/10.1016/j.vetpar.2012.01.029>
4. Reye AL, Arinola OG, Hübschen JM, Muller CP. Pathogen prevalence in ticks collected from the vegetation and livestock in Nigeria. *Appl Environ Microbiol*. 2012;78:2562–8. <http://dx.doi.org/10.1128/AEM.06686-11>
5. Walker AR, Bouattour A, Camicas JL, Estrada-Peña A, Horak IG, Latif A, et al. Ticks of domestic animals in Africa. A guide to identification of species. Edinburgh (UK): Bioscience Reports; 2003.
6. Regnery RL, Spruill CL, Plikaytis BD. Genotypic identification of rickettsiae and estimation of intraspecies sequence divergence for portions of two rickettsial genes. *J Bacteriol*. 1991;173:1576–89.
7. Fournier PE, Roux V, Raoult D. Phylogenetic analysis of spotted fever group rickettsiae by study of the outer surface protein rOmpA. *Int J Syst Bacteriol*. 1998;48:839–49. <http://dx.doi.org/10.1099/00207713-48-3-839>
8. Alberdi MP, Dalby MJ, Rodriguez-Andres J, Fazakerley JK, Kohl A, Bell-Sakyi L. Detection and identification of putative bacterial endosymbionts and endogenous viruses in tick cell lines. *Ticks Tick Borne Dis*. 2012;3:137–46. <http://dx.doi.org/10.1016/j.ttbdis.2012.05.002>
9. Robinson JB, Eremeeva ME, Olson PE, Thornton SA, Medina MJ, Sumner JW, et al. New approaches to detection and identification of *Rickettsia africae* and *Ehrlichia ruminantium* in *Amblyomma variegatum* (Acari: Ixodidae) ticks from the Caribbean. *J Med Entomol*. 2009;46:942–51. <http://dx.doi.org/10.1603/033.046.0429>
10. Kelly PJ, Mason PR, Manning T, Slater S. Role of cattle in the epidemiology of tick-bite fever in Zimbabwe. *J Clin Microbiol*. 1991;29:256–9.

Address for correspondence: Kim Picozzi, Division of Pathway Medicine, Edinburgh University Medical School, The Chancellor's Bldg, 49 Little France Crescent, Edinburgh EH16 4SB, Scotland, UK; email: [kim.picozzi@ed.ac.uk](mailto:kim.picozzi@ed.ac.uk)

## Ongoing Measles Outbreak in Orthodox Jewish Community, London, UK

**To the Editor:** Measles outbreaks have been reported in Orthodox and ultra-Orthodox Jewish communities across Europe and Israel (1–5). We describe an ongoing outbreak within the largest European Orthodox Jewish community (including a Charedi population of 17,587), based in London, focused in Hackney (6). Vaccination coverage within this community is lower than in the general population of London, causing low herd immunity and outbreaks of vaccine-preventable diseases. Vaccination coverage data within the communities cannot be extrapolated, because membership is not classified as an ethnicity and not collected within health electronic recording systems. However, general practice surgeries in Hackney known to have high proportions of Orthodox Jewish patients have considerably lower vaccination coverage (55%–75% of patients 24 months of age had received measles, mumps, rubella [MMR] vaccine in the 3rd quarter of 2012) compared with the London average (87.3%) (7). Health beliefs, family size (the average Charedi household size is 6.3 persons), and underutilization of immunization services contribute to low coverage (8,9).

The outbreak clinical case definition was taken from Public Health England's guidance (10). It also included membership in the Orthodox

# *Rickettsia africae* in *Amblyomma variegatum* Ticks, Uganda and Nigeria

## Technical Appendix

Technical Appendix Table. Results of PCR screening of *Amblyomma variegatum* ticks, Uganda and Nigeria, 2010\*

Study area	No. ticks Identified			No. PCR-positive/total											
	M	F	N	<i>gltA</i>				<i>ompA</i>				<i>gltA and ompA</i>			
				Total	M	F	N	Total	M	F	N	Total	M	F	N
Uganda, n = 39	32	5 (4)	2	16/39	12/32	4 (3)/5 (4)	0/2	26/39	24/32	2 (2)/5 (4)	0/2	13/39	11/32	2 (0)/5 (4)	0/2
Nigeria, n = 141	80	59 (28)	2 (1)	84/141	45/80	38 (19)/59 (28)	1 (1)/2 (1)	88/141	44/80	44 (22)/59 (28)	0/2(1)	68/141	32/80	36 (18)/59 (28)	0/2(1)
Total, n = 180	112	64 (32)	4 (1)	100/180	57/112	42 (22)/64 (32)	1 (1)/4 (1)	114/180	68/112	46 (24)/64 (32)	0/4 (1)	81/180	43/112	38 (18)/64 (32)	0/4(1)

\*M, male; F, female; N, nymph. Numbers in parentheses indicate engorged female ticks and nymphs.