

no comparative uptake data are available to supplement our evaluation of the intervention.

Information relating to the outbreak was placed in 2 Orthodox Jewish newspapers and targeted information for families (in English, Yiddish, and Hebrew) has been disseminated. Finally, all 25 HPTs were alerted to this outbreak and the national Public Health England database (HPZone) has been enhanced to capture notifications from Orthodox Jewish communities.

This ongoing outbreak highlights continued health risks in communities with low vaccination coverage. The outbreak has been largely contained within London's Orthodox Jewish communities, with limited spread outside of the city and to just 1 local non-Orthodox Jewish child. Given the mobility of members, the risk for transmission outside of London is relatively high. The outbreak underscores the need for ongoing evidence-based and culturally appropriate health interventions that seek to improve vaccination coverage.

**Vanessa Baugh, Jose Figueroa,
Joanne Bosanquet,
Philippa Kemsley,
Sarah Addiman,
and Deborah Turbitt**


Author affiliations: Public Health England, London, UK (V. Baugh, J. Bosanquet, S. Addiman, D. Turbitt, J. Figueroa); and Homerton University Hospital Foundation Trust, London (P. Kemsley)

DOI: <http://dx.doi.org/10.3201/eid1910.130258>

References

- Lernout T, Kissling E, Hutse V, Schrijver KD, Top G. An outbreak of measles in Orthodox Jewish communities in Antwerp, Belgium, 2007–2008: different reasons for accumulation of susceptibles. *Euro Surveill.* 2009;14:19087.
- Stein-Zamir C, Abramson N, Shoob H, Zentner G. An outbreak of measles in an ultra-Orthodox Jewish community in Jerusalem, Israel, 2007—an in-depth report. *Euro Surveill.* 2008;13:8045.
- Cohen BJ, McCann R, van den Bosch C, White J. Outbreak of measles in an Orthodox Jewish community. *Euro Surveill.* 2000;4:1675.
- Ashmore J, Addiman S, Cordery R, Maguire H. Measles in North East and North Central London, England: a situation report. *Euro Surveill.* 2007;12:3271.
- Muscat M. Who gets measles in Europe? *J Infect Dis.* 2011;204:S353–65. <http://dx.doi.org/10.1093/infdis/jir067>
- Mayhew L, Harper G, Waples S, Mayhew Harper Associates. Counting Hackney's population using administrative data—an analysis of change between 2007 and 2011 [cited 2013 Aug 2]. <http://www.hackney.gov.uk/Assets/Documents/estimating-and-profiling-the-population-of-hackney.pdf>
- Public Health England. Quarterly vaccine coverage data tables [cited 2013 Aug 2]. http://www.hpa.org.uk/webw/HPAweb&HPAwebStandard/HPAweb_C/1211441442288
- Henderson L, Millett C, Thorogood N. Perceptions of childhood immunization in a minority community: qualitative study. *J R Soc Med.* 2008;101:244–51. <http://dx.doi.org/10.1258/jrsm.2008.070363>
- Purdy S, Jones KP, Sherratt M, Fallon PV. Demographic characteristics and primary health care utilization patterns of strictly orthodox Jewish and non-Jewish patients. *Fam Pract.* 2000;17:233–5.
- Health Protection Agency. HPA national measles guidelines: local and regional services. 2010 [cited 2013 Aug 2]. http://www.hpa.org.uk/webc/HPAwebFile/HPAweb_C/1274088429847

Address for correspondence: Vanessa Baugh, North East and North Central Health Protection Team, 2nd Floor, 151 Buckingham Palace Rd, London SW1W 9SZ, UK; email: vanessa.baugh@phe.gov.uk



Human Infection with Eurasian Avian-like Influenza A(H1N1) Virus, China

To the Editor: We report a human infection with avian-like swine A(H1N1) influenza virus first identified through a surveillance system for influenza like illness (ILI) in mainland China. An influenza virus, isolated from a patient with ILI, was originally subtyped as influenza A(H1N1)pdm09 virus with a hemagglutination inhibition (HI) test, but it was identified as a Eurasian avian-like influenza A(H1N1) virus (EA-H1N1) by full genome sequencing on January 30, 2013. The virus was named A/Hebei-Yuhua/SWL1250/2012 (H1N1v) (HB/1250/12), according to the definition of the World Health Organization (1).

The case-patient was a 3-year-old boy who had symptoms of fever and sore throat; his highest body temperature was 38°C on December 9, 2012. He was brought for medical treatment to an influenza sentinel hospital in the city of Shijiazhuang in Hebei Province, China, on December 12. He recovered within a week without hospitalization and oseltamivir treatment. A throat swab specimen was collected and sent to the local Chinese Center for Disease Control and Prevention for virus isolation and characterization, according to the Guidelines of the Chinese National Influenza Surveillance Network. A retrospective investigation was conducted to identify the potential infection source and any other possible cases. The case-patient was previously healthy and had no history of close contact with animals (live or dead wild birds, poultry, and swine) within 2 weeks before the onset of symptoms, nor a history of travel. He lived with his sister and parents; all other family members did not

develop influenza-like symptoms during the period of the investigation.

Sporadic human infections with swine influenza virus had been reported previously (2,3). Another case-patient, infected by EA-H1N1 influenza virus A/Jiangsu/1/2011(JS11) in early 2011, was reported (4,5). The genome sequences of the viruses isolated from the 2 case-patients showed high homology; the similarity of the polymerase basic protein 2 was 99.1%; of polymerase basic protein 1, 99.3%; of polymerase acidic protein, 98.9%; of hemagglutinin (HA), 99.1%; nucleocapsid protein, 99.1%; neuraminidase protein, 99.2%; matrix protein, 99.6% and of nonstructural protein, 99.2% (Global Initiative on Sharing Avian Influenza Data, GISAID, accession no.EPI301156–63 for JS11 and EPI438417–25 for HB/1250/12). The HB/1250/12 virus has the amino acids D (at site 190) and E (at site 225) within the HA protein, which are reported to be critical for enhancement of the HA affinity in binding to α -2,6-linked sialosides (6). The virus was resistant to amantadine and rimantadine with S31N (Ser→Asn) mutation in M2 gene, but was predicted to be susceptible to the neuraminidase inhibitor drugs oseltamivir and zanamivir on the basis of the neuraminidase gene.

HI test with ferret anti-serum against A(H1N1)pdm09 (CA09), seasonal H3N2(Vic11, BR10/07 and Perth09), classical swine subtype H1N1(NJ76), and the seasonal

influenza subtype H1N1 viruses (BR59/07, SI06) showed that the HB/1250/12 virus is antigenically indistinguishable from NJ76 and CA09, but different from subtype H3N2 viruses (Vic11, BR10/07, and Perth09) and seasonal subtype H1N1 viruses(BR59/07, SI06,) (online Technical Appendix Table; wwwnc.cdc.gov/EID/article/19/10/13-0420-Techapp1.pdf). These findings were consistent with results reported previously (7–9).

To estimate the susceptibility of human population to this virus, and to investigate whether seasonal trivalent inactivated influenza vaccine (TIV) could provide cross-protection, we collected serum samples from children, adults, and elderly adults, before and after 2012–2013 TIV vaccination, and the antibody against HB/1250/12 virus was tested by HI assay. The seroprotection antibody was defined as HI titers ≥ 40 . Before vaccination, 28% of children (3–5 years) and 6.7% of adults (18–59 years) had HI titers ≥ 40 , but elderly adults (>60 years) did not. Samples from 56% of children, 56.7% of adults, and 26.7% of elderly adults had HI titers ≥ 40 after TIV vaccination; however, a 4-fold antibody rise developed in <30% in any age group (Table). These results indicated that a proportion of cross-protective antibody against EA-H1N1 exists in children and adults, whereas elderly adults are the most susceptible to EA-H1N1 infection with no cross-protective antibody, the vaccination

with TIV could not substantially improve the level of cross-reactive EA-H1N1 antibodies.

Antisera from hyperimmune sheep are usually used for influenza virus typing and subtyping, the CA09 sheep antisera reacted well with the HB/1250/12 virus (online Technical Appendix Table). This is the reason why the local Chinese Center for Disease Control and Prevention originally subtyped HB/1250/12 as A(H1N1)pdm09 virus. Such avian-like H1N1 virus could be missed with regular HI test. In addition, a large proportion of swine influenza infection cases are mild and even asymptomatic (2); thus, the human infections with swine influenza virus may have been underestimated in China.

This is the first human case of EA-H1N1 infection identified through the national ILI surveillance network in China, indicating that the influenza surveillance network not only plays a critical role in monitoring the seasonal influenza circulation and the vaccine virus selection, but also is useful for early detection of novel influenza viruses with pandemic potential. This study also highlighted the value of, and urgent demand for, a cost-effective sequencing platform on routine influenza surveillance for pandemic preparedness.

This work was partly supported by National Basic Research Program of China (973 program, no. 2011CB504704) and the China Mega-Project for Infectious Disease (no. 2012ZX10004215).

Table. Cross-reactive antibody response against avian-like influenza A(H1N1) virus in pediatric and adult recipients of seasonal trivalent inactivated influenza vaccines, China, 2013*†

Age group, y	Antigen	Increase ≥ 4 , %‡	Geometric mean titer		%Titer ≥ 40		%Titer ≥ 160	
			Before vac.	After vac.	Before vac.	After vac.	Before vac.	After vac.
Children, 3–5, n = 25	A/California/7/2009	60.0	21.1	121.3	44.0	84.0	16.0	60.0
	HB/1250/12	24.0	12.8	25.7	28.0	56.0	0	24.0
Adults, 18–59, n = 30	A/California/7/2009	70.0	14.8	156.3	26.7	76.7	0	60.0
	HB/1250/12	6.7	8.1	31.7	6.7	56.7	0	23.3
Elderly adults, ≥ 60 , n = 30	A/California/7/2009	46.7	10.5	52.8	10.0	53.3	0	30.0
	HB/1250/12	26.7	5.7	11.5	0	26.7	0	6.7

*Vac., vaccination.

†All children received 2 doses of vaccine with an interval of 1 month. The composition of the trivalent vaccine were A/Christchurch/16/2010(NIB-74xp) (A/California/7/2009-like), A/Victoria/361/2011 (H3N2)IVR-165, and B/Hubei-Wujiagang/158/2009. Serum samples were obtained from vaccine recipients living in northern (children) and southern (adults and elderly adults) China.

‡Increase in antibody titer.

**Da-Yan Wang,¹ Shun-Xiang Qi,¹
Xi-Yan Li, Jun-Feng Guo,
Min-Ju Tan, Guang-Yue Han,
Yan-Fang Liu, Yu Lan,
Lei Yang, Wei-Juan Huang,
Yan-Hui Cheng, Xiang Zhao,
Tian Bai, Zhao Wang,
He-Jiang Wei, Ning Xiao,
and Yue-Long Shu**

Authors affiliations: National Institute for Viral Disease Control and Prevention, China CDC, Beijing, People's Republic of China (D.-Y. Wang, X.-Y. Li, J.-F. Guo, M.-J. Tan, Y. Lan, L. Yang, W.-J. Huang, Y.-H. Cheng, X. Zhao, T. Bai, Z. Wang, N. Xiao, H.-J. Wei, Y.-L. Shu); and Hebei Center for Disease Control and Prevention, Hebei Province, People's Republic of China (S.-X. Qi, G.-Y. Han, Y.-F. Liu)

DOI: <http://dx.doi.org/10.3201/eid1910.130420>

References

1. World Health Organization. Standardization of terminology for the variant A(H3N2) virus recently infecting humans [cited 2013 Mar 12]. http://www.who.int/influenza/gisrs_laboratory/terminology_ah3n2v/en/indexhtml.
2. Myers KP, Olsen CW, Gray GC. Cases of swine influenza in humans: a review of the literature. *Clin Infect Dis*. 2007;44:1084–8. <http://dx.doi.org/10.1086/512813>
3. Adiego Sancho B, Omenaca Teres M, Martínez Cuenca S, Rodrigo Val P, Sanchez Villanueva P, Casas I, et al. Human case of swine influenza A (H1N1), Aragon, Spain, November 2008. *Euro Surveill*. 2009;14: pi.19120.
4. Qi X, Cui L, Jiao Y, Pan Y, Li X, Zu R, et al. Antigenic and genetic characterization of a European avian-like H1N1 swine influenza virus from a boy in China in 2011. *Arch Virol*. 2013; 158: 39–53.
5. Yang H, Qiao C, Tang X, Chen Y, Xin X, Chen H. Human infection from swine-like influenza A (H1N1) viruses in pigs, China. *Emerg Infect Dis*. 2012;18:1144–6. <http://dx.doi.org/10.3201/eid1807.120009>
6. Matrosovich M, Tuzikov A, Bovin N, Gambaryan A, Klimov A, Castrucci MR, et al. Early alterations of the receptor-binding properties of H1, H2, and H3 avian influenza virus hemagglutinins after their introduction into mammals. *J Virol*. 2000;74:8502–12. <http://dx.doi.org/10.1128/JVI.74.18.8502-8512.2000>
7. Vijaykrishna D, Smith GJ, Pybus OG, Zhu H, Bhatt S, Poon LL, et al. Long-term

- evolution and transmission dynamics of swine influenza A virus. *Nature*. 2011;473:519–22. <http://dx.doi.org/10.1038/nature10004>
8. Vijaykrishna D, Poon LL, Zhu HC, Ma SK, Li OT, Cheung CL, et al. Reassortment of pandemic H1N1/2009 influenza A virus in swine. *Science*. 2010;328:1529. <http://dx.doi.org/10.1126/science.1189132>
 9. Zhu H, Zhou B, Fan X, Lam TT, Wang J, Chen A, et al. Novel reassortment of Eurasian Avian-like and pandemic/2009 influenza viruses in swine: infectious potential to humans. *J Virol*. 2011; 85:10432–9. <http://dx.doi.org/10.1128/JVI.05352-11>

Address for correspondence: Yue-long Shu, National Institute for Viral Disease Control and Prevention, Chinese Centre for Disease Control and Prevention, Beijing, 102206, People's Republic of China; email: yshu@cnic.org.cn.

Novel Bat Coronaviruses, Brazil and Mexico

To the Editor: Bats are now recognized as natural reservoirs for many families of viruses that can cross species barriers and cause emerging diseases of humans and animals. Protecting humans against emerging diseases relies on identifying natural reservoirs for such viruses and surveillance for host-jumping events. The emergence of the Middle East respiratory syndrome coronavirus (MERS-CoV) on the Arabian Peninsula (1) further justifies increased surveillance for coronaviruses (CoVs) in bats. MERS-CoV most likely is a zoonotic virus from a bat reservoir and is associated with high case-fatality rates among humans. The existence of a diverse array of alphacoronaviruses in bats in the United States, Canada, and Trinidad has been reported (2–6). Recently, a possible new alphacoronavirus was detected in an urban roost of bats in southern Brazil (7), and a survey of bats in southern Mexico reported 8 novel alphacoronaviruses and 4 novel

betacoronaviruses, 1 with 96% similarity to MERS-CoV (8). These findings expand the diversity and range of known bat coronaviruses and increase the known reservoir for potential emerging zoonotic CoVs.

Expanding on our previous work (2,3), we analyzed samples from 97 bats from Brazil and 75 bats from Mexico (Technical Appendix, wwwnc.cdc.gov/EID/article/19/10/13-0525-Techapp1.pdf). During 2007–2010, intestinal samples were collected from bats of 10 species in northwest São Paulo state in southeastern Brazil. These bats had been submitted to the University Estadual Paulista for rabies testing as a result of epidemiologic surveillance or, in some cases, because of possible or known contact with humans. During 2011–2012, as part of an ongoing rabies surveillance project, intestinal samples were collected from bats of 12 species in their usual habitats in Jalisco state in midwestern Mexico. Bats from a variety of species, including insectivorous, nectarivorous, frugivorous, and hematophagous bats, were included in this study for the purpose of obtaining a diverse array of potential exposures. Intestines were collected and stored, and RNA was purified as described (2). CoV RNA was detected by using a pancoronavirus PCR selective for the RNA-dependent RNA polymerase gene, and amplicons were sequenced as described (3). Virus isolation was not attempted as part of this study.

From 1 of 17 *Molossus rufus* bats and 1 of 8 *Molossus molossus* bats, an identical novel alphacoronavirus was detected (BatCoV-*M.rufus*28/Brazil/2010, GenBank accession no. KC886321). Both specimens were collected in Brazil during 2010 from adult male bats that had been found in urban areas on residential property. The 412-nt sequence of this virus was most closely related to alphacoronaviruses detected in *Eptesicus fuscus* bats in North America (82% nt

¹These authors contributed equally to this article.

Human Infection with Eurasian Avian-like Influenza A(H1N1) Virus, China

Technical Appendix

Technical Appendix Table. Cross-reactive antibody response against avian-like influenza A(H1N1) virus in pediatric and adult recipients of seasonal trivalent inactivated influenza vaccines, China, 2013*†

Age group, y	Antigen	Increase ≥4, %‡	Geometric mean titer		Titer ≥40, %		Titer ≥160, %	
			Before vac	After vac	Before vac	After vac	Before vac	After vac
Children, n = 25	A/California/7/2009	60.0	21.1	121.3	44.0	84.0	16.0	60.0
3–5	HB/1250/12	24.0	12.8	25.7	28.0	56.0	0	24.0
Adults, n = 30	A/California/7/2009	70.0	14.8	156.3	26.7	76.7	0	60.0
18–59	HB/1250/12	6.7	8.1	31.7	6.7	56.7	0	23.3
Elderly adults, n = 30	A/California/7/2009	46.7	10.5	52.8	10.0	53.3	0	30.0
≥60	HB/1250/12	26.7	5.7	11.5	0	26.7	0	6.7

*Vac, vaccination.

†All children received 2 doses of vaccine with an interval of 1 month. The composition of the trivalent vaccine were A/Christchurch/16/2010(NIB-74xp) (A/California/7/2009-like), A/Victoria/361/2011 (H3N2)IVR-165, and B/Hubei-Wujiagang/158/2009. Serum samples were obtained from vaccine recipients living in northern (children) and southern (adults and elderly adults) China.

‡Increase in antibody titer.