

This study was partly supported by grants-in-aid from the Ministry of Health, Labour and Welfare of Japan (H22-Shokuhin-Ippan-012, H23-Shinko-Shitei-020, H24-Shokuhin-Ippan-007, and H24-Shokuhin-Ippan-008).

**Hidemasa Izumiya,  
Jun Terajima, Shouji Yamamoto,  
Makoto Ohnishi,  
Haruo Watanabe, Akemi Kai,  
Takayuki Kurazono,  
Masumi Taguchi, Tetsuo Asai,  
Masato Akiba, Yuko Matsumoto,  
and Yutaka Tamura**

Author affiliations: National Institute of Infectious Diseases, Tokyo, Japan (H. Izumiya, J. Terajima, S. Yamamoto, M. Ohnishi, H. Watanabe); Tokyo Metropolitan Institute of Public Health, Tokyo (A. Kai); Saitama Institute of Public Health, Saitama, Japan (T. Kurazono); Osaka Prefectural Institute of Public Health, Osaka, Japan (M. Taguchi); National Veterinary Assay Laboratory, Tokyo (T. Asai); National Institute of Animal Health, Ibaraki, Japan (M. Akiba); Yokohama City Institute of Health, Kanagawa, Japan (Y. Matsumoto); and Rakuno Gakuen University, Hokkaido, Japan (Y. Tamura)

DOI: <http://dx.doi.org/10.3201/eid1905.121395>

## References

1. Threlfall EJ. Epidemic *Salmonella typhimurium* DT 104—a truly international multiresistant clone. *J Antimicrob Chemother.* 2000;46:7–10. <http://dx.doi.org/10.1093/jac/46.1.7>
2. Briggs CE, Fratamico PM. Molecular characterization of an antibiotic resistance gene cluster of *Salmonella typhimurium* DT104. *Antimicrob Agents Chemother.* 1999;43:846–9.
3. van Belkum A. Tracing isolates of bacterial species by multilocus variable number of tandem repeat analysis (MLVA). *FEMS Immunol Med Microbiol.* 2007;49:22–7. <http://dx.doi.org/10.1111/j.1574-695X.2006.00173.x>
4. Lindstedt BA, Vardund T, Aas L, Kapperud G. Multiple-locus variable-number tandem-repeats analysis of *Salmonella enterica* subsp. *enterica* serovar Typhimurium using PCR multiplexing and multicolor capillary electrophoresis. *J Microbiol Methods.* 2004;59:163–72. <http://dx.doi.org/10.1016/j.mimet.2004.06.014>
5. Larsson JT, Torpdahl M, Petersen RF, Sorensen G, Lindstedt BA, Nielsen EM. Development of a new nomenclature for *Salmonella typhimurium* multilocus variable number of tandem repeats analysis (MLVA). *Euro Surveill.* 2009;14:19174 <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19174>
6. Anderson ES, Ward LR, Saxe MJ, de Sa JD. Bacteriophage-typing designations of *Salmonella typhimurium*. *J Hyg (Lond).* 1977;78:297–300. <http://dx.doi.org/10.1017/S0022172400056187>
7. Hermans AP, Abee T, Zwietering MH, Aarts HJ. Identification of novel *Salmonella enterica* serovar Typhimurium DT104-specific prophage and nonprophage chromosomal sequences among serovar Typhimurium isolates by genomic subtractive hybridization. *Appl Environ Microbiol.* 2005;71:4979–85. <http://dx.doi.org/10.1128/AEM.71.9.4979-4985.2005>

Address for correspondence: Hidemasa Izumiya, 1-23-1 Toyama, Shinjuku-ku, Tokyo, 162-8640, Japan; email: [izumiya@nih.go.jp](mailto:izumiya@nih.go.jp)

## Single Genotype of *Anaplasma phagocytophilum* Identified from Ticks, Camargue, France

**To the Editor:** Granulocytic anaplasmosis is a tickborne zoonosis caused by *Anaplasma phagocytophilum* bacteria, which are emerging in Europe. Besides infecting humans, *A. phagocytophilum* infect a wide range of wild and domestic mammals (1). In Europe, the *Ixodes ricinus* tick is the main vector for the bacteria, but *A. phagocytophilum* has also been detected in association with *Rhipicephalus* and *Dermacentor* spp. ticks (2). The climate and biotopes of the Mediterranean region are particularly favorable for several species of ticks and, therefore, for tickborne diseases.

Although *I. ricinus* ticks are rare or absent in the Mediterranean Basin, serosurveys performed on equine

populations in Camargue, southern France, indicated an *A. phagocytophilum* seroprevalence of ≈10% (3). To investigate the prevalence and diversity of *A. phagocytophilum* bacteria in ticks in Camargue, we collected questing ticks from horse pastures and feeding ticks from horses.

Ticks feeding on horses were collected in randomly selected stables during 2007 (84 stables), 2008 (72 stable), and 2010 (19 stables). The stables were chosen among those where evidence of *A. phagocytophilum* seroconversion in horses had been previously found (3). In 2008 and 2010, questing ticks were collected by the dragging method in 19 pastures, around bushes, and in areas where horses spent the most time. Surveys were conducted in the spring, which represents the peak activity time of *Ixodes* ticks.

A total of 406 adult ticks were collected, representing 6 species: *Rhipicephalus bursa*, *R. sanguineus*, *R. turanicus*, *R. pusillus*, *Dermacentor marginatus*, and *Hyalomma marginatum*. Tick species were identified by morphologic criteria and molecular analyses based on mitochondrial 12S rDNA sequences (4). Total DNA was extracted from the ticks by using the NucleoSpin Tissue Kit (Macherey-Nagel, Düren, Germany) (5). *A. phagocytophilum* was detected by nested PCR targeting the 16S rDNA (online Technical Appendix 1, [wwwnc.cdc.gov/EID/article/19/5/12-1003-Techapp1.pdf](http://wwwnc.cdc.gov/EID/article/19/5/12-1003-Techapp1.pdf)).

Of the 406 ticks, 40 were infected with *A. phagocytophilum*. The infected group included ticks from all 6 collected species except *R. pusillus*. Infection rates among the species ranged from 0 to 22% (online Technical Appendix 2, [wwwnc.cdc.gov/EID/article/19/5/12-1003-Techapp2.pdf](http://wwwnc.cdc.gov/EID/article/19/5/12-1003-Techapp2.pdf)). The prevalence of *A. phagocytophilum* infection did not differ significantly between species (logistic regression model,  $p = 0.76$ ) but was higher among questing ticks than feeding ticks ( $p < 0.001$ ; odds ratio 1.15).




Figure. Phylogenetic networks of *Anaplasma phagocytophilum* based on *msp4* (A) and *ankA* (B) genes and built with SplitsTree4 (version 4.11.3; <http://splitstree.org/>) by the Neighbor-Net method. The sequences of the genotype described in Camargue, France, is framed. Sequences found in the 2 networks are in **boldface**: *A. phagocytophilum* amplified from ticks collected in Combrailles, Auvergne region, France (JX197116–JX197126 and JX197265–JX197270), a human isolate (strain Webster, EU857674 and GU236811), an American roe deer isolate (strain CRT, JX197261 and JX197113), and an American horse isolate (strain MRK, AY530196 and AF153716). Scale bars indicate number of nucleotide substitutions per site.

We amplified 6 loci by nested PCR (online Technical Appendix 1) to characterize *A. phagocytophilum* genetic diversity in positive samples: *ankA*, *msp4*, *pleD*, *typA*, and intergenic regions *hemE*–*APH\_0021* and *APH\_1099*–*APH\_1100* (National Center for Biotechnology Information annotation). The GenBank accession numbers for the nucleotide sequences are JX197073–JX197368. No polymorphism was found among the 6 loci tested in the 40 *A. phagocytophilum*-positive ticks. The genotype identified was 100% identical to the reference sequence (NC\_007797) for loci *msp4*, *pleD*, and *typA* and for intergenic regions *hemE*–*APH\_0021* and *APH\_1099*–*APH\_1100*. The *ankA* sequence was 96% similar (487 nt) to the reference sequence. The relevance of

these loci as markers of diversity was verified (online Technical Appendix 3, [wwwnc.cdc.gov/EID/article/19/5/12-1003-Techapp3.pdf](http://wwwnc.cdc.gov/EID/article/19/5/12-1003-Techapp3.pdf)).

To study the phylogenetic relationships between cognate sequences, we included in our analysis all sequences available in GenBank for genes *ankA* and *msp4*. To account for recombination events that affect *ankA* and *msp4* (data not shown) in phylogenetic analyses, we used Neighbor-Net networks (Figure). Phylogenetic analysis of *msp4* (Figure, panel A) indicated that the genotype of *A. phagocytophilum* from ticks in Camargue was included in a clade that also includes genotypes that infect humans and horses in the United States.

The diversity of *ankA* sequences has been described as 4 phylogenetic

clusters (6). All sequences obtained in our study were included in cluster I, particularly in a branch composed exclusively of sequences of *A. phagocytophilum* isolated from humans in the United States (Figure, panel B).

Previous studies investigating *A. phagocytophilum* have revealed a genetic diversity that is thought to have been caused by sympatric epidemiologic cycles involving different vectors and reservoir hosts (1,6,7). In 5 species of ticks (40 ticks total) that we collected from a 250-km<sup>2</sup> area in southern France, we found only 1 genotype of *A. phagocytophilum*, which we determined to be phylogenetically close to genotypes found in the United States. Sequences phylogenetically related to bacteria in the United States were also observed in Sardinia (8) and Sicily (9).

The low diversity we found could be explained by a recent introduction of the bacteria into the area [although *A. phagocytophilum*–seropositive horses have been found in the area since 2001 (3)] or by a selective sweep linked to the particular ticks and host reservoir in Camargue. The 5 species of ticks that we found positive for *A. phagocytophilum* have been described as potential vectors of *A. phagocytophilum* in the Mediterranean Basin (2,10). Among the tick species in our investigation, *R. bursa* and *R. sanguineus* ticks are the 2 main carriers of *A. phagocytophilum*, and these ticks are likely to feed on humans and, thus, pose a risk of infection to the local population. Further studies are needed to address the potential effect of *A. phagocytophilum*–infected ticks on human health in this area and, more specifically, the relationship between genotype and pathogenicity.

#### Acknowledgments

We thank Curtis Nelson and Friederike von Loewenich for generously providing the DNA of strains HGE1, CRT, and Webster; Véronique Bachy for actively helping us to obtain samples from domestic animals; Magalie René-Martellet and Frédéric Beugnet for helping with the morphological identification of ticks; Gillian Martin for proofreading the manuscript; Nelly Dorr for creating the databases used in this study; and Françoise Rieu-Lesme and Sébastien Masségla who were involved in laboratory work.

**Amélie Chastagner,  
Xavier Bailly, Agnès Leblond,  
Sophie Pradier,<sup>1</sup>  
and Gwenaél Vourc'h**

Author affiliations: National Center for Biotechnology Information (INRA), Saint Genès Champanelle, France (A. Chastagner, X. Bailly, A. Leblond, G. Vourc'h); University of Lyon, Marcy l'Etoile, France (A. Leblond); Ecole Nationale Vétérinaire d'Alfort, Maisons-Alfort, France (S. Pradier); and INRA ENVA ANSES, Maisons-Alfort (S. Pradier)

<sup>1</sup>Current affiliation: Ecole Nationale Vétérinaire de Toulouse, Toulouse, France.

DOI: <http://dx.doi.org/10.3201/eid1905.121003>

#### References

1. Rikihisa Y. Mechanisms of obligatory intracellular infection with *Anaplasma phagocytophilum*. *Clin Microbiol Rev*. 2011;24:469–89. <http://dx.doi.org/10.1128/CMR.00064-10>
2. de la Fuente J, Naranjo V, Ruiz-Fons F, Vicente J, Estrada-Peña A, Almazan C, et al. Prevalence of tick-borne pathogens in ixodid ticks (Acari: Ixodidae) collected from European wild boar (*Sus scrofa*) and Iberian red deer (*Cervus elaphus hispanicus*) in central Spain. *Eur J Wildl Res*. 2004;50:187–96. <http://dx.doi.org/10.1007/s10344-004-0060-1>
3. Leblond A, Pradier S, Pittel PH, Fortier G, Boireau P, Chadoeuf J, et al. An epidemiological survey of equine anaplasmosis (*Anaplasma phagocytophilum*) in southern France [in French]. *Rev Sci Tech*. 2005;24:899–908.
4. Beati L, Keirans JE. Analysis of the systematic relationships among ticks of the genera *Rhipicephalus* and *Boophilus* (Acari: Ixodidae) based on mitochondrial 12S ribosomal DNA gene sequences and morphological characters. *J Parasitol*. 2001;87:32–48.
5. Halos L, Jamal T, Vial L, Maillard R, Suau A, Le Menache A, et al. Determination of an efficient and reliable method for DNA extraction from ticks. *Vet Res*. 2004;35:709–13. <http://dx.doi.org/10.1051/vetres:2004038>
6. Scharf W, Schauer S, Freyburger F, Petrovec M, Schaarschmidt-Kiener D, Liebisch G, et al. Distinct host species correlate with *Anaplasma phagocytophilum ankA* gene clusters. *J Clin Microbiol*. 2011;49:790–6. <http://dx.doi.org/10.1128/JCM.02051-10>
7. Bown KJ, Lambin X, Ogden NH, Begon M, Telford G, Woldehiwet Z, et al. Delineating *Anaplasma phagocytophilum* ecotypes in coexisting, discrete enzootic cycles. *Emerg Infect Dis*. 2009;15:1948–54. <http://dx.doi.org/10.3201/eid1512.090178>
8. Alberti A, Zobba R, Chessa B, Addis MF, Sparagano O, Parpaglia MLP, et al. Equine and canine *Anaplasma phagocytophilum* strains isolated on the island of Sardinia (Italy) are phylogenetically related to pathogenic strains from the United States. *Appl Environ Microbiol*. 2005;71:6418–22. <http://dx.doi.org/10.1128/AEM.71.10.6418-6422.2005>
9. de la Fuente J, Torina A, Caracappa S, Tumino G, Furla R, Almazan C, et al. Serologic and molecular characterization of *Anaplasma* species infection in farm animals and ticks from Sicily. *Vet Parasitol*. 2005;133:357–62. <http://dx.doi.org/10.1016/j.vetpar.2005.05.063>

10. Toledo A, Olmeda AS, Escudero R, Jado I, Valcarcel F, Casado-Nistal MA, et al. Tick-borne zoonotic bacteria in ticks collected from central Spain. *Am J Trop Med Hyg*. 2009;81:67–74.

Address for correspondence: Amélie Chastagner, INRA, UR346 Epidémiologie Animale, Centre de Recherches de Clermont-Ferrand/Theix, F-63122 Saint Genès Champanelle, France; email: [amelie.chastagner@clermont.inra.fr](mailto:amelie.chastagner@clermont.inra.fr)

## West Nile Virus Lineage 2 Strain in Greece, 2012

**To the Editor:** West Nile virus (WNV) has been in Europe at least since the 1960s (1). Before 2010, WNV epidemics in Europe were caused mainly by lineage 1 strains. However, in 2010, a major WNV epidemic in Central Macedonia, Greece, was caused by a lineage 2 strain (Nea Santa-Greece-2010) (2).

This strain also circulated during 2011 (3), causing a second epidemic among humans throughout the country (4). Although the virus was closely related to the goshawk-Hungary-2004 strain circulating in Hungary, Austria, and Italy (5–7), severe epidemics occurred only in Greece; 273 cases of West Nile neuroinvasive disease (WNNND) in humans were reported during the 2 seasons (2,4).

A third epidemic occurred in 2012, and 109 WNNND cases were reported (8). Until mid-August, most cases were in central (Attica; 29 cases) and northeastern Greece (East Macedonia and Thrace; 10 cases). In contrast, during the same period, only 3 cases were confirmed at the location of the 2010 epidemic epicenter, in Central Macedonia (9). This situation led to the question of whether the Nea Santa-Greece-2010 strain was responsible for the third epidemic in Greece.