

for food safety rules and human health risk measures taken by national health and veterinary agencies. Regarding adequate processing of clinical samples and their preservation for morphologic and genetic evaluation, we strongly recommend fixation of positive fecal samples with eggs or segments (proglottids) immediately with 96%–99% molecular grade (i.e., not denatured) ethanol for future molecular diagnosis (1,4,8).

This work was supported by the Czech Science Foundation (grant number P506/12/1632) and the Institute of Parasitology, Biology Centre of the Academy of Sciences of the Czech Republic (grant number RVO: 60077344).

**Roman Kuchta,
José-Guillermo Esteban,
Jan Brabec, and Tomáš Scholz**

Author affiliations: Institute of Parasitology, Biology Centre of the Academy of Sciences of the Czech Republic, České Budějovice, Czech Republic (R. Kuchta, J. Brabec, T. Scholz); and Facultad de Farmacia, Universidad de Valencia, Valencia, Spain (J.-G. Esteban)

DOI: <http://dx.doi.org/10.3201/eid2011.140996>

References

- Scholz T, Garcia HH, Kuchta R, Wicht B. Update on the human broad tapeworm (genus *Diphyllobothrium*), including clinical relevance. *Clin Microbiol Rev*. 2009;22:146–60. <http://dx.doi.org/10.1128/CMR.00033-08>
- de Marval F, Gottstein B, Weber M, Wicht B. Imported diphyllobothriasis in Switzerland: molecular methods to define a clinical case of *Diphyllobothrium* infection as *Diphyllobothrium dendriticum*, August 2010. *Eurosurveillance*. 2013;18:31–6.
- Kuchta R, Brabec J, Kubáčková P, Scholz T. Tapeworm *Diphyllobothrium dendriticum* (Cestoda)—neglected or emerging human parasite? *PLoS Negl Trop Dis*. 2013;7:e2535. <http://dx.doi.org/10.1371/journal.pntd.0002535>
- Kuchta R, Scholz T, Brabec J, Wicht B. Chapter 16. *Diphyllobothrium*, *Diplogonoporus*, and *Spirometra*. In: Xiao L, Ryan U, Feng Y, editors. *Biology of foodborne parasites, section III: Important foodborne helminths*. London: CRC Press; (in press).
- Pastor-Valle J, González LM, Martín-Clemente JP, Merino FJ, Gottstein B, Gárate T. Molecular diagnosis of *diphyllobothriasis* in Spain, most presumably acquired via imported fish, or sojourn abroad. *New Microbes and New Infections*. 2014;2:1–6. <http://dx.doi.org/10.1002/2052-2975.28>
- Colomina J, Villar J, Esteban G. Asymptomatic infection by *Diphyllobothrium latum* in a Spanish 3-year-old child [in Spanish]. *Medicina Clínica (Barcelona)*. 2002;118:279. [http://dx.doi.org/10.1016/S0025-7753\(02\)72359-2](http://dx.doi.org/10.1016/S0025-7753(02)72359-2)
- Esteban JG, Muñoz-Antoli C, Borrás M, Colomina J, Toledo R. Human infection by a “fish tapeworm”, *Diphyllobothrium latum*, in a non-endemic country. *Infection*. 2014;42:191–4. <http://dx.doi.org/10.1007/s15010-013-0491-2>
- Wicht B, Yanagida T, Scholz T, Ito A, Jiménez JA, Brabec J. Multiplex PCR for differential identification of broad tapeworms (Cestoda: *Diphyllobothrium*) infecting humans. *J Clin Microbiol*. 2010;48:3111–6. <http://dx.doi.org/10.1128/JCM.00445-10>
- Gil-Setas A, Mazón A, Pascual P, Sagua H. Helminthiasis in a 71-year-old man, an infrequent condition in our setting [in Spanish]. *Enferm Infecc Microbiol Clin*. 2004;22:553–4. <http://dx.doi.org/10.1157/13067624>
- Salminen K. The effect of high and low temperature on the infestiveness of *Diphyllobothrium latum* with regard to public health [dissertation]. Helsinki: College of Veterinary Medicine; 1970. https://openlibrary.org/works/OL6530111W/The_effect_of_high_and_low_temperature_treatments_on_the_infestiveness_of_Diphyllobothrium_latum_wit

Address for correspondence: Roman Kuchta, Institute of Parasitology, Biology Centre of the Academy of Sciences of the Czech Republic, Branišovská 31, 370 05, České Budějovice, Czech Republic; email: krtek@paru.cas.cz

**EMERGING
INFECTIOUS DISEASES**

Free Online RSS Feed

in PubMed Central

Ahead of print Peer-Reviewed

CME podcasts

GovDelivery

Drug Resistance in *Salmonella enterica* ser. Typhimurium Bloodstream Infection, Malawi

To the Editor: *Salmonella enterica* serotype Typhimurium is one of the most common causes of bloodstream infection in sub-Saharan Africa (1). Among adults, the principal risk factor for invasive nontyphoidal *Salmonella* (iNTS) disease is advanced HIV infection; up to 44% of HIV-infected patients experience bacteremic recurrence through recrudescence of the original infection (2,3). Epidemics of iNTS disease in sub-Saharan Africa have been associated with a novel genotype of *S. enterica* ser. Typhimurium of multilocus sequence type (ST) 313 that is rarely seen outside the region and is associated with multidrug resistance (MDR) to chloramphenicol, cotrimoxazole, and ampicillin (4,5). As a consequence, ceftriaxone has become a key agent in the empirical management of nonfocal sepsis in Malawi (6).

In March 2009, a 40-year-old HIV-infected and antiretroviral therapy-naïve woman sought care in Blantyre, Malawi, with an MDR *S. enterica* ser. Typhimurium bloodstream infection. She was treated with ceftriaxone (2 g intravenously once daily) and discharged with oral ciprofloxacin (500 mg twice daily) for 10 days. She was readmitted 1 month later with recurrent fever. At this time, she had an MDR *S. enterica* ser. Typhimurium bloodstream infection with additional resistance to ceftriaxone and ciprofloxacin. In the absence of a locally available effective antimicrobial drug, she was treated with ceftriaxone, gentamicin, and high-dose ciprofloxacin but died shortly thereafter.

To help clarify how this extended MDR *S. enterica* ser. Typhimurium emerged, we determined the molecular mechanisms underpinning this disturbing pattern of antimicrobial resistance

(online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/20/11/14-1175-Techapp1.pdf>). We conducted phenotypic drug susceptibility testing by disk diffusion on *S. enterica* ser. Typhimurium strains A54285 (initial presentation) and A54560 (recurrence); both isolates were resistant to ampicillin, chloramphenicol, and cotrimoxazole, but A54560 exhibited additional resistance to ceftriaxone, ciprofloxacin, and tetracycline.

Paired-end sequencing of isolates A54285 (European Nucleotide Archive [ENA] accession number ERS035867) and A54560 (ENA accession no. ERS035866) that were cultured 1 month apart showed no differences between the conserved regions of these genomes (Figure). The similarity of these *S. enterica* ser.

Typhimurium genomes strongly suggests that this recrudescence occurred after incomplete clearance of the first infection; although re-infection from the same source is unlikely, it cannot be excluded. Comparison of the accessory genomes, however, showed an additional 300 kb DNA in A54560.

Plasmid extraction and gel electrophoresis of genomic DNA identified a plasmid migrating in the gel to a position approximately equivalent to 120 kb, the size of ST313 virulence plasmid pSLT-BT in both strains, but no 300-kb plasmid was visualized in the ceftriaxone- and ciprofloxacin-resistant strain (A54560, data not shown), possibly because of the difficulty large plasmids have entering standard 1% agarose gels. However, ceftriaxone resistance was mobilized

to *Escherichia coli* by conjugation at a frequency 6.5×10^{-2} transconjugants per donor at 26°C. This frequency dropped dramatically to $\approx 1 \times 10^{-7}$ transconjugants per donor when conjugation was performed at 37°C. The presence of an IncHI2 plasmid in the transconjugants was confirmed by PCR for the IncHI2 region (7), and drug susceptibility testing confirmed that transconjugant clones acquired resistance to ceftriaxone, ciprofloxacin, and tetracycline.

These data confirm the presence of an extended-spectrum β -lactamase (ESBL)-producing IncHI2 plasmid in strain A54560 that is capable of conjugative transfer and suggest that the plasmid might have been acquired by residual index strain within the patient by transfer from an unknown donor




Figure. Midpoint-rooted phylogenetic tree of published whole-genome sequence data from D23580-like *Salmonella enterica* serotype Typhimurium sequence type 313s from Malawi based on 204 informative single-nucleotide polymorphisms. A54285 and A54560, highlighted in gray, are indistinguishable. Scale bar indicates nucleotide substitutions per site.

bacterium. Partial decolonization of the patient's gastrointestinal tract by ceftriaxone and fluoroquinolone antimicrobial therapy might have rendered it receptive to colonization by ESBL-producing bacteria, which we hypothesize donated the plasmid to the residual index strain.

The transconjugant plasmid DNA was sequenced by using the PacBio RSII platform (Pacific Biosciences, Menlo Park, CA, USA; <http://www.pacificbiosciences.com>), which assembled as a single contiguous sequence of 309,406 bp, designated pSTm-BTCR (online Technical Appendix Figure, ENA accession no. LK056646). We identified 331 predicted coding sequences, including 109 genes required for replication and transfer and 61 genes predicted to be associated with metabolism, membranes, virulence, antimicrobial resistance, and a toxin/antitoxin addiction system. We found an additional 160 predicted, hypothetical genes. Fifteen putative antimicrobial resistance genes were identified, predicted to encode resistance to; tetracycline (*tetA(C)*, *tetR(C)*), β -lactams (*bla*_{CTX-M15'}, *bla*_{TEM-1b'}, *bla*_{OXA-30}), chloramphenicol (*catB3*, *catA1*), aminoglycosides (*strA*, *strB*, *aadA1*, *aacA4*, *aacC3*), ciprofloxacin (*qnrB1*), ulfonamiides (*sul2*), and trimethoprim (*dfpA14*).

In our experience, ESBL and fluoroquinolone-resistant iNTS remain extremely uncommon in Blantyre, Malawi. This is surprising because diverse ESBL genotypes were observed in other members of *Enterobacteriaceae* in Blantyre within a year after ceftriaxone came into common use locally (8). That IncHI2 plasmids transfer most efficiently at temperatures <30°C (9), a lower temperature than in the human gastrointestinal tract, might explain why the acquisition of ESBL-producing enzymes through IncHI2 plasmids has not been commonly observed within patients with recurrent iNTS disease in this setting. However, rates of

transfer might differ when bacteria are growing in the intestine.

The spread of mobile genetic elements that confer antimicrobial resistance among gram-negative organisms is of considerable concern. Wide dissemination of this strain or the IncHI2 (pSTm-BTCR) plasmid among other salmonellae in sub-Saharan Africa would rapidly render iNTS effectively untreatable with currently available antibacterial drugs.

This work was supported by the Wellcome Trust; N.A.F. holds a Wellcome Research Training Fellowship. The Malawi Liverpool Wellcome Trust Clinical Research Programme and the Wellcome Trust Sanger Institute are core funded by the Wellcome Trust.

**Nicholas A. Feasey,
Amy K. Cain,
Chisomo L. Msefula,
Derek Pickard, Maaïke Alaerts,
Martin Aslett, Dean B. Everett,
Theresa J. Allain,
Gordon Dougan,
Melita A. Gordon,
Robert S. Heyderman,
and Robert A. Kingsley**

Author affiliations: Liverpool School of Tropical Medicine, Liverpool, UK (N.A. Feasey, R.S. Heyderman); Wellcome Trust Sanger Institute, Cambridge, UK (N.A. Feasey, A.K. Cain, D. Pickard, M. Aslett, G. Dougan, R.A. Kingsley); University of Malawi College of Medicine, Blantyre, Malawi (N.A. Feasey, C.L. Msefula, M. Alaerts, D.B. Everett, T.J. Allain, R.S. Heyderman); University of Liverpool, Liverpool (D.B. Everett, M.A. Gordon); and Institute of Food Research, Colney, Norwich, UK (R.A. Kingsley)

DOI: <http://dx.doi.org/10.3201/eid2011.141175>

References

1. Feasey NA, Dougan G, Kingsley RA, Heyderman RS, Gordon MA. Invasive non-typhoidal *Salmonella* disease: an emerging and neglected tropical disease in Africa. *Lancet*. 2012;379:2489–99. [http://dx.doi.org/10.1016/S0140-6736\(11\)61752-2](http://dx.doi.org/10.1016/S0140-6736(11)61752-2)

2. Okoro CK, Kingsley RA, Quail MA, Kankwatira AM, Feasey NA, Parkhill J, et al. High-resolution single nucleotide polymorphism analysis distinguishes recrudescence and reinfection in recurrent invasive nontyphoidal *Salmonella* Typhimurium disease. *Clin Infect Dis*. 2012;54:955–63. <http://dx.doi.org/10.1093/cid/cir1032>
3. Gordon MA, Banda HT, Gondwe M, Gordon SB, Boeree MJ, Walsh AL, et al. Non-typhoidal *Salmonella* bacteraemia among HIV-infected Malawian adults: high mortality and frequent recrudescence. *AIDS*. 2002;16:1633–41. <http://dx.doi.org/10.1097/00002030-200208160-00009>
4. Gordon MA, Graham SM, Walsh AL, Wilson L, Phiri A, Molyneux E, et al. Epidemics of invasive *Salmonella enterica* serovar Enteritidis and *S. enterica* serovar Typhimurium infection associated with multidrug resistance among adults and children in Malawi. *Clin Infect Dis*. 2008;46:963–9. <http://dx.doi.org/10.1086/529146>
5. Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, et al. Epidemic multiple drug resistant *Salmonella* Typhimurium causing invasive disease in sub-Saharan Africa have a distinct genotype. *Genome Res*. 2009;19:2279–87. <http://dx.doi.org/10.1101/gr.091017.109>
6. Feasey NA, Houston A, Mukaka M, Komrower D, Mwalukomo T, Tenthani L, et al. A reduction in adult blood stream infection and case fatality at a large African hospital following antiretroviral therapy roll-out. *PLoS ONE*. 2014;9:e92226. <http://dx.doi.org/10.1371/journal.pone.0092226>
7. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods*. 2005;63:219–28. <http://dx.doi.org/10.1016/j.mimet.2005.03.018>
8. Gray KJ, Wilson LK, Phiri A, Corkill JE, French N, Hart CA. Identification and characterization of ceftriaxone resistance and extended-spectrum β -lactamases in Malawian bacteraemic *Enterobacteriaceae*. *J Antimicrob Chemother*. 2006;57:661–5. <http://dx.doi.org/10.1093/jac/dkl037>
9. Taylor DE, Levine JG. Studies of temperature-sensitive transfer and maintenance of H incompatibility group plasmids. *J Gen Microbiol*. 1980;116:475–84.

Address for correspondence: Nicholas A Feasey, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA, UK; email: nfeasey@liverpool.ac.uk

The opinions expressed by the authors contributing to this journal do not necessarily reflect the opinions of the Centers for Disease Control and Prevention or the institution with which the authors are affiliated.

Drug Resistance in *Salmonella enterica* ser. Typhimurium Bloodstream Infection, Malawi

Technical Appendix

Methods

Bacteria

Salmonella Typhimurium A54285 (index multidrug-resistant) and A54560 (recurrence extended multidrug resistant) were isolated from the same patient at initial presentation and at recurrence 1 month later, respectively. Phenotypic drug susceptibility testing was undertaken by the disk diffusion method (Oxoid, Basingstoke, UK). The sequence of *Salmonella* Typhimurium D23580, an invasive isolate of multilocus sequence type ST313, isolated at Queen Elizabeth Central Hospital in 2004 (1), was used as a reference for sequence comparison. A rifampicin-resistant mutant of *Escherichia coli* C600 containing no plasmids was used as a recipient for conjugation.

Sequencing and Comparative Genome Analysis

Genomic DNA from bacteria was prepared by Wizard genomic DNA purification kit (Promega, Madison, WI, USA). Short-read Illumina sequencing libraries were prepared, and paired-end sequencing was performed on 2 µg gDNA on a HiSeq2000 platform (Illumina, San Diego, CA, USA) generating 150-bp reads (2). The whole-genome sequence of each isolate was assembled by using Velvet (<http://www.ebi.ac.uk/~zerbino/velvet>, version 1.2.03) and aligned against D23580 and its associated plasmids by using Abacas (version 1.3.1) (3). Single-nucleotide polymorphisms (SNPs) were determined with reference to strain D23580 by using SMALT, version 0.5.8 (Wellcome Trust Sanger Institute, Cambridge, UK). Contigs that failed to align with D23580 were considered as potentially unique mobile genetic elements and were further characterized by using BLASTn analysis (<http://www.ncbi.nlm.nih.gov/BLAST>) to predict whether they were plasmids; if so, in silico PCR was used to determine the plasmid incompatibility group (4).

Plasmid Isolation and DNA Extraction

Plasmid DNA was initially extracted by using an alkaline lysis method (Kado and Liu) and plasmids were profiled by visualising on a 1% agarose gel (5).

Individual plasmids were extracted following conjugal transfer from *Salmonella* Typhimurium A54560 to rifampicin-resistant *E. coli* K12 recipient strain c600, as described previously (6). Donor and recipient strains were mated by mixing 100 µL of overnight cultures on Luria-Bertani agar plates and incubated overnight at 26°C. Cells were cultured on MacConkey agar containing rifampicin (100 µg/mL) to select for *E. coli* and ceftriaxone (4 µg/mL) to select for transconjugants. The number of donor cells was determined by culture on Luria-Bertani agar containing rifampicin alone, and the transfer frequency was determined and the antimicrobial drug resistance phenotype of transconjugants was determined (Oxoid, USA), to ensure all resistances from the donor strain transferred.

Plasmid DNA was extracted from transconjugants by using an alkaline lysis method optimized for large plasmids, as previously described (7). PCR-based replicon typing on the transconjugant DNA was used to confirm the incompatibility group suggested by in silico plasmid typing (4).


Plasmid Sequencing

The transconjugant plasmid DNA was sequenced by using the PacBio RSII platform (Pacific Biosciences). Gene prediction and annotation was performed using Prokka (<http://vicbioinformatics.com>).

References

1. Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, et al. Epidemic multiple drug resistant *Salmonella* Typhimurium causing invasive disease in sub-Saharan Africa have a distinct genotype. *Genome Res.* 2009;19:2279–87 . [PubMed](#)
<http://dx.doi.org/10.1101/gr.091017.109>
2. Harris SR, Cartwright EJ, Torok ME, Holden MT, Brown NM, Ogilvy-Stuart AL, et al. Whole-genome sequencing for analysis of an outbreak of meticillin-resistant *Staphylococcus aureus*: a descriptive study. *Lancet Infect Dis.* 2013;13:130–6 . [PubMed](#) [http://dx.doi.org/10.1016/S1473-3099\(12\)70268-2](http://dx.doi.org/10.1016/S1473-3099(12)70268-2)

3. Swain MT, Tsai IJ, Assefa SA, Newbold C, Berriman M, Otto TD. A post-assembly genome-improvement toolkit (PAGIT) to obtain annotated genomes from contigs. *Nat Protoc.* 2012;7:1260–84 . [PubMed](#) <http://dx.doi.org/10.1038/nprot.2012.068>
4. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods.* 2005;63:219–28 . [PubMed](#) <http://dx.doi.org/10.1016/j.mimet.2005.03.018>
5. Kado CI, Liu ST. Rapid procedure for detection and isolation of large and small plasmids. *J Bacteriol.* 1981;145:1365–73. [PubMed](#)
6. Cain AK, Liu X, Djordjevic SP, Hall RM. Transposons related to Tn1696 in IncHI2 plasmids in multiply antibiotic resistant *Salmonella enterica* serovar Typhimurium from Australian animals. *Microb Drug Resist.* 2010;16:197–202 . [PubMed](#) <http://dx.doi.org/10.1089/mdr.2010.0042>
7. Cain AK, Hall RM. Evolution of IncHI2 plasmids via acquisition of transposons carrying antibiotic resistance determinants. *J Antimicrob Chemother.* 2012;67:1121–7 . [PubMed](#) <http://dx.doi.org/10.1093/jac/dks004>



Technical Appendix Figure. Plasmid map of pSTm-BTCR. Circles represent the 309-kbp circular plasmid with *rep1A* close to position 0. The outermost circle shows genes predicted on the forward strand, and the circle inside that shows genes predicted on the reverse strand. Red indicates genes involved in DNA replication; yellow indicates metabolism; purple indicates antimicrobial resistance or virulence; and green marks indicate hypothetical genes. Key resistance genes are annotated. The antimicrobial resistance genes are all located between 128 kbp and 182 kbp. The regions marked in blue in the next track highlights the sites of IS elements. The innermost circle shows the guanine–cytosine plot.