

4. Page LK, Anchor C, Luy E, Kron S, Larson G, Madsen L, et al. Backyard raccoon latrines and risk for *Baylisascaris procyonis* transmission to humans. *Emerg Infect Dis.* 2009;15:1530–1. <http://dx.doi.org/10.3201/eid1509.090128>
5. Kelly TG, Madhavan VL, Peters JM, Kazacos KR, Silvera VM. Spinal cord involvement in a child with raccoon roundworm (*Baylisascaris procyonis*) meningoencephalitis. *Pediatr Radiol.* 2012;42:369–73. <http://dx.doi.org/10.1007/s00247-011-2151-y>
6. Wise ME, Sorvillo FJ, Shafir SC, Ash LR, Berlin OG. Severe and fatal central nervous system disease in humans caused by *Baylisascaris procyonis*, the common roundworm of raccoons: a review of current literature. *Microbes Infect.* 2005;7:317–23. <http://dx.doi.org/10.1016/j.micinf.2004.12.005>
7. Blizzard EL, Yabsley MJ, Beck MF, Harsch S. Geographic expansion of *Baylisascaris procyonis* roundworms, Florida, USA. *Emerg Infect Dis.* 2010;16:1803–4. <http://dx.doi.org/10.3201/eid1611.100549>
8. Miyashita M. Prevalence of *Baylisascaris procyonis* in raccoons in Japan and experimental infections of the worm in laboratory animals. *Journal of Urban Living and Health Association.* 1993;37:137–51.
9. Xie Y, Zhang Z, Niu L, Wang Q, Wang C, Lan J, et al. The mitochondrial genome of *Baylisascaris procyonis*. *PLoS ONE.* 2011;6:e27066. <http://dx.doi.org/10.1371/journal.pone.0027066>
10. Sorvillo F, Ash LR, Berlin OG, Tatabe J, Degiorgio C, Morse SA. *Baylisascaris procyonis*: an emerging helminthic zoonosis. *Emerg Infect Dis.* 2002;8:355–9. <http://dx.doi.org/10.3201/eid0804.010273>

Address for correspondence: Guangyong Yang, Department of Parasitology, College of Veterinary Medicine, Sichuan Agricultural University, 46 Xinkang Rd, Ya'an, Sichuan, 625014, People's Republic of China; email: [guangyoyang@hotmail.com](mailto:guangyoyang@hotmail.com)

**EMERGING INFECTIOUS DISEASES**

Free Online RSS Feed

in PubMed Central

Ahead of print

CME Peer-Review

podcasts wed

GovDelivery



## Novel Divergent Rhabdovirus in Feces of Red Fox, Spain

**To the Editor:** Rhabdoviruses (family *Rhabdoviridae*) are enveloped single-stranded negative-sense RNA viruses belonging to the Mononegavirales order. The International Committee on Taxonomy of Viruses recognizes 11 genera (*Cytorhabdovirus*, *Ephemerovirus*, *Lyssavirus*, *Novirhabdovirus*, *Nucleorhabdovirus*, *Perhabdovirus*, *Sigmavirus*, *Sprivivirus*, *Tibrovirus*, *Tupavirus*, *Vesiculovirus*) (1). In addition, many recently described rhabdoviruses remain unassigned. Rhabdoviruses contain 5 major genes, encoding for nucleoprotein (N), phosphoprotein (P), matrix (M), glycoprotein (G), and RNA-dependent RNA polymerase (L). The *Rhabdoviridae* family includes pathogens of various animal species, humans, and plants. Viruses of the genus *Lyssavirus* are the most relevant to public health because they can cause rabies. Bats are the driving force within this genus; foxes and various other species of wild carnivores also can be infected with lyssaviruses and transmit them to humans and dogs (2).

During a viral metagenomic survey, conducted as described previously (3), of fecal samples collected from 4 red foxes (*Vulpes vulpes*) that were found dead in Álava, Basque Country, Spain, we identified the complete coding sequence and the partial leader and trailer sequence of a novel rhabdovirus, tentatively called red fox fecal rhabdovirus (RFFRV; 15,541 nt, GenBank accession no. KF823814; online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/20/12/14-0236-Techapp1.pdf>) by mapping 8,287 of the 56,519 sequence reads in the sample of a red fox. A proportion of obtained reads contained sequences that were  $\geq 99\%$  identical to mitochondrial DNA of *V. vulpes*, which confirmed

that the sample was collected from a red fox.

The obtained sequence of RFFRV was partially confirmed by specific primers and Sanger sequencing of PCR amplicons. Five major and 3 minor open reading frames (ORFs) were identified that had a genome organization similar to that of other rhabdoviruses (Figure, panel A). No significant hits were obtained by BLAST analysis (<http://blast.ncbi.nlm.gov/Blast.cgi>) of N, P, M, and G nucleotide and amino acid sequences, which was reported previously for novel divergent rhabdoviruses (4).

Predicted N, P, and M genes of RFFRV consist of 1,629, 2,490, and 813 nt, respectively, encoding for 543, 830, and 271 aa (online Technical Appendix Table 1). In addition to the absence of significant hits observed by BLAST analysis, no significant sequence homology was observed with known rhabdovirus proteins in pairwise alignments. Furthermore, no conserved motifs were detected in N, P, and M genes of RFFRV that are commonly observed in rhabdoviruses. However, intergenic regions between all major ORFs contained relatively conserved motifs that could be transcription termination/polyadenylation sequences (A/U) CU<sub>7</sub>, similar to other rhabdoviruses (5). Adjacent to this termination signal was a stretch of conserved nucleotides that might function as a transcription initiation signal (online Technical Appendix Table 1).

The amino acid sequence of the G protein consisted of 669 aa and contained an N terminal signal peptide (1-MYHLIVLLVMLGQRAVA-17), a noncytoplasmic domain (aa 18–646), a transmembrane domain (647-ITAILMPLLSLAVVVGIMCC-667), and a cytoplasmic tail of 2 aa, similar to other rhabdovirus G proteins as predicted by using Phobius and TMHMM (<http://www.cbs.dtu.dk/services/TMHMM>) (6,7). We predicted 3 potential glycosylation sites in the ectodomain at positions 38–40

(NKT), 554–556 (NAS), and 592–594 (NIS) using NetNGlyc 1.0 (<http://www.cbs.dtu.dk/services/NetNGlyc>).

Between the G and L genes, a complex intergenic region was present that contained 3 ORFs of 246 nt (7,413–7,658 aa), 231 nt (7,716–7,946 aa), and 459 nt (7,893–8,355 aa), of which 2 were overlapping frames (U1–3). Additional ORFs between G and L genes were detected previously in other rhabdoviruses (8,9). We detected transmembrane domains in the amino acid sequences of all 3 additional ORFs, suggesting they might act as viroporin (8,9).

The L gene of RFFRV contained 6,591 nt (2,197 aa). We detected several conserved domains and motifs, including RNA-dependent RNA polymerase, mRNA-capping region, mRNA capping enzyme, and virus-capping methyltransferase. Alignment of the deduced amino acid sequence of the L gene with the L gene of various other viruses belonging to the Mononegavirales order by using MAFFT version 7 (<http://mafft.cbrc.jp/alignment/software/>) and subsequent phylogenetic reconstruction by using a maximum-likelihood tree (WAG+F+I+G model with 100 bootstrap replicates in MEGA5 [<http://www.megasoftware.net>]) suggested that this virus belongs to a novel genus of the *Rhabdoviridae* family. In addition, pairwise identities of the deduced amino acid sequence of the L gene of RFFRV with that of other rhabdoviruses of the *Rhabdoviridae* family were only  $\leq 35\%$  (online Technical Appendix Table 2).

Because the fox was found dead and no tissue samples were collected, whether RFFRV played a role in the animal's death is unknown. In addition, multiple attempts to isolate this virus on various cell lines of eukaryotes (Vero E6, MDCK, CRFK, N2a, and BHK cells, primary fox kidney cells) failed because of the absence of cytopathic effects and viral replication by quantitative reverse transcription PCR,




Figure. Genome organization and phylogenetic analysis of RFFRV. A) Genome organization of RFFRV. Indicated are the locations of the major ORFs (including the positions of the first and last nucleotide) and 3 theoretical minor ORFs between the G and L genes. B) Phylogenetic maximum-likelihood tree using the WAG+F+I+G model and 100 bootstrap replicates in MEGA5 (<http://www.megasoftware.net>) of the deduced amino acid sequence of the L genes of various viruses of the order Mononegavirales. G, glycoprotein; L, RNA-dependent RNA polymerase; M, matrix; N, nucleoprotein; ORF, open reading frame; P, phosphoprotein; RFFRV, red fox fecal rhabdovirus. Only bootstrap values in the close proximity of the branch of the RFFRV are indicated. Scale bar indicates nucleotide substitutions per site. Viruses and GenBank accession numbers are shown in the expanded figure legend online (<http://wwwnc.cdc.gov/EID/article/20/12/14-0236-F1.htm>).

despite a high number of reads in the original sample. The fox might have acquired the virus through spillover from a small prey, such as a bat, and additional studies are required to elucidate the prevalence, original host, and pathogenic potential of this novel virus.

#### Acknowledgments

We thank all researchers and institutions for their invaluable help during sampling and for providing the specimens used in this study, especially Patricia Lizarraga, Laura Elorza, Ricardo Gutierrez, and Luis Javier Chueca.

This work was funded by the European Community's Seventh Framework Program (FP7/2007–2013) under the project "European Management Platform for Emerging and Re-emerging Infectious

Disease Entities" European Community grant agreement no. 223498; the Virgo Consortium; and the Niedersachsen-Research Network on Neuroinfectiology of the Ministry of Science and Culture of Lower Saxony, Germany. In addition, this research was funded partially by the Basque Government through the research group on "Systematics, Biogeography and Population Dynamics" (ref. IT317-10; GIC10/76).

A.R.-G. holds a postdoctoral fellowship awarded by the Department of Education, Universities and Research of the Basque Government (ref. DKR-2012-64) and was awarded a short-visit research grant from the ConGenOmics Research networking program of the European Science Foundation to visit the Department of Viroscience, Erasmus Medical Centre and develop the current research project.

**Rogier Bodewes,<sup>1</sup>  
Aritz Ruiz-Gonzalez,<sup>1</sup>  
Anita C. Schürch,  
Albert D.M.E. Osterhaus,  
and Saskia L. Smits**

Author affiliations: Erasmus Medical Centre, Rotterdam, the Netherlands (R. Bodewes, A.C. Schürch, A.D.M.E. Osterhaus, S.L. Smits); University of the Basque Country, Vitoria-Gasteiz, Spain (A. Ruiz-Gonzalez); National Institute for Environmental Protection and Research, Ozzano dell'Emilia, Italy (A. Ruiz-Gonzalez); and Viroclinics Biosciences, Rotterdam (A.D.M.E. Osterhaus, S.L. Smits)

DOI: <http://dx.doi.org/10.3201/eid2012.140236>

## References

1. International Committee on Taxonomy of Viruses. Virus taxonomy: 2013 release [cited 2014 Feb 9]. <http://www.ictvonline.org/virusTaxonomy.asp>
2. Matha IS, Salunke SR. Immunogenicity of purified Vero cell rabies vaccine used in the treatment of fox-bite victims in India. *Clin Infect Dis*. 2005;40:611–3. <http://dx.doi.org/10.1086/427700>
3. Bodewes R, Ruiz-Gonzalez A, Schapendonk CM, van den Brand JM, Osterhaus AD, Smits SL. Viral metagenomic analysis of feces of wild small carnivores. *Virology*. 2014;11:89. <http://dx.doi.org/10.1186/1743-422X-11-89>
4. Palacios G, Forrester NL, Savji N, Travassos da Rosa AP, Guzman H, Detoy K, et al. Characterization of Farmington virus, a novel virus from birds that is distantly related to members of the family *Rhabdoviridae*. *Virology*. 2013;10:219. <http://dx.doi.org/10.1186/1743-422X-10-219>
5. Albertini AA, Ruigrok RW, Blondel D. Rabies virus transcription and replication. *Adv Virus Res*. 2011;79:1–22. <http://dx.doi.org/10.1016/B978-0-12-387040-7.00001-9>
6. Coll JM. The glycoprotein G of rhabdoviruses. *Arch Virol*. 1995;140:827–51. <http://dx.doi.org/10.1007/BF01314961>
7. Käll L, Krogh A, Sonnhammer EL. A combined transmembrane topology and signal peptide prediction method. *J Mol Biol*. 2004;338:1027–36. <http://dx.doi.org/10.1016/j.jmb.2004.03.016>
8. Gubala AJ, Proll DF, Barnard RT, Cowled CJ, Crameri SG, Hyatt AD, et al. Genomic characterisation of Wongabel virus reveals novel genes within the *Rhabdoviridae*. *Virology*. 2008;376:13–23. <http://dx.doi.org/10.1016/j.virol.2008.03.004>
9. McWilliam SM, Kongsuwan K, Cowley JA, Byrne KA, Walker PJ. Genome organization and transcription strategy in the complex GNS-L intergenic region of bovine ephemeral fever rhabdovirus. *J Gen Virol*. 1997;78:1309–17.

Address for correspondence: Rogier Bodewes, Erasmus Medical Centre, Department of Viroscience Dr. Molewaterplein 50, 3015GE Rotterdam, the Netherlands; email: [r.bodewes@erasmusmc.nl](mailto:r.bodewes@erasmusmc.nl)

## Ngari Virus in Goats during Rift Valley Fever Outbreak, Mauritania, 2010

**To the Editor:** Ngari virus (NRIV) is a single-stranded RNA virus belonging to the family *Bunyaviridae*, genus *Orthobunyavirus*. The genome comprises 3 segments, the small (S), medium (M), and large (L) segments, which encode the nucleocapsid (N) protein, the 2 glycoproteins Gn and Gc, and the RNA-dependent RNA-polymerase, respectively. Sequence analysis showed that NRIV is a reassortant between Bunyamwera virus (BUNV) and Batai virus (BATV), both from the genus *Orthobunyavirus*. S and L segments derived from BUNV, and the M segment derived from BATV (1,2). NRIV is more virulent than BUNV and BATV and is associated with hemorrhagic fever. NRIV was first isolated from *Aedes simpsoni* mosquitoes in 1979 and from humans in 1993, both in Senegal (3). During 1997 and 1998, humans were affected with hemorrhagic fever diseases in Kenya and Somalia that were caused by Rift Valley fever virus (RVFV) and by NRIV (2,4).

In 2010, during an ongoing RVFV outbreak in Mauritania, we collected 163 serum samples (62 from camels, 8 from cattle, and 93 from small ruminants) (5). RVFV RNA was isolated from serum samples as described previously (5). Further molecular testing of the samples was conducted by a SYBRGreen-based real-time reverse transcription PCR (RT-PCR) adapted from a conventional RT-PCR and based on generic primers (bun\_group\_forw 5'-CTGCTAACACCAGCAGTACTTTTGAC-3' and bun\_group\_rev 5'-TGGAGGGTAAGACCATCGTCAGGAAGT-3') that target a 250-nt sequence of the S segment of Bunyamwera serogroup members (6). Real-time RT-PCR was performed in a CFX 96 real-time PCR system (Bio-Rad, Hercules, CA, USA) by using 5  $\mu$ L RNA with a QuantiTect SYBR Green RT-PCR Kit (QIAGEN, Hilden Germany) in a final volume of 25  $\mu$ L. Cycling conditions included RT at 50°C for 30 min and 95°C for 15 min, followed by amplification with 44 cycles of 95°C for 15 s, 55°C for 25 s, 72°C for 30 s, and 77°C for 5 s. A melting curve analysis was then performed starting with 95°C for 60 s, and a temperature gradient was conducted from 68°C to 94°C in increments of 0.2°C.

Of the 163 serum samples tested, 2 samples from goats resulted in a positive signal with cycle thresholds of 23 (sample 51) and 28 (sample 65), respectively. Both samples showed similar melting peaks at  $\approx$ 78.2°C and shared the identical partial nucleotide sequence of the S segment. The sequence belongs to the Bunyamwera serogroup, but the short partial sequence was not sufficient for accurate virus determination and identification. For this reason, both serum samples were used to inoculate cell monolayers of Vero E6 cells that were assayed for virus replication. Only sample 51 displayed a cytopathic effect after 72 h and was further analyzed. We isolated the viral RNA from cell culture with TRIzol reagent

<sup>1</sup>These authors contributed equally to this article.