

Zoonotic *Bartonella* Species in Cardiac Valves of Healthy Coyotes, California, USA

Spencer P. Kehoe, Bruno B. Chomel, Matthew J. Stuckey, Rickie W. Kasten, Nandhakumar Balakrishnan, Benjamin N. Sacks, and Edward B. Breitschwerdt

We investigated whether *Bartonella* spp. could cause endocarditis in coyotes or localize to cardiac valves before lesions develop. *Bartonella* DNA was amplified more often from coyote cardiac valves than spleen. *Bartonella* infection apparently leads to cardiac valve tropism, which could cause endocarditis, an often lethal complication in mammals, including humans.

Bartonellae are vector-borne gram-negative, aerobic, intracellular bacteria with a tropism for erythrocytes and endothelial cells (1). These bacteria, many of which are zoonotic, infect a wide range of domestic and wild animal species, causing a spectrum of disease manifestations and pathologies (2). Bartonellae, especially *Bartonella vinsonii* subsp. *berkhoffii* (*B. v. berkhoffii*), cause valvular endocarditis, especially of the aortic valve in mammals, including humans, dogs, cats, and cattle (1,3). Fleas and possibly ticks can vector *B. v. berkhoffii* (4). *Bartonella* species, typically observed in 5- to 7-year-old mid-sized to large dogs, account for ≈28% of endocarditis in dogs (3,5). Bartonellae, including *B. v. berkhoffii*, account for ≈3% of human endocarditis cases (1,6). In dogs and humans, these bacteria appear to have a specific tropism for aortic and mitral valves (1). Similar to lesions that develop with *Coxiella burnetii* endocarditis (7), valvular vegetative lesions can result from chronic *Bartonella* infection.

In California, coyotes (*Canis latrans*) are a major reservoir for *B. v. berkhoffii* (8). Natural *Bartonella* reservoir hosts are often asymptomatic, but to our knowledge, the possible role of *Bartonella*-induced endocarditis in

coyotes has never been investigated. We hypothesized that *B. v. berkhoffii* or other *Bartonella* species could cause endocarditis in coyotes. We also hypothesized that bartonellae might preferentially localize to the aortic and/or mitral valves before vegetative lesions develop. Hence, coyotes served as a naturally occurring epidemiologic and physiologic sentinel model for studying infection kinetics and pathology induced by this bacterium in a reservoir host (coyotes).

The Study

During the early 2000s, a total of 70 coyotes trapped in 9 northern California counties as part of a depredation control program were assessed for zoonotic heartworm (*Dirofilaria immitis*) disease (9). Coyote hearts and spleens were collected at that time and stored at -70°C in a manner to avoid DNA carryover during handling, storage, and processing. In 2012 and 2013, the hearts were dissected for macroscopic evidence of aortic and mitral valve vegetative endocarditis lesions. A board-certified veterinary pathologist examined possible valvular lesions or thickened regions; however, because the samples had been frozen, microscopic histopathologic examination was not conducted. We extracted DNA from aortic and mitral valves and spleens using DNeasy Blood and Tissue Kits (QIAGEN, Hilden, Germany). *B. v. berkhoffii*-spiked rabbit blood was the DNA extraction positive control. We tested extracted samples by PCR for *Bartonella* DNA targeting the citrate synthase gene (*gltA*) (10). PCR of spleen tissue was a substitute for blood culture detection of bacteremia. *B. henselae* DNA and distilled water were PCR-positive and -negative controls, respectively. Partial gene sequencing was performed on PCR-positive tissues. Nineteen aortic valve, mitral valve, and splenic DNA samples from 14 coyotes (*B. v. berkhoffii* PCR-positive animals by *gltA* PCR and sequencing) were genotyped by using primers targeting 16–23S intergenic transcribed spacer (ITS) region, as previously described (11) with minor modifications in annealing temperature (68°C for 15 s) and extension (72°C for 18 s). We conducted statistical analysis for differences in tissue tropism using Epi Info version 6 (Centers for Disease Control and Prevention, Atlanta, GA, USA).

Of the 70 coyotes collected from 9 counties (Figure), 45 (64%) were male. Coyotes' ages ranged from ≤ 1 year (57 [81%]) to >5 years (3 [4%]). Nine (20%) male and 6 (24%) female coyotes were PCR positive for *Bartonella* species. Fourteen (93%) of the 15 *Bartonella*-positive coyotes were ≤ 3 years old, of which 13 (87%) were ≤ 1 year old. Prevalence by county ranged from 0% to 33% (Figure).

We found no gross vegetative aortic or mitral valvular endocarditis lesions. Fifteen (21%) coyotes tested

Author affiliations: University of California, Davis, Davis, California, USA (S.P. Kehoe, B.B. Chomel, M.J. Stuckey, R.W. Kasten, B.N.Sacks); and North Carolina State University, Raleigh, North Carolina, USA (N. Balakrishnan, E.B. Breitschwerdt)

DOI: <http://dx.doi.org/10.3201/eid2012.140578>




Figure. Molecular prevalence of *Bartonella* species in 70 coyotes from 9 counties, California, USA. Shaded areas are counties where coyotes were trapped during the early 2000s. *Bartonella*-positive coyotes were identified from the 9 counties as follows: Yuba, 6 (33%) of 18 trapped coyotes; Santa Clara, 3/22 (14%); Mendocino, 2/11 (18%); Napa, 2/6 (33%); Sonoma, 1/5 (20%); Glenn, 1/4 (25%); Yolo, 0/1; Butte, 0/1; Solano, 0/2.

positive by PCR for *Bartonella gltA* gene. Overall, 8 aortic valves, 5 mitral valves, and 4 spleens were PCR positive. Aortic and mitral cardiac valves of 1 coyote (no. 93) tested positive by PCR for *B. v. berkhoffii*, and the aortic valve and spleen of another coyote (no. 110) were PCR positive (Table). Although a higher percentage of positive cardiac valves were aortic (53%) than mitral (33%), the difference was not significant. However, when we compared the number of *Bartonella*-infected cardiac valves (11 valves) with *Bartonella*-infected spleens (3 spleens), we found that *Bartonella* DNA was amplified 4.16 times (95% CI 1.02–24.12) more often from cardiac valves than from spleens.

Partial DNA sequencing showed that aortic valves from 8 (53%) of 15 coyotes were *B. v. berkhoffii* positive, compared with mitral valves from 4 (27%) and spleens from 3 (20%) coyotes. *B. rochalimae* was amplified from the spleen of coyote no. 99, and *B. henselae* DNA was

amplified from the mitral valve of coyote no. 137 (Table). Of 14 coyotes tested for *B. v. berkhoffii* genotypes by 16–23S ITS PCR, 8 were positive, whereas *Bartonella* DNA was not amplified from the remaining 6 tissue DNA samples by using ITS primers. By sequence analyses, 4 coyotes were infected with *B. v. berkhoffii* genotype I, 3 with genotype II, and 1 with genotype III.

Conclusions

Our study documents the presence of 3 zoonotic *Bartonella* species in heart valves and/or spleen of free-ranging coyotes from northern California. Despite the absence of gross vegetative endocardial lesions, *Bartonella* DNA was amplified and sequenced from >20% of the coyotes, mainly from cardiac valves; only 4 (6%) coyotes had PCR-positive spleens, compared with 12 (17%) coyotes with PCR-positive cardiac valves. We hypothesize that *Bartonella* in the spleen indicated early or ongoing bacteremia, whereas

Table. Coyotes (*Canis latrans*) positive for *Bartonella* species by PCR, California, USA

Coyote no.	Sex/estimated		County	<i>Bartonella</i> PCR-positive tissue	<i>Bartonella</i> species by DNA sequencing
	age, y	Weight, kg			
91	F/1	11.7	Yuba	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type III
92	M/1	13	Yuba	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type I
93	M/1	10.5	Mendocino	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type I
				Mitral valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> *
99	M/<1	10.6	Yuba	Spleen	<i>B. rochalimae</i>
101	M/1	12.3	Yuba	Mitral valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> *
102	M/<1	10	Glenn	Spleen	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type II
106	F/1	12	Yuba	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> *
110	M/<1	11	Yuba	Aortic valve, spleen	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type II
121	F/<1	8.6	Santa Clara	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> *
124	F/9	10.4	Santa Clara	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> *
137	M/<1	10.7	Mendocino	Mitral valve	<i>B. henselae</i>
146	M/3	16.6	Sonoma	Aortic valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type I
152	M/<1	9.7	Napa	Spleen	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type II
156	F/<1	8.1	Santa Clara	Mitral valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> *
164	F/<1	10.1	Napa	Mitral valve	<i>B. vinsonii</i> subsp. <i>berkhoffii</i> type I

*Type not amplified.

bartonellae in the heart valves, in their absence in the spleen, indicated valvular bacterial localization, possibly facilitating persistent infection that could evolve through time to endocarditis. This evolution has been observed for *C. burnetii* infection in humans (12), for which the mean reported interval from illness onset to endocarditis diagnosis is 12–24 months (7). *Bartonella* endocarditis is usually seen in middle-aged dogs (mean age 6.3 years \pm 2.8) (3,5) and in adult humans (mean age 54 years \pm 16) (6). Because 93% of the PCR-positive coyotes were \leq 3 years old, they were very likely too young for vegetative endocarditis to have developed.

Nevertheless, the fact that \approx 20% of the cardiac valve tissues were PCR positive for *Bartonella* perhaps indicates that the bacteria had localized to the valves of infected coyotes. *B. v. berkhoffii* can induce vasoproliferative lesions in animals (13); thus, cases of *Bartonella* endocarditis might represent only a small fraction of infected animals that have chronic cardiac valvular localization. All 3 *B. v. berkhoffii* genotypes identified in these coyotes have been previously involved in humans or dogs with endocarditis (14,15). To our knowledge, *B. henselae* and *B. v. berkhoffii* genotype III have not been previously identified in coyotes; thus, these mammals can be added to the list of susceptible species. Coyotes might be a natural reservoir for *B. v. berkhoffii* genotype III, which so far has been mainly described in California gray foxes (14).

In conclusion, *Bartonella* infection of a natural reservoir appears to lead to cardiac valve tropism. This tropism could result in development of endocarditis, a severe and often lethal complication of *Bartonella* infection.

Mr Kehoe is a fourth-year veterinary student at the University of California, Davis. His primary research interests include zoologic and wildlife medicine and health and emerging zoonotic infections.

References

- Chomel BB, Kasten RW, Williams C, Wey AC, Henn JB, Maggi R, et al. *Bartonella* endocarditis: a pathology shared by animal reservoirs and patients. *Ann N Y Acad Sci.* 2009;1166:120–6. <http://dx.doi.org/10.1111/j.1749-6632.2009.04523.x>
- Boulouis HJ, Chang CC, Henn JB, Kasten RW, Chomel BB. Factors associated with the rapid emergence of zoonotic *Bartonella* infections. *Vet Res.* 2005;36:383–410. <http://dx.doi.org/10.1051/vetres:2005009>
- MacDonald KA, Chomel BB, Kittleson MD, Kasten RW, Thomas WP, Pesavento P. A prospective study of canine infective endocarditis in northern California (1999–2001): emergence of *Bartonella* as a prevalent etiologic agent. *J Vet Intern Med.* 2004;18:56–64.
- Yore K, Digangi B, Brewer M, Balakrishnan N, Breitschwerdt EB, Lappin M. Flea species infesting dogs in Florida and *Bartonella* spp. prevalence rates. *Vet Parasitol.* 2014;199:225–9. <http://dx.doi.org/10.1016/j.vetpar.2013.10.017>
- Sykes JE, Kittleson MD, Pesavento PA, Byrne BA, MacDonald KA, Chomel BB. Evaluation of the relationship between causative organisms and clinical characteristics of infective endocarditis in dogs: 71 cases (1992–2005). *J Am Vet Med Assoc.* 2006;228:1723–34. <http://dx.doi.org/10.2460/javma.228.11.1723>
- Houpikian P, Raoult D. Diagnostic methods. Current best practices and guidelines for identification of difficult-to-culture pathogens in infective endocarditis. *Cardiol Clin.* 2003;21:207–17. [http://dx.doi.org/10.1016/S0733-8651\(03\)00028-6](http://dx.doi.org/10.1016/S0733-8651(03)00028-6)
- Stein A, Raoult D. Q fever endocarditis. *Eur Heart J.* 1995;16 Suppl B:19.
- Chang CC, Kasten RW, Chomel BB, Simpson DC, Hew CM, Kordick DL, et al. Coyotes (*Canis latrans*) as the reservoir for a human pathogenic *Bartonella* sp.: molecular epidemiology of *Bartonella vinsonii* subsp. *berkhoffii* infection in coyotes from central coastal California. *J Clin Microbiol.* 2000;38:4193–200.
- Sacks BN, Chomel BB, Kasten RW, Chang CC, Sanders RK, Leterme SD. Validation for use with coyotes (*Canis latrans*) of a commercially available enzyme-linked immunosorbent assay for *Dirofilaria immitis*. *Vet Parasitol.* 2002;109:45–58. [http://dx.doi.org/10.1016/S0304-4017\(02\)00254-6](http://dx.doi.org/10.1016/S0304-4017(02)00254-6)
- Norman AF, Regnery R, Jameson P, Greene C, Krause DC. Differentiation of *Bartonella*-like isolates at the species level by PCR-restriction fragment length polymorphism in the citrate synthase gene. *J Clin Microbiol.* 1995;33:1797–803.

11. Varanat M, Maggi RG, Linder KE, Breitschwerdt EB. Molecular prevalence of *Bartonella*, *Babesia*, and hemotropic *Mycoplasma* sp. in dogs with splenic disease. *J Vet Intern Med*. 2011;25:1284–91. <http://dx.doi.org/10.1111/j.1939-1676.2011.00811.x>
12. Fenollar F, Fournier PE, Carrieri MP, Habib G, Messana T, Raoult D. Risks factors and prevention of Q fever endocarditis. *Clin Infect Dis*. 2001;33:312–6. <http://dx.doi.org/10.1086/321889>
13. Beerlage C, Varanat M, Linder K, Maggi RG, Cooley J, Kempf VA, et al. *Bartonella vinsonii* subsp. *berkhoffii* and *Bartonella henselae* as potential causes of proliferative vascular diseases in animals. *Med Microbiol Immunol (Berl)*. 2012;201:319–26. <http://dx.doi.org/10.1007/s00430-012-0234-5>
14. Maggi RG, Chomel B, Hegarty BC, Henn J, Breitschwerdt EB. A *Bartonella vinsonii berkhoffii* typing scheme based upon 16S–23S ITS and Pap31 sequences from dog, coyote, gray fox, and human isolates. *Mol Cell Probes*. 2006;20:128–34. <http://dx.doi.org/10.1016/j.mcp.2005.11.002>
15. Cadenas MB, Bradley J, Maggi RG, Takara M, Hegarty BC, Breitschwerdt EB. Molecular characterization of *Bartonella vinsonii* subsp. *berkhoffii* genotype III. *J Clin Microbiol*. 2008;46:1858–60. <http://dx.doi.org/10.1128/JCM.02456-07>

Address for correspondence: Bruno B. Chomel, Department of Population Health and Reproduction, School of Veterinary Medicine, VM3B, 1089 Veterinary Medicine Dr, University of California, Davis, Davis, CA 95616, USA; email: bbchomel@ucdavis.edu

