

Novel Avian Influenza A(H7N9) Virus in Tree Sparrow, Shanghai, China, 2013

Baihui Zhao,¹ Xi Zhang,¹ Wenfei Zhu,¹ Zheng Teng, Xuelian Yu, Ye Gao, Di Wu, Enle Pei, Zhengan Yuan, Lei Yang, Dayan Wang, Yuelong Shu, and Fan Wu

In spring 2013, influenza A(H7N9) virus was isolated from an apparently healthy tree sparrow in Chongming Dongping National Forest Park, Shanghai City, China. The entire gene constellation of the virus is similar to that of isolates from humans, highlighting the need to monitor influenza A(H7N9) viruses in different species.

Since its emergence in China in February 2013, avian influenza A(H7N9) virus has resulted in 217 human infections and 57 deaths (1). The biological features of the virus and its pandemic potential have caused global concern (2). Although the epidemic declined quickly after the closure of live poultry markets in China in April 2013, new cases in humans have reemerged since October 2013. The number of new cases has increased sharply since January 1, 2014, paralleling the peak of the first wave (1,3,4), indicating that subtype H7N9 viruses were circulating asymptotically among natural hosts. Sequence data indicated that the hemagglutinin gene of this novel subtype H7N9 virus might originate from a subtype H7N3 virus in ducks and that the neuraminidase gene probably originated from a subtype H7N9 virus in wild birds (5) or ducks or chickens (6,7). These data suggest that wild birds might play a role in the emergence of subtype H7N9 viruses, similar to the role they played in the geographic spread of avian subtype H5N1 viruses (8). However, although avian influenza A(H7N9) viruses have been isolated from chickens and pigeons, to our knowledge, none have been isolated from wild birds. To better understand the role of wild birds in the emergence and potential dissemination of subtype H7N9

viruses, during spring of 2013, the Shanghai Municipal Center for Disease Control and Prevention, in collaboration with the Shanghai Wildlife Conservation and Management Center, investigated influenza A(H7N9) virus infection among wild birds in Shanghai.

The Study

During April 10–May 15, a total of 2,198 fecal, tissue, cloacal swab, and tracheal swab samples were collected from wild birds in Shanghai. Trained staff captured healthy birds with an approved trapping method, collected samples, and released the birds. Tissue samples were collected from naturally dead wild birds. Information on bird species and sampling places are listed in online Technical Appendix Tables 1 and 2 (wwwnc.cdc.gov/EID/article/20/5/13-1707-Techapp1.pdf). RNA was extracted from each sample and tested by using influenza A universal real-time PCR according to the standard operating procedure of the World Health Organization (9). Influenza A virus–positive specimens were further subtyped by reverse transcription PCR with an avian influenza A virus subtype primer set reported previously (10). Of the 2,198 samples, 28 were positive for influenza A virus. One tracheal sample from an apparently healthy tree sparrow was positive for the novel subtype H7N9 virus, whereas the cloacal swab samples from this bird were negative. The positive sample was inoculated into 11-day-old specific pathogen free embryonated chicken eggs for virus isolation. The isolated virus was termed A/tree sparrow/Shanghai/01/2013 (H7N9). The tree sparrow had been collected from a forest on Chongming Dongping Forest Park, which is 47 km from Dongtan National Nature Reserve, a winter habitat for wild migratory birds (Figure 1).

To explore the genetic relationships between this sparrow-derived influenza A(H7N9) virus and other viruses from humans and poultry, we amplified total genomic segments by using viral RNA directly isolated from the original specimen with the primer sets listed in online Technical Appendix Table 3 and sequenced by Sunny Biotech Co., Ltd. (Shanghai, China). The Chinese National Influenza Center performed the sequencing by using RNA from chicken embryonated cultured viruses in an ABI 3730xl automatic DNA analyzer (Life Technologies, Foster City, CA, USA). Full-genome sequences from the original sample and the embryonated chicken egg isolation were deposited in GenBank (accession nos. KF609524–KF609531 and KJ508887–KJ508894). To facilitate the phylogenetic analysis, we downloaded sequences of the novel subtype H7N9 viruses from 2013 and the avian subtype H7N9 viruses from before 2013 from the Global Initiative on Sharing Avian Influenza Data (<http://platform.gisaid.org/>)

Author affiliations: Shanghai Municipal Center for Disease Control and Prevention, Shanghai, China (B. Zhao, X. Zhang, Z. Teng, X. Yu, Y. Gao, Z. Yuan, F. Wu); National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China (W. Zhu, L. Yang, D. Wang, Y. Shu); and Shanghai Wildlife Conservation and Management Center, Shanghai (D. Wu, E. Pei)

DOI: <http://dx.doi.org/10.3201/eid2005.131707>

¹These authors contributed equally to this article.



Figure 1. Location of tree sparrow from which novel avian influenza A(H7N9) virus was isolated: Chongming National Dongping Forest Park of Shanghai (yellow solid circle), which is located in the Australia–East Asia migratory wild bird flyway. Top right: sampling locations in Shanghai City. Bottom right: sampling location of influenza A(H7N9)-positive tree sparrow. CM, Chongming District; BS, Baoshan District; JD, Jiading District; SH center, Changning, Putuo, and Xuhui Districts; QP, Qingpu District; MH, Minhang District; SJ, Songjiang District; PD, Pudong District; FX, Fengxian District; JS, Jinshan District.

epi3/frontend#46b284). Sequence alignments were performed by using the MegAlign method of Lageregene 7.01 software (www.dnastar.com/t-megalign.aspx). Phylogenetic analysis was analyzed by using the neighbor-joining method in MEGA software version 5.10 (www.megasoftware.net).

Eight gene segments of the tree sparrow virus shared most ($\geq 99.3\%$) similarities with subtype H7N9 virus isolates from humans. Phylogenetic analysis of hemagglutinin genes revealed that subtype H7N9 viruses could be classified into Eurasia and North America lineages. The subtype H7N9 virus in this study shared the same influenza lineage with all novel subtype H7N9 viruses from humans and poultry (Figure 2). Like the hemagglutinin genes, the other 7 genes showed the same evolutionary pattern (online Technical Appendix Figure 1). Homology and phylogenetic analyses indicated that the genetic constellation of the tree sparrow-derived subtype H7N9 virus is similar to that of novel subtype H7N9 avian influenza viruses isolated from humans and poultry in this region.

According to genetic signatures, the tree sparrow-derived subtype H7N9 virus acquired the ability to bind to human-like receptors, for which substitutions G186V and Q226L in hemagglutinin protein (H3) are responsible, similar to most human and avian subtype H7N9 viruses. A 69–73-aa deletion was also found in its neuraminidase gene (N2). Amino acid 292R was maintained in neuraminidase

genes, indicating its sensitivity to neuraminidase inhibitors. However, an S31N mutation in the matrix 2 protein confers resistance to adamantane. Asp at polymerase basic 2 protein (PB2) residue 701 was associated with reduced transmissibility. Mixed E/K at residue 627 in PB2 and V/I at residue 31 in matrix 1 protein were detected from the original sample (online Technical Appendix Figure 2). Because all previously reported influenza A (H7N9) viruses isolated from birds or the environment acquired PB2 627E, the mixed amino acids of 627E/K in the sample from the tree sparrow suggested that the PB2 E627K substitution might have occurred during replication of the virus in birds.

Conclusions

The high similarity of genes from the avian influenza A(H7N9) virus from an apparently healthy tree sparrow in Shanghai and influenza A(H7N9) viruses from humans and poultry in this region indicate that avian influenza A(H7N9) virus might be transmitted from poultry to tree sparrows or vice versa. Earlier reports documented that influenza A viruses, including subtypes H5N1 and H3N2, have been isolated from sparrows (11,12). A serologic survey also suggested that rates of influenza A virus infection were high among sparrows (13), which might result from abundant distribution of avian influenza virus receptor SA α 2,3Gal in the respiratory tracts of sparrows (14). The novel subtype




Figure 2. Phylogenetic tree of the hemagglutinin (A) and neuraminidase (B) genes of influenza A(H7N9) viruses. Multiple alignments were constructed by using the MUSCLE algorithm of MEGA software version 5.10 (www.megasoftware.net). Phylogenetic trees were constructed by using the neighbor-joining method with bootstrap analyses of 1,000 replications. Bootstrap values >60% are shown in the nodes. Sequences of human influenza A(H7N9) viruses are shown in purple, novel subtype H7N9 viruses from poultry (chickens, ducks, and pigeons) in blue, novel subtype H7N9 viruses from the environment in green, and novel subtype H7N9 viruses from wild birds in red. Scale bar indicates base substitutions per site.

H7N9 virus expands not only the number of influenza virus subtypes that infect tree sparrows but also range of hosts for subtype H7N9 viruses. Our finding of only 1 subtype H7N9-positive sample among 2,198 samples is consistent with recent findings that subtype H7N9 in wild birds is rare (15).

Tree sparrows are abundant and widely distributed in China. They are frequently in contact with humans and poultry. Prevalence of avian influenza viruses among tree sparrows could increase opportunities for them to carry influenza viruses from aquatic birds to domestic farms and even to humans. Hence, such expansion of influenza A(H7N9) virus host ranges undoubtedly increased the seriousness of the threat of this novel subtype.

Tree sparrows have been shown to be susceptible to influenza A(H5N1) viruses, and they might have the ability to disseminate subtype H5N1 viruses (11). Dongping National Forest Park, where the novel subtype H7N9-positive tree sparrow was captured, is on Chongming Island, China's third largest island, which is located in the Australia-East Asia migratory wild bird flyway. Dongping National Forest Park is adjacent to Dongtan National Nature Reserve, where hundreds of species of migratory

and domestic birds gather for winter. Whether migratory birds became infected through contact with tree sparrows and then disseminated subtype H7N9 virus to other geographic regions merits further investigation. Isolation of novel influenza A(H7N9) virus in a tree sparrow emphasizes the need to expand influenza surveillance to not only domestic birds but also wild and terrestrial birds.

Acknowledgments

We thank the data submitters from the Global Initiative on Sharing Avian Influenza Data for their H7N9 sequences and Ren Lili for her assistance with the sequencing.

This study was supported by Shanghai Municipal Commission of Health and Family Planning (nos. 2013QLG001 and 2013QLG003), Ministry of Science and Technology of People's Republic of China (no. KJYJ-2013-01-01), and Science and Technology Committee of Shanghai Municipality (no. 10XD1403600).

Dr Zhao works in the microbiology laboratory of the Shanghai Municipal Center for Disease Control and Prevention. Her research interests include respiratory viruses and emerging diseases.

References

- World Health Organization. Global alert and response. Avian influenza A(H7N9) virus [cited 2014 Jan 20]. <http://www.who.int/csr/don/archive/year/2014/en/index.html>
- Zhou J, Wang D, Gao R, Zhao B, Song J, Qi X, et al. Biological features of novel avian influenza A (H7N9) virus. *Nature*. 2013;499:500–3. <http://dx.doi.org/10.1038/nature12379>
- World Health Organization. Avian influenza A (H7N9) in China [cited 2014 Jan 22]. <http://www.wpro.who.int/china/en/index.html>
- Center for Infectious Disease Research and Policy. H7N9 cases in China surge past 200 [cited 2014 Jan 21]. <http://www.cidrap.umn.edu/news-perspective/2014/01/h7n9-cases-china-surge-past-200>
- Liu D, Shi W, Shi Y, Wang D, Xiao H, Li W, et al. Origin and diversity of novel avian influenza A H7N9 viruses causing human infection: phylogenetic, structural, and coalescent analyses. *Lancet*. 2013;381:1926–32. [http://dx.doi.org/10.1016/S0140-6736\(13\)60938-1](http://dx.doi.org/10.1016/S0140-6736(13)60938-1)
- Lam TT, Wang J, Shen Y, Zhou B, Duan L, Cheung CL, et al. The genesis and source of the H7N9 influenza viruses causing human infections in China. *Nature*. 2013;502:241–4. <http://dx.doi.org/10.1038/nature12515>
- Lee RT, Gunalan V, Van TD, Le LT, Eisenhaber F, Maurer-Stroh S. A new piece in the puzzle of the novel avian-origin influenza A (H7N9) virus. *Biol Direct*. 2013;8:26. <http://dx.doi.org/10.1186/1745-6150-8-26>
- Fang LQ, de Vlas SJ, Liang S, Looman CW, Gong P, Xu B, et al. Environmental factors contributing to the spread of H5N1 avian influenza in mainland China. *PLoS ONE*. 2008;3:e2268. <http://dx.doi.org/10.1371/journal.pone.0002268>
- World Health Organization. Manual for the laboratory diagnosis and virological surveillance of influenza [cited 2014 Jan 22]. http://www.who.int/influenza/gisrs_laboratory/manual_diagnosis_surveillance_influenza/en/index.html
- Tsukamoto K, Javier PC, Shishido M, Noguchi D, Pearce J, Kang HM, et al. SYBR green-based real-time reverse transcription-PCR for typing and subtyping of all hemagglutinin and neuraminidase genes of avian influenza viruses and comparison to standard serological subtyping tests. *J Clin Microbiol*. 2012;50:37–45. <http://dx.doi.org/10.1128/JCM.01195-11>
- Liu Q, Ma J, Kou Z, Pu J, Lei F, Li T, et al. Characterization of a highly pathogenic avian influenza H5N1 clade 2.3.4 virus isolated from a tree sparrow. *Virus Res*. 2010;147:25–9. <http://dx.doi.org/10.1016/j.virusres.2009.09.014>
- Han Y, Hou G, Jiang W, Han C, Liu S, Chen J, et al. A survey of avian influenza in tree sparrows in China in 2011. *PLoS ONE*. 2012;7:e33092. <http://dx.doi.org/10.1371/journal.pone.0033092>
- Arenas A, Carranza J, Perea A, Miranda A, Maldonado A, Hermoso M. Type A influenza viruses in birds in southern Spain: serological survey by enzyme-linked immunosorbent assay and haemagglutination inhibition tests. *Avian Pathol*. 1990;19:539–46. <http://dx.doi.org/10.1080/03079459008418706>
- Gutiérrez RA, Som S, Nicholls JM, Buchy P. Eurasian tree sparrows, risk for H5N1 virus spread and human contamination through Buddhist ritual: an experimental approach. *PLoS ONE*. 2011;6:e28609. <http://dx.doi.org/10.1371/journal.pone.0028609>
- Olson SH, Gilbert M, Cheng MC, Mazet JA, Joly DO. Historical prevalence and distribution of avian influenza virus A(H7N9) among wild birds. *Emerg Infect Dis*. 2013;19:2031–3. <http://dx.doi.org/10.3201/eid1912.130649>

Address for correspondence: Fan Wu, Shanghai Municipal Center for Disease Control and Prevention, No. 1380 West Zhongshan Rd, Changning District, Shanghai 200336, China; email: fwu@scdc.sh.cn



Scan this QR Code with your smartphone and enjoy listening to our podcasts about the latest emerging infectious diseases.

<http://wwwnc.cdc.gov/eid/podcasts.htm>

**EID
PODCAST**

www.cdc.gov/ncidod/EID/podcast/

Novel Avian Influenza A(H7N9) Virus in Tree Sparrow, Shanghai, China, 2013

Technical Appendix

Technical Appendix Table 1. The information of sampling places in Shanghai, 2013

Sources of samples	wild migrate bird				wild domestic bird				bird in zoo		environment	unknown
	ardeidae	wader bird	Wild geese and ducks	aquatic bird	wild chicken	sparrow	predator	owl	parrot	peacock	bird living environment	unknown
Swab	123	216	0	56	2	778	36	4	10	0	0	16
Feces	387	0	45	0	0	38	0	0	9	5	58	3
Tissues	0	0	0	0	2	333	65	4	0	0	0	8
Total	510	216	45	56	4	1149	101	8	19	5	58	27







Technical Appendix Table 2. Geographic distributions of sampling places in this study

Sampling places	wild migrate bird				wild domestic bird				bird in zoo		environment	unknown
	wild geese				wild chicken	sparrow	predator	owl	parrot	peacock	bird living	
	ardeidae	wader bird	and ducks	aquatic bird							environment	unknown
Baoshan district (BS)	50	0	45	0	0	2	1	0	0	5	0	0
Chongming district (CM)	55	216	0	48	0	313	10	0	0	0	34	0
Fengxian district (FX)	0	0	0	0	0	20	2	0	0	0	0	2
Jiading district (JD)	199	0	0	0	0	0	0	0	0	0	1	0
Jinshan district (JS)	52	0	0	0	0	0	2	0	0	0	0	0
Minhang district (MH)	50	0	0	0	0	0	0	0	0	0	0	0
Pudong district (PD)	28	0	0	6	4	709	82	8	0	0	0	20
Qingpu district (QP)	26	0	0	0	0	34	2	0	0	0	11	2
Shanghai center (SH center)	50	0	0	2	0	71	2	0	19	0	12	3
Total	510	216	45	56	4	1149	101	8	19	5	58	27


Technical Appendix Table 3. The primer sets for H7N9 whole genome PCR

Target gene	1st round		Forward primer		Reverse Primer	
	RT-PCR Primers	2nd round PCR Primers	name	Sequence	name	Sequence
HA	1F/1680R	1F/693R	HA1F	ATGAACACTCAAATCCTGGTATTCG	HA7-1680R	TTATATACAAATAGTGCACCGCATGTTTC
		589F/1157R	HA7-589F	ACTGCAGAGCAAACCAAGCTATATG	HA7-693R	TTGTGGTCTCGCTCCTGGACT
		1043F/1218R	HA7-1043F	TCATTGAAAATGGATGGGAAGG	HA7-1218R	GTCTATCAACTCAAATTGTTGGTTGGT
		1043F/1680R	HA7-1680R	TTATATACAAATAGTGCACCGCATGTTTC	HA7-1157R	TGATCAATTGCCGATTGAGTGC
NA	NA1F/1398 R	1F/827R	NA9-1F	ATGAATCAAATCAGAAGATTCTATGC	NA9-827R	GAGCATTCTTCAATATGCTTAGCAG
		790F/1271R	NA9-790F	TCTCTGACTGGAAGTCTAAG	NA9-1271R	ATCAACTCCACATAAAAAACAG
		790F/1398R	NA9-1398R	TTAGAGGAAGTACTCTATTTTAGCCCC		
PA	PA1F/G215 1R	1F/708R	PA-1F	ATGGAAGAYTTTGTGCGAC	PA-708R	GAATCCATCCACATAGGCTCT
		13F/896R	PA-13F	GTGCGACAGTGCTTCAATCCA	PA-896R	CTTCGTGGCTCGGGTCTCA
		688F/1550R	PA688F	AGAGCCTATGTGGATGGATTC	PA-1550R	ACGTCGGTATCATTCTCAA
		1531F/2017R	PA1531F	TTGAGGAATGATACCGACGT	PA-2017R	GCTACTCATTGTTGAGGCGCTTA
		G1465F/G2151R	PAG1465F	TGTAGAACCAAAGAAGGAAGACG	PAG2151R	CTATCTTAGTGCATGTGTGAGGAAGG
PB1	PB1F/2274 R	1F/873R	PB1-1F	ATGGATGTCAATCCGACTTT	PB1-873R	AGTCATCATCTTCTCACAACA
		8F/851R	PB1-8F	TCAATCCGACTTTACTTTTCT	PB1-851R	TTTGCCAATTTAGCTTTCTTCTCA
		852F/1774R	PB1-852F	TGTTGTGAGGAAGATGATGACT	PB1-1774R	AAACCAACAGTCTGCCTTC
		1755F/2274R	PB1-1755F	GAAGGCAGGACTGTTGGTTT	PB1-2274R	CTATTTTTGCCGTCTGAGCT
PB2	PB2F/2280 R	1F/833R	PB2-1F	ATGGAAAGAATAAAAGAAGTAAAGAG	PB2-833R	GCCAACGGGTCTGCTGATACTGT
		802F/1390R	PB2-802F	AGAAGAGCAACAGTATCAGCAGA	PB2-1390R	ATATTCCGATCATCCCCATTACA
		1281F/1796R	PB2-1281F	GCTAAATCCCATGCATCAACTCCT	PB2-1796R	AGAACCCTCACGAACCCACTA
		1686F/2221R	PB2-1686F	TCAATGGTCCCAAGATCCTAC	PB2-2221R	AGTCCCGTTTCCGTTTCATCACCA
		G1968F/G2208R	PB2G1968F	CAACTACAACAAGGCAACCAAGAGG	PB2GG2208R	TTAATTGATGGCCATCCGAATCC
		PB22280R	ATTCGACACTAATTGATGGC			
NP	1F/G1497R	1F/642R	NP-1F	ATGGCGTCTCAAGGCACCAAACGATC	NP-624R	CCTTCTCCATTTTCGCCTCTCCAGA

Target gene	1st round		Forward primer		Reverse Primer	
	RT-PCR Primers	2nd round PCR Primers	name	Sequence	name	Sequence
NS	1F/838R	485F/1081R	NP-485F	GGATGTGCTCTCTGATGCAAGGAT	NP-1081R	TAGTGGATAGCTGTCTCTTGGGA
		926F/1292R	NP-926F	ACAGCCAGGTCTTTAGTCTCATTA	NP-1292R	CCTGTAAATGCTGCCATAATGGTTGC
		G1221F/G1497R	NPG-1221F	CGTTCAACCCACTTTCTCAGTA	NP-G1497R	TCAATTGTCATACTCCTCTGCATTG
		1F/471R	NSF1	ATGGATTCCAATACTGTGTCAAGCT	NS471R	CTACAATTGCTCCTTCTTCCGTAA
		1F/477R	NS447R	AGCTCTAAGTAGTATCAGGGCTT	NS381F	CATCACATTGAAAGCAAATTC
		381F/838R	NS838R	TCATTAATAAGCTGAAACGAG	NS342F	CCTATGCATTAGAATGGACCAAGC
M	F0/982R	342F/791R	NS791R	CTTATCTCTTGCTCCACTTCAAGC		
		1F/772R	MF0	AGCAAAAGCAGGTAGATG	M982R	TTACTTCAGCTCTATGTTGACA
		1F/521R	M1F	ATGAGTCTTCTAACCGAGGTCGAA	M772R	AACGACTAGAGGCTCACTTGAAC
		429F/982R	M521R	GCTAGTACCATTCTATTCTCATGC	M429F	TCTTGGACTAGTATGTGCCACTT
			M982R	TTACTTCAGCTCTATGTTGACA		



Technical Appendix Figure 1. Phylogenetic trees of full-length polymerase basic (PB)2, PB1, polymerase acidic, nucleoprotein, matrix, and nonstructural genes of the tree sparrow–derived influenza A (H7N9) viruses in China, 2013. Sequences of human influenza A(H7N9) viruses are shown in purple, novel subtype H7N9 viruses from poultry (chickens, ducks, and pigeons) in blue, novel subtype H7N9 viruses from the environment in green, and novel subtype H7N9 viruses from wild birds in red. Scale bars indicate base substitutions per site.



Technical Appendix Figure 2. Sequencing maps showing mixed population of 627E/K in polymerase basic 2 proteins from influenza A (H7N9) viruses. A, adenine phosphate (green); C, cytosine (blue); G, guanine (black). The square shows mixed nucleotides AAG (amino acid lysine, K) and GAG (amino acid glutamic acid, E). Red and green arrows indicated bidirectional sequencing.