

susceptibility to SBV infection, but differences in seroprevalence suggest different roles for sympatric ruminants in SBV epidemiology. The role of vector species in the transmission of SBV in alpine ecosystems should be analyzed.

Acknowledgments

We thank Mariano Domingo for assistance during necropsy studies of still-born calves, Joan Planas for technical assistance during livestock sample collection, and the rangers and staff of the Freser-Setcases National Hunting Reserve for their collaboration.

This study was supported by the Departament d'Agricultura, Ramaderia, Pesca, Alimentació i Medi Natural de la Generalitat de Catalunya, and funded by the research project CGL2009-11631 of the Spanish Ministerio de Ciencia e Innovación.

**Xavier Fernández-Aguilar,
Joan Pujols,
Roser Velarde, Rosa Rosell,
Jorge R. López-Olvera,
Ignasi Marco, Marti Pumarola,
Joaquim Segalés,
Santiago Lavín,
and Oscar Cabezón**

Author affiliations: Centre de Recerca en Sanitat Animal, Barcelona, Spain (X. Fernández-Aguilar, J. Pujols, R. Rosell, J. Segalés, O. Cabezón); Universitat Autònoma de Barcelona, Barcelona (X. Fernández-Aguilar, R. Velarde, J.R. López-Olvera, I. Marco, M. Pumarola, J. Segalés, S. Lavín, O. Cabezón); Institut de Recerca i Tecnologia Agroalimentàries, Barcelona (J. Pujols); and Generalitat de Catalunya Departament d'Agricultura, Barcelona (R. Rosell)

DOI: <http://dx.doi.org/10.3201/eid2006.130961>

References

- Hoffmann B, Scheuch M, Hoper D, Jungblut R, Holsteg M, Schirremeier H, et al. Novel orthobunyavirus in cattle, Europe, 2011. *Emerg Infect Dis.* 2012;18:469–72. <http://dx.doi.org/10.3201/eid1803.111905>

- Rasmussen LD, Kristensen B, Kirkeby C, Rasmussen TB, Belsham GJ, Bodker R, et al. Culicoids as vectors of Schmallenberg virus. *Emerg Infect Dis.* 2012;18:1204–6. <http://dx.doi.org/10.3201/eid1807.120385>
- Bilk S, Schulze C, Fischer M, Beer M, Hlinak A, Hoffmann B. Organ distribution of Schmallenberg virus RNA in malformed newborns. *Vet Microbiol.* 2012;159:236–8. <http://dx.doi.org/10.1016/j.vetmic.2012.03.035>
- Loeffen W, Quak S, Boer-Luijze E, Hulst M, Van der Poel W, Bouwstra R, et al. Development of a virus neutralization test to detect antibodies against Schmallenberg virus and serological results in suspect and infected herds. *Acta Vet Scand.* 2012;54:44. <http://dx.doi.org/10.1186/1751-0147-54-44>
- van Maanen C, van der Heijden H, Wellenberg GJ, Witteveen G, Lutikholt S, Bouwstra R, et al. Schmallenberg virus antibodies in bovine and ovine fetuses. *Vet Rec.* 2012;171:299. <http://dx.doi.org/10.1136/vr.101061>
- Maxie MG, Youssef S. Nervous system. In: Maxie MG, editor. Jubb, Kennedy, and Palmer's pathology of domestic animals. Philadelphia: Elsevier Limited; 2007. p. 281–457.
- Linden A, Desmecht D, Volpe R, Wirtgen M, Gregoire F, Pirson J, et al. Epizootic spread of Schmallenberg virus among wild cervids, Belgium, Fall 2011. *Emerg Infect Dis.* 2012;18:2006–8. <http://dx.doi.org/10.3201/eid1812.121067>
- Kaufmann C, Steinmann IC, Hegglin D, Schaffner F, Mathis A. Spatio-temporal occurrence of *Culicoides* biting midges in the climatic regions of Switzerland, along with large scale species identification by MALDI-TOF mass spectrometry. *Parasit Vectors.* 2012;5:246. <http://dx.doi.org/10.1186/1756-3305-5-246>

Address for correspondence: Oscar Cabezón, Servei d'Ecopatologia de Fauna Salvatge, Wildlife Diseases Research Group, Departament de Medicina i Cirurgia Animals, Edifici V. Facultat de Veterinària, Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona), Spain; email: oscar.cabezón@cresa.uab.cat

Novel Henipa-like Virus, Mojiang Paramyxovirus, in Rats, China, 2012

To the Editor: The genus *Henipavirus* (family *Paramyxoviridae*) contains 3 established species (*Hendra virus*, *Nipah virus*, and *Cedar virus*) and 19 newly identified species, including 1 full-length sequenced virus, Bat Paramyxovirus Eidhel/GH-M74a/GHA/2009 (1,2). The zoonotic pathogens *Hendra virus* and *Nipah virus* have been associated with lethal neurologic and respiratory diseases in humans, horses, and pigs (3–5). The known natural reservoirs of henipaviruses are fruit bats (1,3); these viruses have not been reported in other wild animals. We report on a novel henipa-like virus, Mojiang paramyxovirus (MojV), in rats (*Rattus flavipectus*) in China.

In June 2012, in Mojiang Hani Autonomous County, Yunnan Province, China, severe pneumonia without a known cause was diagnosed in 3 persons who had been working in an abandoned mine; all 3 patients died. Half a year later, we investigated the presence of novel zoonotic pathogens in natural hosts in this cave. For the investigation, we collected anal swab samples from 20 bats (*Rhinolophus ferrumequinum*), 9 rats (*R. flavipectus*), and 5 musk shrews (*Crocidura dracula*) from the mine for virome analysis.

All samples were processed by using a virus particle-protected nucleic acid purification method, followed by sequence-independent PCR amplification of extracted RNA and DNA (6). The amplified viral nucleic acid libraries were then sequenced by using an Illumina Genome Analyzer II (Illumina Trading, Beijing, China) for a single read of 81 bp. All raw reads were then aligned to the nonredundant protein database of the National Center for Biotechnology Information (www.ncbi.nlm.nih.gov/RefSeq/) by using BLASTx (<http://blast.ncbi.nlm.nih.gov/>).

Find emerging infectious disease information on

facebook

<http://www.facebook.com>

nih.gov/Blast.cgi) after filtering reads as described (6). The taxonomy of the aligned reads was parsed by using the MEGAN4 MetaGenome Analyzer (7).

On the basis of the nonredundant protein alignment results, we identified 38 sequence reads that were classified as *Henipavirus* spp. However, the sequences shared low nucleotide and amino acid identities with known henipaviruses. The reads were then used for reads-based PCR to identify the partial genome of this virus. The remaining genomic sequences were determined by using genome walking. The 5' and

3' untranslated regions were obtained by nested PCR with combined specific primers and henipavirus-specific degenerate primers as described (8), and the exact sequences of the 5' and 3' genome termini were determined by rapid amplification of cDNA ends.

MojV shares similar features with known henipaviruses. The virus has a genome length of 18,404 nt (submitted to GenBank under accession no. KF278639), and has the characteristic henipavirus gene order: 3'-nucleocapsid (N) protein (539 aa); P/V/W/C proteins (phosphoprotein; 694 aa, 464

aa, 434 aa, 177 aa); matrix protein (340 aa); fusion protein (545 aa); attachment glycoprotein (625 aa); and large (L) protein (2,277 aa)-5' (online Technical Appendix Figure, wwwnc.cdc.gov/EID/article/20/6/13-1022-Techapp1.pdf). The predicted conserved sequences between genes showed features characteristic of henipaviruses (online Technical Appendix Table). The central domain of the N protein contains 3 conserved motifs common in all paramyxoviruses: QXW [I/V] X₃K [A/C] XT, FX₂T[I/L][R/K]Φ[G/A][L/I/V] XT, and FX₄YPX₂ΦSΦAMG, where Φ




Figure. Phylogenetic trees based on the nucleocapsid proteins (A) and large proteins (B) of Mojiang paramyxovirus (MojV) and other previously reported paramyxoviruses. Bold font indicates MojV and *Henipavirus* spp. Scale bars indicate nucleotide substitutions per site.

is an aromatic amino acid (9). In addition, the RNA editing site (AAAAGG) for the processing of V and W proteins conserved in the phosphoprotein gene sequences of Hendra virus and Nipah virus was found, and 6 conserved domains within the L proteins of the order *Mononegavirales* (8) were found in the MojV L protein.

The nucleotide identities of predicted MojV genes exhibited similarity with genes of known henipaviruses: N (53.0%–57.0% identity), phosphoprotein (37.8%–43.0% identity), matrix (59.5%–63.4% identity), fusion (47.5%–51.4% identity), attachment glycoprotein (36.6%–41.8% identity), and L (55.9%–58.6% identity) genes. Using MEGA5 (10), we used the phylogenetic trees based on N and L proteins to describe the evolutionary relationships between MojV and members of the family *Paramyxoviridae* (Figure). MojV clustered with the 4 members of the genus *Henipavirus* and was distant from other clusters. Thus, considering the similar genome features between MojV and other henipaviruses, we confirmed that MojV could be classified as a new species closely related to *Henipavirus* spp.

Specific nested primer sets targeting the L gene of MojV were designed to separately re-evaluate the 34 anal swab samples and some tissue samples. Of 9 anal swab samples from the *R. flavipectus* rats, 3 were positive for MojV, and a tissue sample from 1 of the 3 MojV-positive rats was also MojV positive (tissue was not collected from the other 2 rats). All 20 samples from *R. ferrumequinum* bats and all 5 samples from *C. dracula* musk shrews were MojV negative. The 3 MojV-positive anal swab samples were cultured in Vero E6, Hep2, and BHK21 cells for virus isolation; no cytopathic effects or viral replication was detected after 2 blind subculture passages.

Our study showed the presence of a rodent-origin, henipa-like virus, MojV, in China. *R. flavipectus* rats are the natural reservoir of MojV. This finding and its context indicate that

Henipavirus spp. viruses might infect more mammalian hosts than previously thought and that bats may not be the only hosts of henipaviruses.

This work was supported by a National S&T Major Project (China Mega-Project for Infectious Disease; grant no. 2011ZX10004-001) from the People's Republic of China, and by a Basic Research and Operating Expenses grant (no. 2013IPB301) from the Institute of Pathogen Biology, Chinese Academy of Medical Sciences and Peking Union Medical College.

**Zhiqiang Wu,¹ Li Yang,¹
Fan Yang,¹ Xianwen Ren,¹
Jinyong Jiang, Jie Dong,
Lilian Sun, Yafang Zhu,
Hongning Zhou, and Qi Jin**

Author affiliations: Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China (Z. Wu, L. Yang, F. Yang, X. Ren, J. Dong, L. Sun, Y. Zhu, Q. Jin); Institute of Pathogen Biology, Beijing (Z. Wu, L. Yang, F. Yang, X. Ren, J. Dong, L. Sun, Y. Zhu, Q. Jin); and Yunnan Institute of Parasitic Diseases, Puer, China (J. Jiang, H. Zhou)

DOI: <http://dx.doi.org/10.3201/eid2006.131022>

References

- Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, Binger T, et al. Bats host major mammalian paramyxoviruses. *Nat Commun.* 2012;3:796. <http://dx.doi.org/10.1038/ncomms1796>
- Marsh GA, de Jong C, Barr JA, Tachedjian M, Smith C, Middleton D, et al. Cedar virus: a novel henipavirus isolated from Australian bats. *PLoS Pathog.* 2012;8:e1002836. <http://dx.doi.org/10.1371/journal.ppat.1002836>
- Smith I, Wang LF. Bats and their virome: an important source of emerging viruses capable of infecting humans. *Curr Opin Virol.* 2013;3:84–91. <http://dx.doi.org/10.1016/j.coviro.2012.11.006>
- Mendez DH, Judd J, Speare R. Unexpected result of Hendra virus outbreaks for veterinarians, Queensland, Australia. *Emerg Infect Dis.* 2012;18:83–5. <http://dx.doi.org/10.3201/eid1801.111006>
- Sazzad HM, Hossain MJ, Gurley ES, Ameen KM, Parveen S, Islam MS, et al. Nipah virus infection outbreak with nosocomial and corpse-to-human transmission, Bangladesh. *Emerg Infect Dis.* 2013;19:210–7. <http://dx.doi.org/10.3201/eid1902.120971>
- Wu Z, Ren X, Yang L, Hu Y, Yang J, He G, et al. Virome analysis for identification of novel mammalian viruses in bat species from Chinese provinces. *J Virol.* 2012;86:10999–1012. <http://dx.doi.org/10.1128/JVI.01394-12>
- Huson DH, Mitra S, Ruscheweyh HJ, Weber N, Schuster SC. Integrative analysis of environmental sequences using MEGAN4. *Genome Res.* 2011;21:1552–60. <http://dx.doi.org/10.1101/gr.120618.111>
- Miller PJ, Boyle DB, Eaton BT, Wang LF. Full-length genome sequence of Mossman virus, a novel paramyxovirus isolated from rodents in Australia. *Virology.* 2003;317:330–44. <http://dx.doi.org/10.1016/j.virol.2003.08.013>
- Lau SK, Woo PC, Wong BH, Wong AY, Tsoi HW, Wang M, et al. Identification and complete genome analysis of three novel paramyxoviruses, Tuhoko virus 1, 2 and 3, in fruit bats from China. *Virology.* 2010;404:106–16. <http://dx.doi.org/10.1016/j.virol.2010.03.049>
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 2011;28:2731–9. <http://dx.doi.org/10.1093/molbev/msr121>

Address for correspondence: Qi Jin, No. 6 Rongjing East St, Yizhuang, Beijing, 100176, China; email: zdsys@vip.sina.com

The Public Health Image Library (PHIL)



The Public Health Image Library (PHIL), Centers for Disease Control and Prevention, contains thousands of public health-related images, including high-resolution (print quality) photographs, illustrations, and videos.

PHIL collections illustrate current events and articles, supply visual content for health promotion brochures, document the effects of disease, and enhance instructional media.

PHIL Images, accessible to PC and Macintosh users, are in the public domain and available without charge.

Visit PHIL at <http://phil.cdc.gov/phil>

¹These authors contributed equally to this article.

Novel Henipa-like Virus, Mojiang Paramyxovirus, in Rats, China, 2012

Technical Appendix



Figure. Genomic organization of Mojiang paramyxovirus. N, nucleocapsid protein; L, large protein; P/V/W/C, phosphoprotein; M, matrix protein; F, fusion protein; G, attachment glycoprotein.

Table. Conserved nucleotide sequences for the gene start, intergenic region, and gene stop of Mojiang paramyxovirus and henipaviruses*

Variable	Gene stop	Intergenic region	Gene start
MojV, gene			
/N		CTT	AGGATTCAGG
N/P	TTAAACAAAA	CTT	AGGATCCAAG
P/M	TCATAAAAAA	CTT	AGGAGTCAAG
M/F	ATATAAAAAA	CTT	AGGTGTCAGG
F/G	TTAATAAAAA	CTT	AGGAGTCAGG
G/L	TTACAAAAAA	CTT	AGGATTCACG
Consensus sequences for henipaviruses			
	TWAHRAAAAA	CTT	AGGANMCARG

*N, nucleocapsid; P, phosphoprotein (P/V/W/C); M, matrix; F, fusion; G, glycoprotein; L, large protein.