

***Bartonella* spp. and *Yersinia pestis* Reservoirs, Cusco, Peru**

To the Editor: *Bartonella* spp. are gram-negative alphaproteobacteria that are transmitted between the reservoir and mammal host by hematophagous insects (1). The genus *Yersinia* comprises 11 species, of which *Y. pestis* is the causative agent of plague, a deadly rodent-associated, fleaborne zoonosis (2). Despite the large number of plague cases reported in humans and the large amount of data about human-infecting *Bartonella* spp. in Peru (3), no data have been published about which rodent species are reservoirs of these pathogens in this country.

La Convención Province, where cases of bartonellosis occurred during 1998 (4), is located in the northeastern part of Cusco, Peru. Although, to our knowledge, no human cases of plague have been reported in this province, a plague outbreak was recently detected in Junin Province (M.A. Quispe-Riclude, pers. comm.), which is located northwest of La Convención Province.

A total of 28 rodents were captured during 2010–2011 in 3 villages (Alto Ivochote, Aguas Calientes, and Yomentoni) in the Echarate District, La Convención Province. Traps had been set in intradomiciliary, peridomiciliary, and extradomiciliary settings. Spleens of animals were obtained, and DNA was isolated by using the Illustra Tissue and Cells Genomic Prep Mini Spin Kit (GE Healthcare, Little Chalfont, UK).

Rodents were examined for *Bartonella* spp. DNA by using a PCR and primers CS443f and CS1210r specific for a 767-bp fragment of the citrate synthase gene (5). Screening for plague was performed by using PCR primers Yp1 and Yp2 specific for a plasminogen activator protein (*pla*) encoded by the *Y. pestis*-specific pPLA plasmid (6).

New and published *Bartonella* spp. and *Y. pestis* sequences were obtained from GenBank and compared by using the nucleotide-nucleotide basic local sequence alignment tool (BLAST) (blastn) program (www.ncbi.nlm.nih.gov/Class/MLACourse/Modules/BLAST/nucleotide_blast.html). The χ^2 test was used to determine statistical differences in the prevalence of both pathogens among host species and villages.

Overall prevalences for *Y. pestis* and *Bartonella* spp. were 17.9% and 21.4%, respectively (Table). Co-infections with both bacteria were found in 3 (10.7%) rodents: 2 *Hylaeamys perenensis* rodents and 1 *Oecomys* spp. rodent. *Bartonella* prevalence was higher in *H. perenensis* rats than in *Rattus rattus* rodents ($p < 0.001$). Rodents positive for *Bartonella* spp. were found in the 3 study villages, and prevalence for Aguas Calientes was higher than that for Alto Ivochote ($p < 0.001$). One of 8 rodents trapped inside houses and 1 of 2 rodents trapped at peridomestic sites were positive for *Bartonella* spp.

Sequence analysis identified 3 citrate synthase gene sequences (GenBank accession nos. KF021602–KF021604) that had 98% and 99% sequence similarity to genotypic variant A3 of the undescribed *Bartonella*

genogroup A, which was obtained from *Oryzomys palustris* rats in the southeastern United States (7). One genotype (isolate B259) was identified in *H. perenensis* rats, and 2 other genotypes were identified in 1 *H. perenensis* rodent and 1 *Oecomys* spp. rodent (isolates B273 and B280, respectively). We propose that the genotype of isolates B273 and B280 is variant A6 and the genotype of isolate B259 is variant A7. A previous study reported that the A, B, and C genogroups contain independent species (8).

The *pla* amplicons (GenBank accession nos. KF214264–KF214266) had 98% sequence identity with *Y. pestis* reference sequences. Plague prevalence was higher in *H. perenensis* rats than in *R. rattus* rats ($p < 0.05$). Infected rodents were found in all villages studied except Yomentoni, and prevalence in Aguas Calientes was higher than in Alto Ivochote ($p < 0.01$). Two (25%) of 8 rodents trapped inside houses were infected with *Y. pestis*.

This study suggests that infections of rodents with *Bartonella* spp. and *Y. pestis* are common and widespread throughout the Echarate District. It also shows the role of *H. perenensis* and *Oecomys* spp. rodents as reservoirs of both pathogens. This role was confirmed by amplifying the chromosomal ferric iron uptake regulation gene by using PCR primers Yp1 and Yp2, as described by Hinnebusch et al. (9). The epidemiologic role of rodent-borne *Bartonella* spp. as a cause of disease in humans is emerging in the Americas. This role has been suggested by identification of a novel rodent-associated *Bartonella* strain causing febrile illness in the

Table. Prevalence of *Bartonella* spp. and *Yersinia pestis* in rodents from Echarate District, Cusco, Peru

Study area	Rodent host species (no.)	No. (%) positive for <i>Yersinia pestis</i>	No. (%) positive for <i>Bartonella</i> spp.
Alto Ivochote	<i>Rattus rattus</i> (20)	3 (15)	0 (0)
Alto Ivochote	<i>Hylaeamys perenensis</i> (1)	0	1 (100)
Aguas Calientes	<i>Hylaeamys perenensis</i> (2)	2 (100)	2 (100)
Aguas Calientes	<i>Oecomys</i> spp. (1)	1 (100)	1 (100)
Yomentoni	<i>Rattus rattus</i> (4)	0	1 (25)
Total	(28)	6 (21.4)	5 (17.9)

rural southwestern United States (10) and a strain of pathogenic *B. elizabethae*, a bacteria that can cause human endocarditis, in the Huayllacallán Valley in Peru (3).

Because most identified *Bartonella* spp. have been reported as infectious agents for humans, our results should prompt public health concern. However, our findings require further investigation about the pathogenicity of these *Bartonella* genotypes. The detection of both pathogens in intradomestic and peridomestic areas where humans are in close contact with rodents could indicate that the incidence of both diseases in humans from Echarate District might be underestimated.

This study was supported by Agencia Española para la Cooperación Internacional y el Desarrollo under Programa de Cooperación Interuniversitaria (A1/037176/11), the Spanish Ministry of Foreign Affairs and Cooperation (project Red de Investigación Colaborativa de Centros de Enfermedades Tropicales; RD06/0021/0005); and the Spanish Ministry of Health, Madrid. A.M.-A. was supported by a PhD grant from Agencia Canaria de Investigación, Innovación y Sociedad de la Información. M.A.Q.-R. was supported by a research contract from Centro de Excelencia Internacional–Plataforma Atlántica para el Control de las Enfermedades Tropicales.

**Aarón Martín-Alonso,
Mayday Soto, Pilar Foronda,
Elsa Aguilar,
Guillermo Bonnet,
Rosa Pacheco,
Basilio Valladares,
and María A. Quispe-Ricalde**

Author affiliations: University of La Laguna, Canary Islands, Spain (A. Martín-Alonso, P. Foronda, G. Bonnet, B. Valladares, M. A. Quispe-Ricalde); and National University of San Antonio Abad, Cusco, Peru (M. Soto, E. Aguilar, R. Pacheco)

DOI: <http://dx.doi.org/10.3201/eid2006.131194>

References

1. Kamani J, Morick D, Mumcuoglu Y, Harrus S. Prevalence and diversity of *Bartonella* species in commensal rodents and ectoparasites from Nigeria, west Africa. *PLoS Negl Trop Dis*. 2013;7:e2246. <http://dx.doi.org/10.1371/journal.pntd.0002246>
2. Gage KL, Kosoy MY. Natural history of plague: perspectives from more than a century of research. *Annu Rev Entomol*. 2005;50:505–28. <http://dx.doi.org/10.1146/annurev.ento.50.071803.130337>
3. Birtles RJ, Canales J, Ventosilla P, Alvarez E, Guerra H, Llanos-Cuentas A, et al. Survey of *Bartonella* species infecting intradomestic animals in the Huayllacallán Valley, Ancash, Peru, a region endemic for human bartonellosis. *Am J Trop Med Hyg*. 1999;60:799–805.
4. Parola P, Shpynov S, Montoya M, Lopez M, Houpiqian P, Zeaiter Z, et al. First molecular evidence of new *Bartonella* spp. in fleas and a tick from Peru. *Am J Trop Med Hyg*. 2002;67:135–6.
5. Billeter SA, Gundi VA, Rood MP, Kosoy MY. Molecular detection and identification of *Bartonella* species in *Xenopsylla cheopis* (Siphonaptera: Pulicidae) collected from *Rattus norvegicus* in Los Angeles, California. *Appl Environ Microbiol*. 2011;77:7850–2. <http://dx.doi.org/10.1128/AEM.06012-11>
6. Hinnebusch J, Schwan TG. New method for plague surveillance using polymerase chain reaction to detect *Yersinia pestis* in fleas. *J Clin Microbiol*. 1993;31:1511–4.
7. Kosoy MY, Regnery R, Tzianabos T. Distribution, diversity, and host specificity of *Bartonella* in rodents from the southeastern United States. *Am J Trop Med Hyg*. 1997;57:578–88.
8. Chan KS, Kosoy M. Analysis of multi-strain *Bartonella* pathogens in natural host population—do they behave as species or minor genetic variants? *Epidemics*. 2010;2:165–72. <http://dx.doi.org/10.1016/j.epidem.2010.08.002>
9. Hinnebusch BJ, Gage KL, Schwan TG. Estimation of vector infectivity rates for plague by means of a standard curve-based competitive polymerase chain reaction method to quantify *Yersinia pestis* in fleas. *Am J Trop Med Hyg*. 1998;58:562–9.
10. Iralu J, Bai Y, Crook L, Tempest B, Simpson G, McKenzie T, et al. Rodent-associated *Bartonella* febrile illness, southwestern United States. *Emerg Infect Dis*. 2006;12:1081–6. <http://dx.doi.org/10.3201/eid1207.040397>

Address for correspondence: Pilar Foronda, Institute of Tropical Diseases and Public Health of the Canary Islands, University of La Laguna, Avda. Fco. Sanchez s/n, 38203, Tenerife, Canary Islands, Spain; email: pforonda@ull.edu.es

Buruli Ulcer Disease in Republic of the Congo

To the Editor: Buruli ulcer, which is caused by the *Mycobacterium ulcerans* bacterium, is a severe disabling necrotic disease of the skin, occurring mainly in swampy rural areas of western and central Africa. This tropical disease is neglected, despite being the third most common mycobacterial disease of humans, after tuberculosis and leprosy. The disease has become substantially more frequent over the past decade, particularly around the Gulf of Guinea, and has been detected or suspected in at least 31 countries (1). Clinical diagnosis of Buruli ulcer disease should be confirmed by PCR, as recommended by the World Health Organization (WHO); and case-patients should be treated with rifampin/streptomycin daily for 8 weeks (therapy available since 2004), combined, if necessary, with surgery.

Although confirmed cases of Buruli ulcer disease have been reported in all countries neighboring the Republic of the Congo (hereafter called Congo) (2–4), only 1 report of a confirmed case in Congo has been published (5) (Figure, panel A). During 2007–2012, a total of 573 clinical cases of Buruli ulcer disease were reported to WHO by the National Leprosy, Buruli Ulcer and Yaws Control Program in Congo. We report 108 cases (19% of all cases reported) that