

# Tembusu-Related Flavivirus in Ducks, Thailand

**Aunyaratana Thontiravong,  
Patchareeporn Ninvilai, Wikanda Tunterak,  
Nutthawan Nonthabenjawan,  
Supassma Chaiyavong,  
Kingkarn Angkabkingkaew,  
Chatthapon Mungkundar, Woranuch Phuengpho,  
Kanisak Oraveerakul, Alongkorn Amonsin**

Since 2013, outbreaks of disease caused by duck Tembusu virus (DTMUV) have been observed in layer and broiler duck farms in Thailand. The virus is closely related to Chinese DTMUVs and belongs to the Ntaya group of mosquito-borne flaviviruses. These findings represent the emergence of DTMUV in ducks in Thailand.

In 2010, a severe contagious disease emerged in layer and breeder duck farms in China (1). The infected ducks typically exhibited a dramatic reduction in egg production and severe neurologic disorders. The causative agent of this emerging disease was identified as the new duck Tembusu virus (DTMUV), a member of the Ntaya virus group in the genus *Flavivirus* (1,2). In addition to China, new DTMUV was recently detected among ducks in Malaysia (3). In Thailand, a severe contagious disease affecting ducks has newly emerged since 2013. The disease rapidly spread through duck farms in high-density duck-producing areas, causing economic losses for both traditional and agro-industrial duck businesses. This study reports the emergence of DTMUV infection among domestic ducks in Thailand.

## The Study

Since 2013, several layer and broiler duck farms located in high-density duck-producing areas of Thailand have had an emerging, contagious disease characterized by severe neurologic dysfunction and dramatically decreased egg production among domestic ducks. Outbreaks have been reported on farms in the northeastern (Nakhon Ratchasima), eastern (Prachinburi and Chonburi), and central (Suphanburi) provinces of Thailand. At least 7 duck farms were affected, and outbreaks occurred throughout the year (August 2013–September 2014). However, the disease occurred more frequently during the rainy season (July–December). We estimated the mean prevalence of the outbreaks at 17.19% (online


Technical Appendix Figure 1, panels A, B, [http://wwwnc.cdc.gov/EID/article/21/12/15-0600\\_Techapp1.pdf](http://wwwnc.cdc.gov/EID/article/21/12/15-0600_Techapp1.pdf)). Clinical signs were usually observable in broiler ducks >3 weeks of age and in layer ducks during their production period. Infected ducks typically exhibited neurologic signs, including ataxia, reluctance to walk, and progressive paralysis (Figure 1, panels A, B). A remarkable drop in egg production was usually observed among layer ducks. The main pathologic changes were ovaritis, ovarian hemorrhage, and ovarian atrophy (Figure 1, panel C). Splenic enlargement was observed in some ducks. Histopathologic analysis showed moderate multifocal gliosis and perivascular cuffing in the brain (cerebellum) and spinal cord of most sick ducks (Figure 1, panels D, E). Rates of illness and death ranged from 20% to 50% and 10% to 30%, respectively, correlating positively with secondary bacterial infection.

We identified 22 DTMUVs through reverse transcription PCR using E gene-specific primers (1) (online Technical Appendix Table 1). One virus (DK/TH/CU-1) was inoculated into embryonated chicken eggs. The embryos died within 3–5 days after inoculation, with severe cutaneous hemorrhages (Figure 1, panel F). The allantoic fluid tested negative through hemagglutination test and PCR for common duck viruses, including avian influenza virus, Newcastle disease virus and duck herpesvirus 1. In addition, 5 representative viruses (DK/TH/CU-2, DK/TH/CU-3, DK/TH/CU-4, DK/TH/CU-5, DK/TH/CU-6) from duck farms located in the northeastern (3 farms) and the eastern (2 farms) provinces were selected for partial E gene sequencing (Table). The nucleotide sequences of the Thai DTMUVs used in this study were submitted to GenBank under accession nos. KR061333–KR061338.

To characterize Thai DTMUV, DK/TH/CU-1 was subjected to whole-genome sequencing. The whole-genome length of DK/TH/CU-1 is 10,278 nt, encoding 3,426 aa. BLAST analysis (<http://www.ncbi.nlm.nih.gov/blast>) showed that the polyprotein gene sequences of DK/TH/CU-1 shared very high identity (98.3%) with GX2013E, a Chinese DTMUV strain isolated in 2013. Phylogenetic analysis of the polyprotein gene sequence using the neighbor-joining and maximum-likelihood algorithms showed that DK/TH/CU-1 is grouped into the major cluster with mosquito-borne flaviviruses (65.2%–77% nt identity with viruses in the Ntaya group) and is most closely related to Chinese DTMUVs (97.3%–98.3% nt identity). DK/TH/CU-1 shared only 90.3% and 89.4% nt identity with MM1775 strain and Sitiawan virus, which are Tembusu viruses isolated from mosquitos and chickens, respectively (Figure 2, panel A; online Technical Appendix Table 2).

Author affiliations: Chulalongkorn University, Bangkok, Thailand (A. Thontiravong, P. Ninvilai, W. Tunterak, N. Nonthabenjawan, S. Chaiyavong, K. Oraveerakul, A. Amonsin); Animal Health and Technical Service Office, Bangkok (P. Ninvilai, K. Angkabkingkaew, C. Mungkundar, W. Phuengpho)

DOI: <http://dx.doi.org/10.3201/eid2112.150600>



**Figure 1.** Clinical signs and pathologic lesions of duck Tembusu virus (DTMUV)-infected ducks, Thailand. A, B) Clinical signs; DMTUV-infected ducks showed neurologic signs, including inability to stand, ataxia, and paralysis. C) Gross lesion; severe hemorrhage and regression of ovarian follicles. D, E) Histopathologic lesion; moderate multifocal gliosis (black arrows) and perivascular cuffing (white arrows) in cerebellum (D) and spinal cord (E). Scale bars indicate 100  $\mu$ m (D) and 50  $\mu$ m (E). F) Chicken embryos infected with DTMUV strain DK/TH/CU-1. Normal embryo is shown at left; infected embryos at right died 3–5 days after inoculation, with severe cutaneous hemorrhage.

Analysis of the partial E gene sequences of the 5 Thai DTMUVs (DK/TH/CU-2 to 6) showed that the viruses are grouped with DK/TH/CU-1 and Chinese DTMUVs (Figure 2, panel B). The partial E gene sequences of Thai DTMUVs shared 97.5%–99.7% and 96.7%–98.9% nt identity with each other and with the Chinese DTMUVs, respectively. However, the nucleotide identities were lower (88.6%–90.6%) than Malaysian DTMUVs. The E gene sequence of a DK/TH/CU-1 shares only 89.1% and 90.9% nt identity with TMUV strains isolated in 2002 from mosquitos and healthy ducks in Thailand, respectively (6). Analysis of partial NS5 gene showed similar findings

with those of polyprotein and E genes (Figure 2, panel C). DK/TH/CU-1 shared 96.4%–98.1% and 92.7%–93% nt identity with Chinese DTMUVs and Malaysian DTMUVs, respectively.

**Conclusions**

Since 2013, outbreaks of a severe contagious disease among domestic ducks have been occurring and spreading in the high-density duck-producing areas of Thailand, causing substantial economic losses in the agricultural sector. On the basis of pathologic examinations, virus isolation, virus identification and genetic characterization, we found an association with the new DTMUV.




Despite lack of the experimental pathogenicity testing of the virus isolates, our observations on clinical signs and pathologic findings were consistent with previously reported findings of DTMUV infections in China and Malaysia (1,3). Therefore, the isolated DTMUVs can be considered as the causative agent. Because DTMUV is a mosquito-borne flavivirus, it can be transmitted to ducks from mosquitos. Our data indicated that the disease caused by DTMUV occurred most frequently during rainy season, when mosquito activity in Thailand is highest. A previous study detected TMUV in *Culex* mosquitos in Thailand in 2002. The *Culex* mosquito has also proven to be a vector for transmitting TMUV to chickens (6). DTMUV transmission through the fecal–oral route also has been reported (2,7,8). However, the pathogenicity and transmission routes of Thai DTMUV were not determined in this study. Further studies on the Thai DTMUVs should be conducted.

Genetic analyses of polyprotein sequences of the Thai DTMUVs showed higher nucleotide identity with DTMUVs reported from China (97.9%) than with those reported from Malaysia (90.3%), indicating that Chinese DTMUVs are possible ancestors of Thai DTMUVs. Phylogenetic analyses based on polyprotein, E gene and NS5 gene using 2 algorithms (neighbor-joining and maximum-likelihood) have displayed similar results that the Thai isolates were grouped with the Chinese DTMUV with high bootstraps value. The Malaysian DTMUVs were grouped into a subcluster apart from Thai and Chinese DTMUV. Although TMUV strains were isolated from Thai mosquitos and healthy ducks

**Table.** Detailed description of DTMUVs characterized in study of DTMUV in ducks, Thailand\*

Virus name	Study designation	Time of collection	Duck age	Duck type	Location in Thailand	Genome sequencing	GenBank accession no.
DTMUV strain DK/TH/CU-1	DK/TH/CU-1	2013 Nov	39 d	Broiler	Nakhon Ratchasima	WG	KR061333
DTMUV strain DK/TH/CU-2	DK/TH/CU-2	2014 Aug	38 wk	Layer	Chonburi	Partial E	KR061334
DTMUV strain DK/TH/CU-3	DK/TH/CU-3	2014 Aug	35 d	Broiler	Nakhon Ratchasima	Partial E	KR061335
DTMUV strain DK/TH/CU-4	DK/TH/CU-4	2014 Aug	42 d	Broiler	Nakhon Ratchasima	Partial E	KR061336
DTMUV strain DK/TH/CU-5	DK/TH/CU-5	2013 Sep	24 d	Broiler	Nakhon Ratchasima	Partial E	KR061337
DTMUV strain DK/TH/CU-6	DK/TH/CU-6	2013 Oct	35 d	Broiler	Prachinburi	Partial E	KR061338

\*All samples were pooled organs (i.e., brain, spinal cord, spleen, lung, kidney, proventriculus, and intestine). DTMUV, duck Tembusu virus; partial E, partial E gene sequence; WG, whole-genome.



**Figure 2.** Phylogenetic analysis of the nucleotide sequences of polyprotein gene (10,278 bp) (A), partial envelope gene (361 bp) (B), and partial nonstructural 5 gene (900 bp) (C) of duck Tembusu viruses (DTMUVs) from ducks in Thailand and selected reference strains of flaviviruses. The nucleotide sequences were aligned by using Muscle version 3.6 (4). The phylogenetic trees were constructed in MEGA version 6.0 by using the neighbor-joining algorithm with the Kimura-2 parameter model applied to 1,000 replications of bootstrap (5). Circle indicates Thai DTMUVs. Similar results were observed when applying the maximum-likelihood algorithm (online Technical Appendix Figure 2, panels A–C, <http://wwwnc.cdc.gov/EID/article/21/12/15-0600-Techapp1.pdf>). Scale bars indicate nucleotide substitutions per site.

in 2002, the nucleotide sequences of those viruses were less similar to Thai DTMUVs than those of Chinese DTMUVs. Nevertheless, the source of the novel DTMUV emergence in Thailand remains unknown and requires further investigation.

As a member of the *Flavivirus* genus, DTMUV has a high potential to become a zoonotic pathogen that threatens public health. Thus far, DTMUV has not been reported to cause illness in humans. However, DTMUV-specific antibodies and DTMUV RNA were detected in duck farm workers in China (9). Therefore, a novel DTMUV that can cause disease in humans possibly could emerge. Previous studies have reported that DTMUV can infect a wide variety of avian species, including geese, chickens, pigeons, and house sparrows, indicating the continued expansion of its host range (7,10–12). Thus, the continued monitoring of DTMUV in animals and humans is essential to preventing economic losses in animal production as well as zoonotic potential in humans. In summary, our data collectively demonstrate that a newly emerged, contagious disease among ducks in Thailand is caused by DTMUV. Our findings highlight the necessity of systemic surveillance of DTMUVs in animals and in humans for early detection and prevention.

#### Acknowledgments

We thank the staff of the Virology Unit, Department of Veterinary Microbiology, Faculty of Veterinary Science, Chulalongkorn University; Rachod Tantilertcharoen for his contribution in histopathologic analysis; and Joshua A. Felber for reviewing the manuscript.

This research was supported by the Ratchadaphiseksomphot Endowment Fund 2013 of Chulalongkorn University (CU-56-642-HR and WCU-005-HR-57). The authors also thank the Thailand Research Fund (TRG5780010) and Grants for Development of New Faculty Staff, Rachadaphiseksomphot Endowment Fund, Chulalongkorn University. Chulalongkorn University provided financial support to the Center of Excellence for Emerging and Re-emerging Infectious Diseases in Animals, and Thailand Research Fund provided financial support to the TRF Senior Scholar to A.A. (RTA5780006).

Dr. Thontiravong is an instructor at the Faculty of Veterinary Science, Chulalongkorn University, Bangkok. Her research interests include virology of emerging viruses in animals.

#### References

1. Su J, Li S, Hu X, Yu X, Wang Y, Liu P, et al. Duck egg-drop syndrome caused by BYD virus, a new Tembusu-related flavivirus. *PLoS ONE*. 2011;6:e18106. <http://dx.doi.org/10.1371/journal.pone.0018106>
2. Cao Z, Zhang C, Liu Y, Ye W, Han J, Ma G, et al. Tembusu virus in ducks, China. *Emerg Infect Dis*. 2011;17:1873–5. <http://dx.doi.org/10.3201/eid1710.101890>
3. Homonnay ZG, Kovacs EW, Banyai K, Albert M, Feher E, Mato T, et al. Tembusu-like flavivirus (Perak virus) as the cause of neurological disease outbreaks in young Pekin ducks. *Avian Pathol*. 2014;43:552–60. <http://dx.doi.org/10.1080/03079457.2014.973832>
4. Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics*. 2004;5:113. <http://dx.doi.org/10.1186/1471-2105-5-113>
5. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731–9. <http://dx.doi.org/10.1093/molbev/msr121>
6. O'Guinn ML, Turell MJ, Kengluetcha A, Jaichapor B, Kankaew P, Miller RS, et al. Field detection of Tembusu virus in western Thailand by RT-PCR and vector competence determination of select *Culex* mosquitoes for transmission of the virus. *Am J Trop Med Hyg*. 2013;89:1023–8. <http://dx.doi.org/10.4269/ajtmh.13-0160>
7. Tang Y, Diao Y, Yu C, Gao X, Ju X, Xue C, et al. Characterization of a Tembusu virus isolated from naturally infected house sparrows (*Passer domesticus*) in northern China. *Transbound Emerg Dis*. 2013;60:152–8. <http://dx.doi.org/10.1111/j.1865-1682.2012.01328.x>
8. Yan P, Zhao Y, Zhang X, Xu D, Dai X, Teng Q, et al. An infectious disease of ducks caused by a newly emerged Tembusu virus strain in mainland China. *Virology*. 2011;417:1–8. <http://dx.doi.org/10.1016/j.virol.2011.06.003>
9. Tang Y, Gao X, Diao Y, Feng Q, Chen H, Liu X, et al. Tembusu virus in human, China. *Transbound Emerg Dis*. 2013;60:193–6. <http://dx.doi.org/10.1111/tbed.12085>
10. Chen S, Wang S, Li Z, Lin F, Cheng X, Zhu X, et al. Isolation and characterization of a Chinese strain of Tembusu virus from Hy-Line Brown layers with acute egg-drop syndrome in Fujian, China. *Arch Virol*. 2014;159:1099–107. <http://dx.doi.org/10.1007/s00705-013-1931-0>
11. Han K, Huang X, Li Y, Zhao D, Liu Y, Zhou X, et al. Complete genome sequence of goose tembusu virus, isolated from jiangnan white geese in Jiangsu, China. *Genome Announc*. 2013;1:e0023612. <http://dx.doi.org/10.1128/genomeA.00236-12>
12. Liu P, Lu H, Li S, Wu Y, Gao GF, Su J. Duck egg drop syndrome virus: an emerging Tembusu-related flavivirus in China. *Sci China Life Sci*. 2013;56:701–10. <http://dx.doi.org/10.1007/s11427-013-4515-z>

Address for correspondence: Alongkorn Amonsin, Department of Veterinary Public Health, Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand, 10330; email: [alongkornamonsin1@gmail.com](mailto:alongkornamonsin1@gmail.com)

## Like our podcasts?

Sign up to receive email announcements when a new podcast is available.

[www.cdc.gov/ncidod/eid/subscribe.htm](http://www.cdc.gov/ncidod/eid/subscribe.htm)

# Tembusu-Related Flavivirus in Ducks, Thailand

## Technical Appendix

### Methods

#### Outbreak Investigations

During August 2013–September 2014, we investigated outbreaks of a contagious duck disease among ducks characterized by severe neurologic dysfunction and dramatic decreases in egg production in layer and broiler duck farms in Thailand. Epidemiologic information, clinical observations, postmortem examinations, samples collection, and laboratory testing were recorded and analyzed to determine the etiology of the outbreaks.

#### Virus Isolation and Identification

Visceral organ samples were collected from affected ducks, including brain, spinal cord, spleen, lung, kidney, proventriculus, and intestine. Each sample was homogenized in sterile phosphate-buffered saline at a 10% suspension (w/v), centrifuged at  $3,000 \times g$  for 15 min, then filtered through 0.2- $\mu\text{m}$  filters. The filtered suspensions were inoculated into the allantoic cavities of 9-day-old embryonated chicken eggs. The allantoic fluids and tissue suspensions were then examined for the presence of duck Tembusu virus (DTMUV) by reverse transcription PCR (RT-PCR) by using E gene–specific primers (1).

The samples were also tested for avian influenza virus (2), Newcastle disease virus (3,4), and duck herpesvirus (5) to rule out other common viruses that can cause similar symptoms. The tissue suspensions and virus isolates were also tested by hemagglutination tests against 1% chicken erythrocytes at 25°C, pH 7.4 to exclude avian hemagglutinating viruses, including avian influenza virus and Newcastle disease virus.

#### Whole-Genome Sequencing and Phylogenetic Analysis of Thai DTMUV

In this study, 1 DTMUV isolate from Thailand (DK/TH/CU-1) was selected and subjected to whole-genome sequencing. Additionally, 5 other isolates (DK/TH/CU-2,

DK/TH/CU-3, DK/TH/CU-4, DK/TH/CU-5, and DK/TH/CU-6) were subjected to partial E gene sequencing. Viral RNAs were extracted from allantoic fluid by using NucleoSpin Extract Viral RNA Kit (Macherey-Nagel, Düren, Germany) in accordance with the manufacturer's instructions. RT-PCR was performed in a single-step reaction by using the AccessQuick RT-PCR System (Promega, Madison, WI, USA). Primers were designed on the basis of the complete genome sequences of DTMUV available in GenBank. The oligonucleotide primer sequences are shown in online Technical Appendix Table 1. The amplicons were then subjected to DNA sequencing (1st Base Laboratories Sdn Bhd, Malaysia). The validated nucleotide sequences were assembled by using SeqMan software v.5.03 (DNASTAR Inc., Madison, Wisconsin, USA). These nucleotide sequences of the Thai DMTUV were submitted to GenBank under accession nos. KR061333–8.

Phylogenetic analysis was performed by comparing the nucleotide sequences of polyprotein, E gene and NS genes of the Thai DMTUVs with those of other reference flaviviruses available in GenBank. The nucleotide sequences were aligned by using Muscle v.3.6 (7). The phylogenetic trees were constructed in MEGA v.6.0 by using the neighbor-joining and maximum-likelihood algorithms (8). To determine nucleotide identities, the nucleotide sequences of Thai, Malaysian, and Chinese DTMUVs were aligned and compared in MegAlign software v.5.03 (DNASTAR Inc.) with other reference flaviviruses.

## References

1. Su J, Li S, Hu X, Yu X, Wang Y, Liu P, et al. Duck egg-drop syndrome caused by BYD virus, a new Tembusu-related flavivirus. PLoS ONE. 2011;6:e18106. [PubMed](http://dx.doi.org/10.1371/journal.pone.0018106) <http://dx.doi.org/10.1371/journal.pone.0018106>
2. Spackman E, Senne DA, Myers TJ, Bulaga LL, Garber LP, Perdue ML, et al. Development of a real-time reverse transcriptase PCR assay for type A influenza virus and the avian H5 and H7 hemagglutinin subtypes. J Clin Microbiol. 2002;40:3256–60. [PubMed](http://dx.doi.org/10.1128/JCM.40.9.3256-3260.2002) <http://dx.doi.org/10.1128/JCM.40.9.3256-3260.2002>
3. Ke GM, Yu SW, Ho CH, Chu PY, Ke LY, Lin KH, et al. Characterization of newly emerging Newcastle disease viruses isolated during 2002–2008 in Taiwan. Virus Res. 2010;147:247–57. [PubMed](http://dx.doi.org/10.1016/j.virusres.2009.11.008) <http://dx.doi.org/10.1016/j.virusres.2009.11.008>

4. Liu H, Wang Z, Wu Y, Zheng D, Sun C, Bi D, et al. Molecular epidemiological analysis of Newcastle disease virus isolated in China in 2005. *J Virol Methods*. 2007;140:206–11. [PubMed](#)  
<http://dx.doi.org/10.1016/j.jviromet.2006.10.012>
5. Li H, Liu S, Kong X. Characterization of the genes encoding UL24, TK and gH proteins from duck enteritis virus (DEV): a proof for the classification of DEV. *Virus Genes*. 2006;33:221–7.  
[PubMed](#) <http://dx.doi.org/10.1007/s11262-005-0060-6>
9. Huang X, Han K, Zhao D, Liu Y, Zhang J, Niu H, et al. Identification and molecular characterization of a novel flavivirus isolated from geese in China. *Res Vet Sci*. 2013;94:774–80. [PubMed](#)  
<http://dx.doi.org/10.1016/j.rvsc.2012.11.014>
7. Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics*. 2004;5:113. [PubMed](#) <http://dx.doi.org/10.1186/1471-2105-5-113>
8. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731–9. [PubMed](#) <http://dx.doi.org/10.1093/molbev/msr121>

**Technical Appendix Table 1.** Primers used for whole-genome sequencing of a Thai duck Tembusu virus

Primer name	Sequence (5'→3')	Reference
P1f	AGAAGTTCRYCTGTGTGA	(6)
DF_R638	CAGCAGTCTATGTCTTCAGG	This study
DF_F441	CGATAGTTGCTGGGCTGAAGC	This study
DF_R1115	GCAGTAAGATCTCACAACCGC	This study
DF_F954	GCTTCAGCTGTCTGGGGATGC	This study
DF_R1650	CAATGACTCTTTGTTTTGCCACG	This study
DF_F1480	CTACACYGCTGAGATGGAGG	This study
DF_R2463	GCCAAGTCGATTGAGCACCCC	This study
DF_F2353	GGCACTGCTATTGTGGATGGG	This study
DF_R3339	GGTGGGGTGGTGCAAGACC	This study
DF_F3302	GGAACAACCTGCACAGTAACG	This study
DF_R4694	GCATGACTCCCACTCCAGCC	This study
DF_F4406	GCATCACAGAGATTTGATGTGG	This study
DF_R5162	CCTGAACCTGGATGTAGGTCC	This study
DF_F4874	GCAAGTCATCGTCGTGCAACC	This study
DF_R5582	GCTCTTCAATGTCTGTTATTGGC	This study
DF_F5399	GCTCACACCTCAGCGAGTGC	This study
DF_R6249	GGTCATTGTAACCTATCCCAGC	This study
DF_F5928	CCAGTTCCTATAACATCAGCC	This study
DF_R6678	GCTCCAAGACCTGTCTTCCC	This study
DF_F6494	CGCTCACAGAATGACAGAATCC	This study
DF_R7348	GGAACATCTGTAGCCACTATGC	This study
DF_F6807	GAACCAGAGAGACAGAGATCGC	This study
DF_R8158	CCCTAGCTAGCCATTCTCGG	This study
DF_F7940	GCAGGTTCAAGGAAGTGAGAGG	This study
DF_R8536	GGATTGTCTTGGTCATAATGCC	This study
DF_F8383	GGATGCACAAAACCAACCGC	This study
DF_R9215	GGCCGAGATGTCACGCAGC	This study
DF_F8084	GCTGTGTGACATAGGTGAAGC	This study
DF_R9449	CCACTTCCCCTCTGGTCTTCC	This study
DF_F9274	GGGACACTAGAATAACCAAGGC	This study
DF_R10485	CCAACATCCGGTGGCAGGG	This study


**Technical Appendix Table 2.** Nucleotide identities of the polyprotein gene sequences of a Thai DTMUV (DK/TH/CU-1) with reference viruses of Ntaya virus group

	DK/TH/CU-1	WFZ_2012	CJD05	xz_2012	YY5	goose	pigeon	DEDSV-									ITV	ROCV	ILHV	ZIKV
								byd1	BYD-1	MM1775	STWV	NTAV	BAGV							
DK/TH/CU-1	100	97.7	97.6	97.7	97.9	97.8	97.9	97.9	97.9	90.3	89.4	76.9	76.7	77	69.4	68.8	65.2			
WFZ_2012			99.1	99.2	99.4	99.3	99.3	99.4	99.4	90.6	89.4	76.9	76.5	77	69.4	68.6	65.2			
CJD05				99.1	99.3	99.2	99.3	99.4	99.4	90.6	89.3	76.9	76.5	76.8	69.3	68.6	65.3			
xz_2012					99.4	99.3	99.3	99.4	99.4	90.7	89.5	77	76.5	76.8	69.3	68.7	65.4			
YY5						99.5	99.6	99.6	99.6	90.7	89.5	77	76.6	76.9	69.4	68.7	65.3			
goose							99.9	99.6	99.6	90.6	89.5	77	76.5	76.8	69.4	68.7	65.3			
pigeon								99.6	99.6	90.7	89.5	77	76.5	76.9	69.4	68.7	65.3			
DEDSV-byd1									100	90.7	89.5	77	76.6	76.9	69.4	68.7	65.3			
BYD-1										90.7	89.5	77	76.6	76.9	69.4	68.7	65.3			
MM1775											94	76.5	76.6	76.9	69.3	68.6	65.1			
STWV												76.5	76.5	76.8	69.2	68.4	65.1			
NTAV													79.9	80.2	69	68.5	65.2			
BAGV														94.9	69.3	69.1	65.2			
ITV															69.6	69.1	65.2			
ROCV																74.9	65.4			
ILHV																	65.4			
ZIKV																		100		


\*BAGV, Bagaza virus; DK/TH/CU-1, Thai duck tembusu virus; ILHV, Ilheus virus; ITV, Israel turkey meningoencephalomyelitis virus; WFZ\_2012, CJD05, xz\_2012, YY5, goose, pigeon, DEDSV-byd1, BYD-1, Chinese duck tembusu viruses; MM1775, Tembusu virus strain MM1775; NTAV, Ntaya virus; ROCV, Rocio virus; STWV, Sitiawan virus; ZIKV, Zika virus.




**A**




**B**




**Technical Appendix Figure 1.** Details of DTMUV outbreaks in Thailand during 2013–2014. A) Locations of DTMUV outbreaks in Thailand. Circles represent provinces where the DMTUV isolated. NR, Nakhon Ratchasima; PC, Prachinburi; CB, Chonburi; SB, Suphanburi. B) Monthly prevalence of DTMUV in farmed ducks, Thailand, 2013–2014. DTMUV, duck Tembusu virus.



Partial E gene



## NS5 gene



**Technical Appendix Figure 2.** Phylogenetic analysis of the nucleotide sequences of polyprotein gene (10,278 bp), partial E gene (361 bp), and partial NS gene (900 bp) of Thai DTMUVs and selected reference strains of flaviviruses. The nucleotide sequences were aligned by using Muscle v.3.6 (7). The phylogenetic trees were constructed in MEGA v.6.0 by using the maximum-likelihood algorithm with the Hasegawa-Kishino-Yano (HKY) model applied to 500 replications of bootstrap (8). Circle indicates Thai DTMUVs.