

Transmission Potential of Influenza A(H7N9) Virus, China, 2013–2014

Adam J. Kucharski,¹ Harriet L. Mills,¹
Christl A. Donnelly, Steven Riley

To determine transmission potential of influenza A(H7N9) virus, we used symptom onset data to compare 2 waves of infection in China during 2013–2014. We found evidence of increased transmission potential in the second wave and showed that live bird market closure was significantly less effective in Guangdong than in other regions.

From February 19, 2013, through April 22, 2014, a total of 429 cases of influenza A(H7N9) virus infection in humans in China were reported and occurred in 2 outbreak waves. During the first wave in spring 2013, live bird markets were closed in several parts of China (1,2); these market closures substantially reduced the risk for infection in affected regions (3). During a second wave in autumn 2013 (4), markets were again closed in some provinces (5–7). Analysis of the largest clusters of subtype H7N9 virus infection in 2013 suggested that the basic reproduction number (R_0 , the average number of secondary cases generated by a typical infectious host in a fully susceptible population) was higher in some clusters than in others (8,9), although the absence of sustained transmission implied that R_0 was less than the critical value of 1. To determine the transmission potential of influenza A(H7N9) virus in the first and second waves in 2013, we compared symptom onset data. We also measured the extent to which market closures in 2014 reduced spillover hazard (i.e., risk for animal-to-human infection).

The Study

We focused on the locations of the 6 largest outbreaks: Shanghai, Zhejiang, and Jiangsu (first wave) and Guangdong, Zhejiang, and Jiangsu (second wave). To infer market hazard and human-to-human transmission potential, we used a statistical model of infection spillover (9). We assumed that cases could be generated in 1 of 2 ways: on each day, the expected number of reported cases was equal to the sum of animal exposure and secondary cases generated by earlier infectious hosts (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/21/5/14-1137-Techapp1.pdf>).

Author affiliations: London School of Hygiene and Tropical Medicine, London, UK (A.J. Kucharski); National Institutes of Health, Bethesda, Maryland, USA (A.J. Kucharski); Imperial College London, London (A.J. Kucharski, H.L. Mills, C.A. Donnelly, S. Riley)

DOI: <http://dx.doi.org/10.3201/eid2105.141137>

Use of such a framework enables estimation of the degree of human-to-human transmission from symptom onset data and of exposure hazard from markets; the accuracy of these estimates is greatly improved when the timing of a sudden change in hazard, such as a market closure, is known (9). We therefore constrained the timing of the drop in exposure hazard to reported market closure dates (online Technical Appendix Table 1). We also estimated R_0 for each of the 6 outbreaks. For patients with known exposure, cluster reports suggest that the serial interval (time delay between symptom onset in primary and secondary case-patients) could be 7–8 days (online Technical Appendix Table 2). We therefore assumed a serial interval of 7 days for our main analysis and tested a range of values from 3 to 9 days during sensitivity analysis. We adjusted for potential delays between symptom onset and case report on the basis of the distribution of delays to date (online Technical Appendix Figure 1).

During the first wave, cases were initially concentrated around Shanghai; reports centered on the city and neighboring Zhejiang and Jiangsu (Figure 1, panel A). A wave-like relationship between location and onset timing was apparent; distance between the location of the first case-patient in Shanghai and subsequent case-patients increased over time (Figure 1, panel B). The pattern of cases at the start of the second wave suggests that infection did not spread outward from a single source; in October 2013, initial cases occurred in Guangdong and Zhejiang.

We used our statistical model to estimate the relative contributions of animal-to-human and human-to-human transmission. In Zhejiang, Shanghai, and Guangdong, market hazard clearly increased and decreased at the start and end of the outbreak, respectively (Figure 2). We also estimated R_0 for different regions over the 2 outbreak waves (Table). Although our estimates for Jiangsu did not change significantly between the 2 waves, for Zhejiang, R_0 was significantly higher for the second wave than for the first wave in spring 2013 ($p = 0.045$). We estimated R_0 to be 0.06 (95% credible interval [CrI] 0.00–0.25) in the first wave and 0.35 (95% CrI 0.15–0.65) in the second.

Using our estimates for R_0 and market hazard, we estimated the number of cases in each outbreak that resulted from human-to-human rather than animal-to-human transmission. We found evidence of a small but significant amount of transmission between humans in the first and second waves (Table). Our findings agree with reports of possible human clusters in the first wave (1,10–12) and corroborate media reports of possible human clusters in Zhejiang and Guangdong

¹These authors contributed equally to this article.




Figure 1. Spatial and temporal distribution of reported cases of influenza A(H7N9) virus infection among humans, China, 2013–2014. Onset of the first case in wave 1 was February 19, 2013 (although the case was not reported until the end of March 2013); onset of the last case in wave 1 was July 27, 2013; only 4 cases occurred in May–July 2013. Onset of the first case in wave 2 was October 7; onset of the last case in our time series was April 17, 2014. A) Case onset reports across all regions. Colors indicate the 4 largest geographic clusters; black indicates all other cases. B) Spatial pattern of reported cases. Points show geodesic distance between the first reported case of influenza A(H7N9) virus infection (in Shanghai) and location of each subsequent reported case. Cases are colored by region as in panel A.

during 2013–2014. We identified 5 clusters during the first wave (February–April 2013) and 8 clusters during the second wave (November 2013–May 2014); the clusters in both waves had median size of 2 cases per cluster (online Technical Appendix Table 2). These conclusions were robust under different assumptions about the duration of serial interval (online Technical Appendix Figures 2, 3).

During the second wave, market closures in Zhejiang began on January 22, 2014, and ended on January 26, 2014 (Table). The reduction in spillover hazard after these closures was significant. We estimated that closures for a serial interval of 7 days reduced hazard by 97% (95% CrI 92%–99%). During 2013, estimated effectiveness was similar in Zhejiang (99%; 95% CrI 97%–100%) and Shanghai (99%; 95% CrI 95%–100%). These estimates are in agreement with those from other analyses for the first wave (3). The 95% CrI was broader for Jiangsu, however, where estimated effectiveness was 97% (95% CrI 80%–100%). In Guangdong, Guangzhou markets closed on February 16, 2014, and reopened on February 28; markets in other cities in Guangdong closed around the same time for 2 weeks. Our results suggest that these closures reduced hazard by 73% (95% CrI 53%–89%). This reduction was significantly smaller than that for Shanghai and Zhejiang ($p < 0.01$). Our result was robust at different serial intervals of infection (online Technical Appendix Figure 4).

Despite the effectiveness of closures during the first wave, interventions in most regions were delayed until

after the Chinese New Year (January 31, 2014). Some regions are investigating alternative market practices: Guangzhou has implemented a trial of a permanent ban on live poultry sales in certain markets, potentially to extend over the entire city by 2024 (5). Our results support recommendations made after the first wave of outbreaks in 2013 (3), which suggest that prompt closure of markets could lead to substantially fewer infections. However, our finding that the relative effectiveness of the shorter closure in Guangdong was lower suggests that such interventions are needed for a sufficiently long time to prevent recurrence.

Our study has limitations. First, case data were insufficient for us to jointly infer serial interval and transmissibility. We therefore tested our results against a wide range of plausible assumptions about the serial interval of infection (online Technical Appendix). We also assumed that the market hazard increased and decreased in a simple stepwise manner (Figure 2). Local market density could also influence the size of spillover hazard and, hence, effectiveness of interventions (13). If the market hazard could be better characterized (e.g., by longitudinal serologic surveillance [14]), the accuracy of our estimates would probably be improved (9). When estimating R_0 , we did not incorporate individual-level variability in transmission and potential superspreading events. However, the framework that we used can still produce reliable estimates of R_0 when a population contains superspreaders (9).




Figure 2. A posteriori probability estimates of spillover hazard for influenza A(H7N9) virus infection in China, by region. Black dots show total number of reported influenza A(H7N9) virus cases for which symptom onset occurred on a given date. Red shading shows a posteriori probability estimate of spillover hazard (i.e., the expected number of cases resulting from animal-to-human transmission on each day). A serial interval of 7 days was assumed. A) Zhejiang, 2013–2014; B) Jiangsu, 2013–2014; C) Shanghai, first outbreak wave, 2013; D) Guangdong, second outbreak wave, 2013–2014.

Conclusions

We found no evidence of reduced human-to-human transmission between the 2 waves. For a serial interval of 7 days, we estimated that R_0 increased in Zhejiang. Furthermore, the effectiveness of live bird market closures varied between regions; short-term closures were substantially less effective than interventions in other regions. These results emphasize the value of prompt and sustainable control measures during outbreaks of influenza A(H7N9) virus infection.

This work was funded by the Wellcome Trust (project grant 093488/Z/10/Z); the Medical Research Council (United Kingdom, project grant MR/J008761/1; Fellowship MR/K021524/1); National Institute for General Medical Sciences (United States, MIDAS U01 GM110721-01); the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 278433-PREDEMICS; the Fogarty International Center (United States, R01 TW008246-01); and the Research and Policy for Infectious Disease Dynamics program also from

Table. Estimates of human-to-human transmission and effectiveness of live bird market closures, China, 2013–2014*

Region, outbreak wave	Total no. cases	R_0 (95% CrI)	Human-to-human transmission, no. cases (95% CrI)	Hazard reduction, % (95% CrI)
Shanghai, first	29	0.32 (0.06–0.60)	11.0 (2.3–14.8)	99 (95–100)
Jiangsu				
First	23	0.24 (0.03–0.69)	6.7 (2.0–12.2)	97 (80–100)
Second	26	0.13 (0.01–0.41)	2.9 (0.1–8.7)	NC
Zhejiang				
First	46	0.06 (0.00–0.25)	3.8 (0.8–12.4)	99 (97–100)
Second	92	0.35 (0.15–0.65)	32.5 (17.3–48.9)	97 (92–99)
Guangdong, second	103	0.16 (0.01–0.54)	16.7 (1.0–48.6)	73 (53–89)

*A serial interval of 7 days was assumed. For sensitivity analysis, see online Technical Appendix (<http://wwwnc.cdc.gov/EID/article/21/5/14-1137-Teachapp1.pdf>). CrI, credible interval; NC, not calculated; R_0 , reproduction number (average number of secondary cases generated by a typical infectious host in a fully susceptible population).

Fogarty International Center with the Science and Technology Directorate, Department of Homeland Security.


Dr. Kucharski is a research fellow in infectious disease epidemiology at the London School of Hygiene and Tropical Medicine. His research focuses on the dynamics of emerging infections and how population structure and social behavior shape disease transmission at the animal–human interface. Dr. Mills is a postdoctoral research associate in the Medical Research Council Centre for Outbreak Analysis and Modelling, Department of Infectious Disease Epidemiology, Imperial College London. Her research includes examination of human mobility patterns and their influence on disease transmission and the analysis of emerging epidemics.

References

- Li Q, Zhou L, Zhou M, Chen Z, Li F, Wu H, et al. Epidemiology of human infections with avian influenza A(H7N9) virus in China. *N Engl J Med*. 2014;370:520–32. <http://dx.doi.org/10.1056/NEJMoa1304617>
- Murhekar M, Arima Y, Horby P, Vandemaekel KA, Vong S, Zijian F et al. Avian influenza A (H7N9) and the closure of live bird markets. *Western Pac Surveill Response*. 2013;4:4–7. <http://dx.doi.org/10.5365/wpsar.2013.4.2.008>
- Yu H, Wu JT, Cowling BJ, Liao Q, Fang VJ, Zhou S, et al. Effect of closure of live poultry markets on poultry-to-person transmission of avian influenza A H7N9 virus: an ecological study. *Lancet*. 2014;383:541–8. [http://dx.doi.org/10.1016/S0140-6736\(13\)61904-2](http://dx.doi.org/10.1016/S0140-6736(13)61904-2)
- Chen E, Chen Y, Fu L, Chen Z, Gong Z, Mao H, et al. Human infection with avian influenza A (H7N9) virus re-emerges in China in winter 2013. *Eurosurveill*. 2013;18:pii:20616. <http://dx.doi.org/10.2807/1560-7917.ES2013.18.43.20616>
- Lau M, Wei L. Guangzhou begins trial ban of live poultry sales after H7N9 wet market tests. *South China Morning Post*. 2014 May 5; *News/China/Health*.
- Huixin W. Some live poultry markets closed permanently. *Shanghai Daily*. 2014 Feb 24.
- Wu P, Jiang H, Wu JT, Chen E, He J, Zhou H, et al. Poultry market closures and human infection with influenza A(H7N9) virus, China, 2013–14. *Emerg Infect Dis*. 2014;20:1891–4. <http://dx.doi.org/10.3201/eid2011.140556>
- Chowell G, Simonsen L, Towers S, Miller MA, Viboud C. Transmission potential of influenza A/H7N9, February to May 2013, China. *BMC Med*. 2013;11:214. <http://dx.doi.org/10.1186/1741-7015-11-214>
- Kucharski AJ, Mills HL, Pinsky A, Fraser C, Van Kerkhove MD, Donnelly CA, et al. Distinguishing between reservoir exposure and human-to-human transmission for emerging pathogens using case onset data. *PLoS Curr*. 2014;6:pii:ecurrents.outbreaks.e1473d9bfc99d080ca242139a06c455f.
- Liu T, Bi Z, Wang X, Li Z, Ding S, Bi Z, et al. One family cluster of avian influenza A (H7N9) virus infection in Shandong, China. *BMC Infect Dis*. 2014;14:98. <http://dx.doi.org/10.1186/1471-2334-14-98>
- Hu J, Zhu Y, Zhao B, Li J, Liu L, Gu K, et al. Limited human-to-human transmission of avian influenza A(H7N9) virus, Shanghai, China, March to April 2013. *Eurosurveill*. 2014;19:pii:20838.
- Xiao C, Li KB, Chen ZQ, Di B, Yang ZC, Yuan J, et al. Transmission of avian influenza A(H7N9) virus from father to child: a report of limited person-to-person transmission, Guangzhou, China, January 2014. *Euro Surveill*. 2014;19:pii:20837.
- Gilbert M, Golding N, Zhou H, Wint GRW, Robinson, TP et al. Predicting the risk of avian influenza A H7N9 infection in live-poultry markets across Asia. *Nat Commun*. 2014;5:4116.
- Chen Z, Li K, Luo L, Lu E, Yuan J, Liu H, et al. Detection of avian influenza A(H7N9) virus from live poultry markets in Guangzhou, China: a surveillance report. *PLoS ONE*. 2014;9:e107266. <http://dx.doi.org/10.1371/journal.pone.0107266>

Address for correspondence: Adam J. Kucharski, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK; e-mail: adam.kucharski@lshtm.ac.uk

A History of the *Emerging Infectious Diseases* Journal



Dr. James Hughes and Dr. D. Peter Drotman discuss the history of the *Emerging Infectious Diseases* journal



<http://www2c.cdc.gov/podcasts/player.asp?f=8635993>

Transmission Potential of Influenza A(H7N9) Virus, China, 2013–2014

Technical Appendix

Data

Using WHO reports and news reports, we collated a line list of reported influenza A/H7N9 cases between 19th February 2013 and 22nd April 2014. In this period there were 429 cases in total, split into two outbreak waves: 144 cases in the spring 2013 wave, which started on 19th February 2013, and 285 cases in the 2013/2014 wave, which began on 7th October 2013 (Figure 1A).

Transmission model

In the model, human cases could be generated in one of two ways [1]. First, they could come from exposure to live bird markets (LBMs). We defined $h_A(t)$ to be the expected number of new human cases with onset on day t due to market exposure. We assumed this to be a step function with S steps and $S - 1$ change points. Cases could also come from human-to-human transmission. In our model, infected individuals had an infectiousness profile described by a Poisson distribution with mean λ , the serial interval of the disease. The number of new infections generated by each infectious individual was dependent on R_0 ; because there were few total infections relative to the population size, we assumed that depletion of the susceptible pool did not affect the dynamics [2]. We defined $h_H(t)$ to be the expected number of new human cases with onset on day t due to previous human cases,

$$h_H(t) = \sum_{i=1}^{I_t} R_0 \frac{\lambda^{t-d_i} e^{-\lambda}}{(t-d_i)!} \quad (1)$$

where d_i was the time infected, hence $t - d_i$ was the time since individual i was infected, and I_t was the total number of infected individuals at time t .

We assumed that the number of new human cases on a given day, N_t , followed a

Poisson distribution with mean $h_A(t) + h_H(t)$. Hence the expected number of cases on day t was given by:

$$m_t = \begin{cases} h_A(t, q) & \text{if } t = 0; \\ \prod_{i=1}^{\min(k, t)} R_0 N_{t-i+1} \frac{t^i e^{-t}}{i!} + h_A(t, q) & \text{if } t > 0 \end{cases} \quad (2)$$

where k is the maximum value the generation time distribution can take.

We used a likelihood-based approach to estimate epidemiological parameters. For a time series of observed human onsets $\{N_t\}_{t=1}^T$, the likelihood of our parameter set is [3]:

$$L(q | N) = \prod_{t=0}^{T-1} \frac{m_t^{N_{t+1}} e^{-m_t}}{N_{t+1}!} \quad (3)$$

The expected number of cases, μ_t , depends both on the shape of the spillover hazard function, $h_A(t, \theta)$, and human-to-human transmission parameters, R_0 and λ . For five of the outbreaks, we assumed that the temporal change in market hazard followed a step-wise hazard function with three steps. The hazard function had five parameters: 3 parameters controlling the relative amplitude of spillover infections, and 2 controlling the timing of the increase and decrease in hazard. We constrained the timing of the drop based on reported market closure dates (Table S1). In the first wave, we assumed that market hazard decreased on a date within 7 days either side of 6th April 2013; in Shanghai, we assumed closure occurred on on 10th April 2013 (± 7 days) in Jiangsu and on 16th April 2013 (± 7 days) in Zhejiang. During the second wave, we assumed that hazard dropped on 26th January 2014 (± 7 days) in Zhejiang and on 16th February 2014 (± 7 days) in Guangdong. As we could not find reports of market closures in Jiangsu in 2014, we used a two-step hazard function for this outbreak, with only an increase in hazard. As well as market hazard, we estimated the basic reproduction number, R_0 , for each of the six outbreaks.

For individual sets of parameter estimates, we used a fixed serial interval, λ . For patients with known exposure, the incubation period of H7N9 infection had a median of 6 days [4] and cluster reports suggest serial interval could be around 7-8 days (Table S2). In our main analysis, we therefore assumed a serial interval of 7 days. However, there is evidence that serial interval for seasonal influenza can be as low as 3-4 days [5]. During sensitivity analysis, we tested a

range of values from 3 to 9 days. We also adjusted for potential delays between symptom onset and case report based on the observed distribution of reporting delays (Figure S1). We assumed that the delay between onset and report followed a normal distribution: based on H7N9 cases reported up to 22nd April 2014, the reporting delay has a mean of 9.0 days and standard deviation of 3.3 days.

Model inference was performed using the full likelihood and Markov Chain Monte Carlo (MCMC) over the space of possible parameter values. We assumed that each parameter was positive, with a flat prior distribution.

The size distribution of human clusters can also be used to estimate the reproduction number of an infection [6]. However, estimation of R_0 from the total outbreak size distribution is implicitly conditional on the infection having so far failed to cause a large epidemic. This condition means it is not possible identify whether R_0 is greater or less than one, and hence whether it has pandemic potential [7]. Moreover, cluster size analysis does not account for change in exposure hazard over time, which can affect the accuracy of transmissibility estimates [8]. The method we here is robust to both of these issues: we did not make the implicit assumption that $R_0 < 1$, and we incorporated information on the temporal change in market hazard when estimating transmission potential.

Calibration of animal-to-human component of model

Before estimating R_0 , we calibrated the market exposure component of the model without the presence of human-to-human transmission. LBMs were closed in Guangdong and Zhejiang in spring 2014. Previous work has shown that a 3 step hazard function performed best according the Bayesian Information Criterion (BIC) for the first wave [1]. We also found most support for 3 step function in 2014 (Table S3). Because we found no reports of closures in Jiangsu in 2014, we assumed a 2 step hazard function for this region.

Technical Appendix Table 1. Details of LBM closures in China in 2014*

Province	City	District	Markets closed	Markets open	Notes
Zhejiang	Hangzhou	Various	22/01/2014, 24/01/2014 and 26/01/2014	unreported	Also closed circuses with live animals
	Hangzhou	Jiangan, Xiacheng, Shangcheng, Gongshu and Xihu	15/02/2014	permanent ban expected	Permanent ban on trading of all birds, for meat and pets. Frozen poultry only to be sold from the end of February.
	Hangzhou	Xiaoshan and Yuhang	15/02/2014	15/05/2014 (expected)	Ban on trading of all birds, for meat and pets.
	Ningbo	Main city districts: Haishu, Jiangdong, Jiangbei, Yinzhou	26/01/2014	unreported	Also stopped people from flying their homing pigeons, closed zoos and scenic bird tours
	Jinhua		by 26/01/2014	unreported	
	Shaoxing		by 26/01/2014	unreported	
Anhui	Anqing	Urban area of Anqing city including Susong County	09/02/2014		
Guangdong	Guangzhou		16/02/2014	28/02/2014	
	Zhongshan		10/02/2014	23/02/2014	
	Shenzhen		31/01/2014	13/02/2014	
Shanghai	Shanghai		31/01/2014	01/05/2014	Extra measures now in place: with extra measures - now will close one day a week for cleaning / sterilization
Hong Kong	Hong Kong		29/01/2014	19/02/2014	The ban on live chicken imports from China will continue for four months. The government intends to screen imported poultry at a holding site.

* In the model, we assumed the drop in hazard resulting from LBM closure in the second wave occurred in Guangdong on 16th February 2014 (± 7 days); and in Zhejiang on 24th January 2014 (± 7 days). For the first wave, based on reported closures [1], we assumed a drop in hazard in Shanghai on 6th April 2013 (± 7 days); in Jiangsu on 23rd March 2013 (± 7 days); and Zhejiang on 31st March 2013 (± 7 days). Dates as reported in public news sources, including Shanghai Daily, Xinhuanet, Guangzhou Daily, Anhui News, China Daily.

Technical Appendix Table 2. Possible human clusters as identified from linelist data and news reports*

Cluster ID	Region	Case ID	Onset Date	Notes
1	Shanghai	1	19-Feb-2013	Father of ID 73 and 76
		73	unknown	Son of ID1
		76	unknown	Son of ID1
2	Shanghai	12	27-Mar-2013	Wife of ID45
		45	02-Apr-2013	Husband of ID12
3	Beijing	44	11-Apr-2013	Child
		88	none	Child, asymptomatic contact of ID44 (parents had bought chickens from parents of ID44)
4	Shandong	106	16-Apr-2013	Father of ID127
		127	27-Apr-2013	Child of ID106
5	Jiangsu	6	21-Mar-2013	Daughter of ID120
		120	08-Mar-2013	Father of ID6
6	Guangdong	165	03-Jan-2014	Father of ID202
		202	14-Jan-2014	Reported daughter of ID165
7	Zhejiang	212	13-Jan-2014	Suspected family cluster (father)
		229	20-Jan-2014	Suspected family cluster (daughter)
		254	23-Jan-2014	Suspected family cluster (mother)
8	Hunan	280	24-Jan-2014	Father of ID286
		286	30-Jan-2014	Daughter of ID280
9	Guangxi	300	27-Jan-2014	Mother of ID310. Traveller, developed fever in Guangdong
		310	03-Feb-2014	Son of ID300. Unclear whether he also travelled.
10	Guangdong	284	27-Jan-2014	Father of ID289
		289	31-Jan-2014	Daughter of ID284
11	Guangdong	274	24-Jan-2014	Probable cluster (father)
		279	29-Jan-2014	Probable cluster (child, cousin of ID293)
		293	26-Jan-2014	Probable cluster (child, cousin of ID279)
12	Zhejiang	141	20-Nov-2013	Father in law of ID144
		144	29-Nov-2013	Son in law of ID141
13	Shandong	446	06-May-2014	Father of ID447
		447	15-May-2014	Son of ID446

*Data from news sources (CIDRAP, Recombinomics, Xinhua Net, South China Morning Post) and journal papers [9, 10]. Case ID refer to the linelist IDs.

Technical Appendix Table 3. Comparison of different market hazard functions in the absence of human-to-human transmission

Outbreak	Model	Likelihood	Parameters	BIC
Guangdong (2nd wave)	3 step	-150.0	5	326.7
	4 step	-146.1	7	329.5
	5 step	-141.9	9	331.8
	6 step	-139.6	11	337.7
	7 step	-140.2	13	349.6
Zhejiang (2nd wave)	3 step	-101.6	5	229.8
	4 step	-98.0	7	233.3
	5 step	-97.8	9	243.5
	6 step	-95.4	11	249.3
	7 step	-96.3	13	261.7

Technical Appendix Table 4. Estimated change in R_0 between 2014 and 2013 for influenza A/H7N9 outbreaks in Jiangsu and Zhejiang provinces*


Outbreak	Serial interval	$R_0^{2014} - R_0^{2013}$ (95%CI)	p-value
Jiangsu	3	-0.14(-0.73-0.48)	0.606
	5	-0.08(-0.56-0.50)	0.721
	7	-0.10(-0.61-0.23)	0.581
	9	-0.01(-0.45-0.48)	0.957
Zhejiang	3	0.25(-0.50-0.72)	0.448
	5	0.27(-0.16-0.62)	0.181
	7	0.28(0.01-0.61)	0.045
	9	0.20(0.04-0.44)	0.020

*Two-sided p-values are given for null hypothesis that R_0 is the same in both outbreak waves.


Technical Appendix Table 5. Estimated difference in market hazard reduction between Guangdong and other geographic regions*

Region	Outbreak wave	Serial interval	Difference in hazard reduction	p-value
Shanghai	1st	3	0.28(0.10-0.59)	0.006
		5	0.26(0.09-0.49)	0.003
		7	0.26(0.09-0.46)	0.003
		9	0.25(0.09-0.45)	0.002
Jiangsu	1st	3	0.25(-0.02-0.56)	0.059
		5	0.23(-0.02-0.46)	0.061
		7	0.24(0.01-0.45)	0.044
		9	0.23(-0.01-0.43)	0.053
Zhejiang	1st	3	0.29(0.12-0.59)	0.001
		5	0.27(0.10-0.49)	0.002
		7	0.26(0.09-0.46)	0.002
		9	0.26(0.09-0.45)	0.001
	2nd	3	0.27(0.09-0.58)	0.007
		5	0.25(0.08-0.47)	0.005
		7	0.24(0.07-0.45)	0.008
		9	0.23(0.07-0.43)	0.006


*Two-sided p-values are given for null hypothesis that there is no difference in hazard between Guangdong and specified region.




Technical Appendix Figure 1. Distribution of delay between case onset and report. We fitted a normal distribution (blue line) to influenza A/H7N9 cases reported between 19th February 2013 and 17th April 2014 (grey bars).



Technical Appendix Figure 2. Estimates of basic reproduction number in different regions as serial interval, λ , varies. Blue point, median of posterior estimate; blue line, 95% credible interval.



Technical Appendix Figure 3. Estimated human-to-human cases in different regions as serial interval, λ , varies. Dashed line, total reported cases; green point, estimated non-index cases; green line, 95% credible interval. (A) Shanghai (1st outbreak wave), (B) Jiangsu (1st wave), (C) Jiangsu (2nd wave), (D) Guangdong (2nd wave), (E) Zhejiang (1st wave), (F) Zhejiang (2nd wave).



Technical Appendix Figure 4. Estimates of reduction in market hazard in different regions as serial interval, λ , varies.