

Highly Pathogenic Avian Influenza A(H5N8) Viruses Reintroduced into South Korea by Migratory Waterfowl, 2014–2015

Jung-Hoon Kwon,¹ Dong-Hun Lee,¹
David E. Swayne, Jin-Yong Noh, Seong-Su Yuk,
Tseren-Ochir Erdene-Ochir, Woo-Tack Hong,
Jei-Hyun Jeong, Sol Jeong, Gyeong-Bin Gwon,
Chang-Seon Song

Highly pathogenic avian influenza A(H5N8) viruses were isolated from migratory waterfowl in South Korea during fall 2014–winter 2015, a recurrence after initial introduction in winter 2014. These reappeared viruses were phylogenetically distinct from isolates circulating in poultry farms in South Korea.

Since the Asian-lineage subtype H5 highly pathogenic avian influenza (HPAI) virus was first detected in China in 1996, outbreaks of infection caused by this virus in poultry have been continuous. The HPAI (H5) viruses have evolved and continue to evolve into many genetic lineages and multiple clades (1). In January 2014, novel reassortant HPAI viruses of subtype H5N8, clade 2.3.4.4, were detected in poultry and wild bird carcasses in South Korea (2). Closely related viruses were also detected in Japan (3) and China (4). Genetic analysis showed that this virus was generated by reassortment of HPAI viruses of eastern China. Subsequently, HPAI (H5N8) viruses spread to Europe and North America and were then reintroduced into South Korea and Japan (5). The HPAI (H5N8) viruses identified in South Korea in early 2014 were divided into groups A (A/Baikal teal/Korea/Donglim3/2014 strain-like) and B (A/breeder duck/Korea/Gochang1/2014-like). Group A viruses further evolved into 3 distinct subgroups: icA1 (Europe/Japan), icA2 (North America/Japan), and icA3 (South Korea/Japan) (5). Wild birds were suspected of being a source of intercontinental transmission because the timing and direction of the outbreak coincided with the migratory route of wild birds (5,6). We sequenced and genetically analyzed the complete genomes of 11 HPAI (H5N8) viruses isolated from wild migratory waterfowl in South Korea during December 2014 and February 2015 and

compared these isolates with other HPAI (H5N8) isolates, including isolates identified from South Korea poultry farms in late 2014.

The Study

A total of 11 HPAI (H5N8) viruses were isolated from 980 samples of wild bird feces and 102 swab samples collected from wild bird habitats in South Korea where active surveillance was conducted during December 2014 and February 2015 (Table). Eight of 65 fecal samples (K14-362–K14-374) collected on December 2014, one of the 50 fecal samples (N15-99) collected on February 2015, one of the 17 swab samples from healthy common teals (KU-12) collected on January 2015, and one of the 13 swab samples from healthy mallards (KU3-2) collected on February 2015 were positive for influenza A virus by egg inoculation and matrix (M) gene real-time reverse transcription PCR performed as described (8). The hosts of the influenza A virus–positive fecal samples were identified as mandarin ducks, greater white-fronted geese, and mallards by DNA barcoding techniques, as described (7). Full-genome sequencing was performed by next-generation sequencing using the Ion Torrent Personal Genome Machine system (Thermo Fisher Scientific, Grand Island, NY, USA) (online Technical Appendix 1, <http://wwwnc.cdc.gov/EID/article/22/3/15-1006-Techapp1.pdf>). The viruses were subtyped as H5N8 by using a BLAST search, and the multibasic cleavage site of the hemagglutinin (HA) gene (PLRERRRKR/GLF) was detected.

For phylogenetic analysis, we constructed a maximum-likelihood tree in MEGA6 software (<http://www.megasoftware.net>) using the Hasegawa-Kishino-Yano (HKY) model. A median-joining phylogenetic network was constructed by using NETWORK version 4.613 (www.fluxus-engineering.com), and Bayesian analysis was performed by using BEAST version 1.8.1 (<http://beast.bio.ed.ac.uk>). A maximum clade credibility tree was generated for each dataset by using TreeAnnotator in BEAST (online Technical Appendix 1).

Each genome segment of 11 HPAI (H5N8) viruses shared high nucleotide sequence identities ranging from 99.1% to 100%: polymerase basic protein 2, 99.4%–100%; polymerase basic protein 1, 99.3%–100%; polymerase acidic protein, 99.5%–100%; HA, 99.1%–100%;

Author affiliations: Konkuk University, Seoul, South Korea (J.-H. Kwon, J.-Y. Noh, S.-S. Yuk, T.-O. Erdene-Ochir, W.-T. Hong, J.-H. Jeong, S. Jeong, G.-B. Gwon, C.-S. Song); US Department of Agriculture, Athens, Georgia, USA (D.H. Lee, D.E. Swayne)

DOI: <http://dx.doi.org/10.3201/eid2103.151006>

¹These authors contributed equally to this article.

Table. HPAI (H5N8) isolates and total wild bird samples collected in South Korea, December 2014–February 2015*

Collection date	Location	Sample type	No. HPAI (H5N8) positive/no. total	Host†	Strain
2014					
Sep 26	36°44' N, 127°07' E	Feces	0/55		
Sep 27	36°37' N, 126°21' E	Feces	0/110		
Nov 6	36°37' N, 126°21' E	Feces	0/335		(1 LPAI)
Nov 7	36°44' E, 127°07' E	Feces	0/105		
Nov 22	36°44' N, 127°07' E	Feces	0/260		(3 LPAI)
Dec 24	36°44' N, E127°07' E	Feces	8/65	Mandarin duck	K14-363-1
				Greater white-fronted goose	K14-366-1
					K14-367-1
					K14-367-4
					K14-369-3
					K14-371-4
					K14-372-2
					K14-374-1
2015					
Jan 22	36°47' N, 127°03' E	Swab	1/17	Common teal	KU-12
Jan 29	35°18' N, 128°40' E	Swab	1/13	Mallard	KU3-2
			0/30	Northern pintail	(1 LPAI)
Feb 6	37°32' N, 127°01' E	Feces	1/50	Mallard	N15-99
Feb 11	36°42' N, 126°27' E	Swab	0/14	Mallard	
Feb 25	37°23' N, 129°14' E	Swab	0/13	Black-tailed gull	
Mar 16	35°53' N, 127°01' E	Swab	0/15	Eurasian wigeon	
Total			11/1,082		

*HPAI, highly pathogenic influenza virus; LPAI, low pathogenicity avian influenza; bold, period of study.

†The hosts of the HPAI-positive fecal samples were identified by using DNA barcoding techniques as described (7).

nucleoprotein, 99.6%–100%; neuraminidase, 99.2%–100%; M protein, 99.4%–100%; and nonstructural protein, 99.2%–100%. Phylogenetic analysis showed that the 4 different subtype H5N8 virus clusters, icA 1–3 and the South Korea poultry farm cluster, most likely evolved from H5N8 virus identified from South Korea in early 2014. All H5N8 isolates collected in South Korea during winter 2014–15 identified in this study clustered with isolates from Japan, including the A/chicken/Miyazaki/7/2014 strain, and were characterized as subgroup icA3. Isolates obtained from South Korea poultry farms in late 2014 were phylogenetically distinct from isolates in other subgroups (Figure 1; online Technical Appendix 1 Figures 1, 2).

Group A (H5N8) viruses have been detected on South Korea poultry farms since the first outbreak in January 2014, including during the summer season. A second wave of the HPAI (H5N8) outbreak started in September 2014. Although the growing HPAI outbreak in September 2014 coincided with the fall migration of migratory waterfowl, phylogenetic analyses suggest that the HPAI (H5N8) viruses detected on South Korean poultry farms in late 2014 are not related to the icA3 viruses carried by wild waterfowl but have instead evolved from viruses circulating on poultry farms or among resident wild birds in South Korea since early 2014.

By the beginning of the fall 2014 migration of migratory waterfowl, new subgroups of H5N8 viruses (icA1, icA2, icA3) were detected in wintering sites of migratory waterfowl, including South Korea and Japan, in late 2014 and early 2015 (5,9). The icA1 subgroup is composed

of HPAI (H5N8) viruses from Europe, South Korea, and Japan, whereas the icA2 subgroup is composed of HPAI (H5N8) viruses from North America, Taiwan, and Japan and the icA3 subgroup is composed of HPAI (H5N8) viruses isolated in South Korea and Japan. Markov chain Monte Carlo analyses showed that the substitution rates estimated for HPAI (H5N8) viruses identified from South Korea are 9.23×10^{-3} (95% highest posterior density range 7.43×10^{-3} to 1.11×10^{-2}) nt substitutions/site/year, which is higher than previous estimates for the HA gene of H5N1 viruses from China from 1996 through 2012 (4.378×10^{-3} nt substitutions/site/year) (10). The interval estimated from most recent common ancestor of the icA3 cluster from South Korea and Japan was 0.44 years (95% highest posterior density range 0.33–0.55 months, corresponds to August 2014) (Figure 2, <http://wwwnc.cdc.gov/EID/article/22/3/15-1006-F2.htm>).

Conclusions

These results suggest that HPAI (H5N8) viruses circulated in wild bird populations and evolved into subgroups during the breeding season. Detection of subtype H5N8 viruses in healthy wild birds (12,13; this study) and subclinical infection with viral shedding among migratory waterfowl experimentally infected with HPAI (H5N8) viruses (11) support the theory of long-term circulation of HPAI (H5N8) viruses in wild bird population.

This study also found that subtype icA3 viruses, derived from HPAI (H5N8) viruses from South Korea and reintroduced by migratory waterfowl, were genetically distinct from the HPAI (H5N8) viruses that continued to circulate




Figure 1. Median-joining phylogenetic network of highly pathogenic avian influenza A(H5N8) virus isolates identified in South Korea during 2014–2015 showing relationships with other virus isolates. The median-joining network was constructed from the hemagglutinin gene and includes all the most parsimonious trees linking the sequences. Each unique sequence is represented by a circle sized relative to its frequency in the dataset. Branch length is proportional to the number of mutations. Isolates are colored according to the origin of the sample: red inner circle, poultry farm isolates; yellow inner circle, wild bird isolates. Red asterisks indicate isolates from South Korea and blue asterisks indicate isolates from Japan identified during December 2014–February 2015.

in poultry farms. In the previous 4 HPAI (H5N8) virus outbreaks in South Korea and Japan, migratory waterfowl were identified as the source of HPAI outbreaks (14,15); however, related HPAI viruses were not reintroduced into South Korea and Japan after the initial outbreak season. The phylogenetic analysis described here shows that HPAI (H5N8) viruses isolated from migratory wild birds in the winter of 2014–15 are phylogenetically distinct from isolates from South Korean poultry farms. HPAI (H5N8) viruses thus independently evolved in wild bird populations and poultry farms in South Korea until late 2014.

Our results indicate that HPAI (H5N8) viruses have been circulating in wild waterfowl population since early 2014. Enhanced global active surveillance is needed to monitor the spread of these viruses through wild birds. Such efforts could clarify the epidemiology of HPAI virus and facilitate early recognition of novel genotypes.

GISAID (Global Initiative on Sharing All Influenza Data) acknowledgment tables for laboratory contributions are shown in online Technical Appendix 2 (<http://wwwnc.cdc.gov/EID/article/22/3/15-1006-Techapp2.xlsx>).

This work was funded by the Korea Institute of Animal Disease Management Technology Development program (grant no. 313016032WT011), the Ministry of Agriculture, Food and Rural Affairs, Republic of Korea; and was partially funded by KIST open research program (project no. 2E25720-15-P030).

Mr. Kwon is a Ph.D. candidate at Konkuk University, Seoul, South Korea. His primary research interest is the epidemiology of highly pathogenic avian influenza viruses in wild birds. Dr. Lee is a postdoctoral researcher at the Southeast Poultry Research Laboratory, USDA Agricultural Research Service, Athens, Georgia, USA. His research interests include molecular epidemiology and host–pathogen interaction of avian influenza viruses.

References

1. Sonnerberg S, Webby RJ, Webster RG. Natural history of highly pathogenic avian influenza H5N1. *Virus Res.* 2013;178:63–77. <http://dx.doi.org/10.1016/j.virusres.2013.05.009>
2. Lee YJ, Kang HM, Lee EK, Song BM, Jeong J, Kwon YK, et al. Novel reassortant influenza A(H5N8) viruses, South Korea, 2014. *Emerg Infect Dis.* 2014;20:1087–9. <http://dx.doi.org/10.3201/eid2006.140233>
3. Kanehira K, Uchida Y, Takemae N, Hikono H, Tsunekuni R, Saito T. Characterization of an H5N8 influenza A virus isolated from chickens during an outbreak of severe avian influenza in Japan in April 2014. *Arch Virol.* 2015;160:1629–43. <http://dx.doi.org/10.1007/s00705-015-2428-9>
4. Fan S, Zhou L, Wu D, Gao X, Pei E, Wang T, et al. A novel highly pathogenic H5N8 avian influenza virus isolated from a wild duck in China. *Influenza Other Respir Viruses.* 2014;8:646–53. <http://dx.doi.org/10.1111/irv.12289>
5. Lee DH, Torchetti MK, Winker K, Ip HS, Song CS, Swayne DE. Intercontinental spread of Asian-origin H5N8 to North America through Beringia by migratory birds. *J Virol.* 2015
6. Verhagen JH, Herfst S, Fouchier RA. Infectious disease. How a virus travels the world. *Science.* 2015;347:616–7. <http://dx.doi.org/10.1126/science.aaa6724>


7. Lee DH, Lee HJ, Lee YJ, Kang HM, Jeong OM, Kim MC, et al. DNA barcoding techniques for avian influenza virus surveillance in migratory bird habitats. *J Wildl Dis.* 2010;46:649–54. <http://dx.doi.org/10.7589/0090-3558-46.2.649>
8. Spackman E, Senne DA, Bulaga LL, Myers TJ, Perdue ML, Garber LP, et al. Development of real-time RT-PCR for the detection of avian influenza virus. *Avian Dis.* 2003;47 (Suppl):1079–82. <http://dx.doi.org/10.1637/0005-2086-47.s3.1079>
9. Ozawa M, Matsuu A, Tokorozaki K, Horie M, Masatani T, Nakagawa H, Okuya K, Kawabata T, Toda S, et al. Genetic diversity of highly pathogenic H5N8 avian influenza viruses at a single overwinter site of migratory birds in Japan, 2014/15. *Euro Surveill.* 2015;20:21132. <http://dx.doi.org/10.2807/1560-7917.ES2015.20.20.21132>
10. Wei K, Chen Y, Xie D. Genome-scale evolution and phylodynamics of H5N1 influenza virus in China during 1996–2012. *Vet Microbiol.* 2013;167:383–93. <http://dx.doi.org/10.1016/j.vetmic.2013.09.030>
11. Kang HM, Lee EK, Song BM, Jeong J, Choi JG, Jeong J, et al. Novel reassortant influenza A(H5N8) viruses among inoculated domestic and wild ducks, South Korea, 2014. *Emerg Infect Dis.* 2015;21:298–304. <http://dx.doi.org/10.3201/eid2102.141268>
12. Adlhoeh C, Gossner C, Koch G, Brown I, Bouwstra R, Verdonck F, et al. Comparing introduction to Europe of highly pathogenic avian influenza viruses A(H5N8) in 2014 and A(H5N1) in 2005. *Euro Surveill.* 2014;19:20996. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.50.20996>
13. Jeong J, Kang HM, Lee EK, Song BM, Kwon YK, Kim HR, et al. Highly pathogenic avian influenza virus (H5N8) in domestic poultry and its relationship with migratory birds in South Korea during 2014. *Vet Microbiol.* 2014;173:249–57. <http://dx.doi.org/10.1016/j.vetmic.2014.08.002>
14. Kim HR, Lee YJ, Park CK, Oem JK, Lee OS, Kang HM, et al. Highly pathogenic avian influenza (H5N1) outbreaks in wild birds and poultry, South Korea. *Emerg Infect Dis.* 2012;18:480–3. <http://dx.doi.org/10.3201/1803.111490>
15. Lee DH, Park JK, Youn HN, Lee YN, Lim TH, Kim MS, et al. Surveillance and isolation of HPAI H5N1 from wild Mandarin Ducks (*Aix galericulata*). *J Wildl Dis.* 2011;47:994–8. <http://dx.doi.org/10.7589/0090-3558-47.4.994>

Address for correspondence: Chang-Seon Song, Avian Disease Laboratory, College of Veterinary Medicine, Konkuk University, 1 Hwayang-dong, Gwangjin-gu, Seoul, 143-701, South Korea; email: songcs@konkuk.ac.kr

July 2015: Malaria

Including:

- Disseminated Infections with *Talaromyces marneffe* in Non-AIDS Patients Given Monoclonal Antibodies against CD20 and Kinase Inhibitors
- Macacine Herpesvirus 1 in Long-Tailed Macaques, Malaysia, 2009–2011
- Malaria Prevalence among Young Infants in Different Transmission Settings, Africa
- Transdermal Diagnosis of Malaria Using Vapor Nanobubbles
- Lack of Transmission among Close Contacts of Patient with Case of Middle East Respiratory Syndrome Imported into the United States, 2014
- Monitoring of Ebola Virus Makona Evolution through Establishment of Advanced Genomic Capability in Liberia
- Parechovirus Genotype 3 Outbreak among Infants, New South Wales, Australia, 2013–2014
- MERS-CoV in Upper Respiratory Tract and Lungs of Dromedary Camels, Saudi Arabia, 2013–2014
- Assessment of Arbovirus Surveillance 13 Years after Introduction of West Nile Virus, United States
- Results from the National *Legionella* Outbreak Detection Program, the Netherlands, 2002–2012



<http://wwwnc.cdc.gov/eid/articles/issue/21/07/table-of-contents>