

possibility that dromedaries may not be the ultimate natural reservoir (i.e., the long-term host of a pathogen of an infectious disease). Topography (i.e., mountain chains) may limit camel movements from the Middle East or Africa to Central Asia, although such interchange certainly occurred centuries ago as a consequence of the silk-trade routes through southern Kazakhstan. The only known recent imports to Kazakhstan are dromedaries (Arvana breed), brought from Turkmenistan for cross-breeding with Bactrians to improve milk production (8). The findings that MERS-CoV is not universally endemic in dromedaries raises the hypothesis that certain species of bats or some other animal, the environment, or both, may constitute a maintenance community and be the true natural reservoir of MERS-CoV and that the virus spills over to camels and is maintained within camels for varying periods of time. Further studies on the epidemiology of MERS-CoV infection among camelids from central Asia are warranted.

This work was supported by a research grant from the US National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services (contract no. HHSN272201500006C) and a commissioned grant from the Health and Medical Research Fund, Food and Health Bureau, Government of the Hong Kong Special Administrative Region.

References

1. Reusken CB, Haagmans BL, Müller MA, Gutierrez C, Godeke GJ, Meyer B, et al. Middle East respiratory syndrome coronavirus neutralising serum antibodies in dromedary camels: a comparative serological study. *Lancet Infect Dis*. 2013;13:859–66. [http://dx.doi.org/10.1016/S1473-3099\(13\)70164-6](http://dx.doi.org/10.1016/S1473-3099(13)70164-6)
2. Perera RA, Wang P, Gomaa MR, El-Shesheny R, Kandeil A, Bagato O, et al. Seroepidemiology for MERS coronavirus using microneutralisation and pseudoparticle virus neutralisation assays reveal a high prevalence of antibody in dromedary camels in Egypt, June 2013. *Euro Surveill*. 2013;18:pii=20574. <http://dx.doi.org/10.2807/1560-7917.ES2013.18.36.20574>
3. Memish ZA, Cotten M, Meyer B, Watson SJ, Alshahafi AJ, Al Rabeeah AA, et al. Human infection with MERS coronavirus after exposure to infected camels, Saudi Arabia, 2013. *Emerg Infect Dis J*. 2014;20:1012–5.
4. Reusken CB, Messadi L, Feyisa A, Ularumu H, Godeke GJ, Danmarwa A, et al. Geographic distribution of MERS coronavirus among dromedary camels, Africa. *Emerg Infect Dis*. 2014; 20:1370–4. <http://dx.doi.org/10.3201/eid2008.140590>
5. Chan RW, Hemida MG, Kayali G, Chu DK, Poon LL, Alnaeem A, et al. Tropism and replication of Middle East respiratory syndrome coronavirus from dromedary camels in the human respiratory tract: an in-vitro and ex-vivo study. *Lancet Respir Med*. 2014;2:813–22. [http://dx.doi.org/10.1016/S2213-2600\(14\)70158-4](http://dx.doi.org/10.1016/S2213-2600(14)70158-4)
6. Chan SM, Damdinjav B, Perera RA, Chu DK, Khishgee B, Enkhbold B, et al. Absence of MERS-coronavirus in Bactrian camels, southern Mongolia, November 2014. *Emerg Infect Dis*. 2015;21:1269–71. <http://dx.doi.org/10.3201/eid2107.150178>
7. World Organisation for Animal Health. Infection with coronavirus in camels, Iran [cited 2015 Oct 11]. http://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?page_refer=MapFullEventReport&reportid=16411
8. Knoll EM, Burger P. The encounter between Bactrian and dromedary camels in Central Asia. In: Faye B, Konuspayeva G, editors. *Camels in Asia and North Africa: interdisciplinary perspectives on their past and present significance*. Vienna (Austria): Austrian Academy of Sciences Press. 2012. p. 27–33, photos p. 248–250.
9. Hemida MG, Perera RA, Al Jassim RA, Kayali G, Siu LY, Wang P, et al. Seroepidemiology of Middle East respiratory syndrome (MERS) coronavirus in Saudi Arabia (1993) and Australia (2014) and characterisation of assay specificity. *Euro Surveill*. 2014;19:pii=20828. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.23.20828>
10. Cramer G, Durr PA, Barr J, Yu M, Graham K, Williams OJ, et al. Absence of MERS-CoV antibodies in feral camels in Australia: implications for the pathogen's origin and spread. *One Health*. 2015;1:76–82.

Address for correspondence: Eve Miguel, CIRAD, AGIRs, Avenue Agropolis, 34398 Montpellier CEDEX 5, France; email: eve.miguel@cirad.fr

Novel Reassortant Avian Influenza A(H5N1) Virus in Human, Southern Vietnam, 2014

Ikuyo Takayama,¹ Nguyen Trung Hieu,¹ Masayuki Shirakura, Mina Nakauchi, Seiichiro Fujisaki, Hitoshi Takahashi, Shiho Nagata, Nguyen Thanh Long, Takato Odagiri, Masato Tashiro, Tsutomu Kageyama

Author affiliations: National Institute of Infectious Diseases, Tokyo, Japan (I. Takayama, M. Shirakura, M. Nakauchi, S. Fujisaki, H. Takahashi, S. Nagata, T. Odagiri, M. Tashiro, T. Kageyama); Pasteur Institute, Ho Chi Minh City, Vietnam (N.T. Hieu, N.T. Long)

DOI: <http://dx.doi.org/10.3201/eid2203.151360>

To the Editor: The first case of human infection with highly pathogenic avian influenza A(H5N1) virus in Vietnam was reported in December 2003 (1), and >120 human cases were confirmed through 2013, with a high case-fatality rate (2). In 2013, clade 2.3.2.1a/c H5N1 viruses circulated widely in poultry across the country, although clade 1.1.1/1.1.2 H5N1 viruses predominated in poultry from the Mekong Delta region to central Vietnam (3,4).

In 2014, two cases of human infection with A(H5N1) virus were identified in southern Vietnam. One case was associated with a clade 1.1.2 reassortant virus, A/Vietnam/14012902/2014 (Global Initiative on Sharing

¹These first authors contributed equally to this article.

All Influenza Data [GISAID; <http://www.gisaid.org>] accession nos. EPI624919–EPI624926), which had been previously detected in Cambodia and Vietnam (5,6). We isolated the virus from the other case, performed phylogenetic analysis to identify the clade of this virus, and identified a novel virus that had undergone gene reassortment.

The case-patient was a 52-year-old man who lived in Binh Phuoc Province (140 km northeast of Ho Chi Minh City). On January 11, 2014, he experienced mild fever and general fatigue; high fever developed on January 13. He was hospitalized with dyspnea on January 16 and died 2 days later. He was not given antiviral drug treatment. Dead poultry infected with H5N1 viruses were found scattered near his house during January 1–16, and he buried his 2 dead chickens on January 5. H5N1 virus infection was detected in the patient's throat swab specimen by real-time reverse transcription PCR at the Pasteur Institute in Ho Chi Minh City. Virus was isolated by inoculating the throat swab specimens into 10-day-old embryonated chicken eggs; the resulting isolate, A/Vietnam/14011801/2014 (GISAID accession nos. EPI624911–EPI624918), then underwent gene sequencing. The 8 viral genes were amplified with SuperScript III Reverse Transcriptase Kit (Fisher Scientific, Pittsburg, PA, USA) and Phusion High-Fidelity DNA Polymerase (New England BioLabs, Ipswich, MA, USA) with specific paired primers, according to the manufacturer's instructions, and sequenced on an ABI 3730 automated sequencer with Big-Dye Terminator Cycle Sequencing reagents (Applied Biosystems, Foster City, CA, USA). Whole genome sequence was determined.

By gene sequencing analysis, A/Vietnam/14011801/2014 was found to have the multibasic cleavage site of hemagglutinin (HA) protein, which indicates highly pathogenic avian influenza A(H5N1) viruses, and was shown to predict binding specificity to an avian α 2,3 sialic acid receptor. The neuraminidase gene possessed no amino acid substitutions associated with decreased antiviral activity, nor did the virus have amino acid substitutions associated with increased adaptation, virulence, infectivity, or transmissibility in mammalian hosts, including the E627K and D701N mutations in polymerase basic protein 2 (7).

Phylogenetic analyses of the 8 viral genes of A/Vietnam/14011801/2014 were performed by using databases (GISAID and the Influenza Virus Resource, National Center for Biotechnology Information, Bethesda, MD, USA; <http://www.ncbi.nlm.nih.gov/genomes/FLU/FLU.html>) that contained complete sequences of viral genomes belonging to clades 1.1.1, 1.1.2, and 2.3.2.1 a/b/c, most of which were collected in Vietnam, particularly after 2012. Neighbor-joining and Kimura 2-parameter methods were implemented by using MEGA version 5.0 software (<http://www.megasoftware.net>). Reliability of the phylogenetic

analysis was tested by using 1,000 bootstrap replications. Lineages of the HA gene were defined by using previously described criteria (8). Lineages of the other 7 genes were defined by using criteria and nomenclature of Nguyen et al. (9).

The HA of A/Vietnam/14011801/2014 belonged to clade 2.3.2.1c (online Technical Appendix Figure, panel A, <http://wwwnc.cdc.gov/EID/article/23/3/15-1360-Techapp.pdf>). The neuraminidase, polymerase basic proteins 1 and 2, and polymerase acid protein genes of this virus were also derived from respective lineages of ancestor clade 2.3.2.1c (online Technical Appendix Figure, panels B–E). However, nucleoprotein, matrix, and nonstructural genes were classified as lineages of ancestor clade 2.3.2.1a (online Technical Appendix Figure, panels F–H) and differed from the gene lineages of almost all clade 2.3.2.1c viruses isolated from poultry in Vietnam. As reported in the Influenza Virus Resource, 2 viruses collected in Vietnam in December 2013 (A/muscovy duck/Long An/43/2013 and A/muscovy duck/Long An/46/2013) were similar reassortant viruses of clade 2.3.2.1c (Figure). However, the ancestor of the nonstructural gene lineage of these 2 viruses is clade 2.3.2.1c, which differs from A/Vietnam/14011801/2014.




Figure. Novel reassortant virus (A/Vietnam/14011801/2014) identified in a human case of influenza A(H5N1) virus infection in Vietnam, 2014. Ancestry of genes is denoted in the hemagglutinin clades. Arrows indicate genes that differ from the gene lineages of original clade 2.3.2.1c viruses.

The differences indicate that A/Vietnam/14011801/2014 is a novel reassortant virus between clades 2.3.2.1a and 2.3.2.1c, between clades 1.1.2 and 2.3.2.1c, or both (Figure). This novel reassortant virus has not been reported in poultry in Vietnam, although novel reassortants between clade 1.1.2 and clade 2.3.2.1a viruses have been detected in Vietnam since 2013 (i.e., A/Vietnam/VP13-28H/2013, GISAID accession nos. EPI624927–EPI624934; and A/Vietnam/14012902/2014) (6). These novel reassortment viruses were first identified in human, animal, and environmental samples in Cambodia in 2013 (5). Other novel gene reassortments in clade 2.3.2.1 viruses have been previously reported (10), and new clade 2.3.4.4 viruses have been observed in Vietnam since 2014.

As multiple clade viruses co-circulate, reassortment events occur frequently in Vietnam. Continuous surveillance of avian influenza A(H5N1) viruses, not only in humans but also in poultry and wild birds, is needed for infection control measures during epidemics of these viruses.

This study was partially supported by Grants-in-Aid for Emerging and Reemerging Infectious Diseases from the Ministry of Health, Labour and Welfare, Japan.

References

- Tran TH, Nguyen TL, Nguyen TD, Luong TS, Pham PM, Nguyen V, et al. Avian influenza A (H5N1) in 10 patients in Vietnam. *N Engl J Med*. 2004;350:1179–88. <http://dx.doi.org/10.1056/NEJMoa040419>
- World Health Organization. Cumulative number of confirmed human cases for avian influenza A(H5N1) reported to WHO, 2003–2014 [cited 2015 Jan 6]. http://www.who.int/influenza/human_animal_interface/EN_GIP_20141002CumulativeNumberH5N1cases.pdf?ua=1
- Nguyen DT, Bryant JE, Davis CT, Nguyen LV, Pham LT, Loth L, et al. Prevalence and distribution of avian influenza A(H5N1) virus clade variants in live bird markets of Vietnam, 2011–2013. *Avian Dis*. 2014;58:599–608. <http://dx.doi.org/10.1637/10814-030814-Reg>
- Lee EK, Kang HM, Kim KI, Choi JG, To TL, Nguyen TD, et al. Genetic evolution of H5 highly pathogenic avian influenza virus in domestic poultry in Vietnam between 2011 and 2013. *Poult Sci*. 2015;94:650–61. <http://dx.doi.org/10.3382/ps/pev036>
- Rith S, Davis CT, Duong V, Sar B, Horm SV, Chin S, et al. Identification of molecular markers associated with alteration of receptor-binding specificity in a novel genotype of highly pathogenic avian influenza A(H5N1) viruses detected in Cambodia in 2013. *J Virol*. 2014;88:13897–909. <http://dx.doi.org/10.1128/JVI.01887-14>
- Thor SW, Nguyen H, Balish A, Hoang AN, Gustin KM, Nhung PT, et al. Detection and characterization of clade 1 reassortant H5N1 viruses isolated from human cases in Vietnam during 2013. *PLoS ONE*. 2015;10:e0133867. <http://dx.doi.org/10.1371/journal.pone.0133867>
- Le QM, Sakai-Tagawa Y, Ozawa M, Ito M, Kawaoka Y. Selection of H5N1 influenza virus PB2 during replication in humans. *J Virol*. 2009;83:5278–81. <http://dx.doi.org/10.1128/JVI.00063-09>
- World Health Organization/World Organization for Animal Health/Food and Agriculture Organization H5N1 Evolution Working Group. Revised and updated nomenclature for highly pathogenic avian influenza A (H5N1) viruses. *Influenza Other Respi Viruses*. 2014;8:384–8. <http://dx.doi.org/10.1111/irv.12230>
- Nguyen T, Rivailler P, Davis CT, Hoa do T, Balish A, Dang NH, et al. Evolution of highly pathogenic avian influenza (H5N1) virus populations in Vietnam between 2007 and 2010. *Virology*. 2012;432:405–16. <http://dx.doi.org/10.1016/j.virol.2012.06.021>
- Creanga A, Thi Nguyen D, Gerloff N, Thi Do H, Balish A, Dang Nguyen H, et al. Emergence of multiple clade 2.3.2.1 influenza A (H5N1) virus subgroups in Vietnam and detection of novel reassortants. *Virology*. 2013;444:12–20. <http://dx.doi.org/10.1016/j.virol.2013.06.005>

Address for correspondence: Tsutomu Kageyama, Influenza Virus Research Center, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashimurayama-shi, Tokyo 208-0011, Japan; email: tkage@nih.go.jp

***Mycobacterium arupense* as an Emerging Cause of Tenosynovitis**

Fiorella Krapp Lopez, Madeline Miley, Babafemi Taiwo

Author affiliations: Northwestern University, Chicago, Illinois, USA (F. Krapp Lopez, B. Taiwo); Northwestern Memorial Hospital, Chicago (M. Miley)

DOI: <http://dx.doi.org/10.3201/eid2203.151749>

To the Editor: *Mycobacterium arupense* was identified in 2006 as a novel species within the *M. terrae* complex with close similarity to *M. nonchromogenicum* (1). Since then, 8 cases describing clinically notable disease have been published (2–8), including 5 cases of tenosynovitis. We report *M. arupense* tenosynovitis in an immunocompromised person who received the selective interleukin (IL) 1 β -inhibitor canakinumab.

In July 2014, a 62-year-old man sought treatment at the emergency department, Northwestern Memorial Hospital (Chicago, Illinois, USA), after 1 week of pain and swelling in the right hand. During the previous 5 years, he had received multiple immunomodulatory drugs for treatment of natural killer cell deficiency, hyper-IL-6 syndrome, recurrent polychondritis, and Sweet syndrome. His medications were prednisone (42.5 mg/d), intravenous immunoglobulin (400 mg/kg monthly), and subcutaneous canakinumab (180 mg every 8 weeks, which began 3 weeks before onset of symptoms).

His first symptom was a tender red nodule on the right palm that increased in size and became extremely tender over the following week (Figure, panels A, B). He did not recall any trauma and denied fever or chills. No improvement was seen after he received oral linezolid for 5 days. A

Novel Reassortant Avian Influenza A(H5N1) Virus in Human, Southern Vietnam, 2014

Technical Appendix


A

HA gene




B


NA gene



C
PB2 gene




D
PB1 gene




E

PA gene




F


NP gene



G
M gene



H
NS gene



Technical Appendix Figure. Phylogenetic analyses of the genes of the avian influenza A(H5N1) virus described in this study: A) hemagglutinin (HA) gene; B) neuraminidase (NA) gene; C) polymerase basic (PB) 2 gene; D) PB1 gene; E) polymerase acid (PA) protein gene; F) nucleoprotein (NP) gene; G) matrix (M) gene; and H) nonstructural (NS) gene. Trees were constructed by using the neighbor-joining method. The evolutionary distances were computed by using the Kimura 2-parameter method. Bootstrap values were calculated from 1,000 replicates and values >60% are shown next to branches. Sequences of the A(H5N1) virus isolated in this study are in bold font and inside a box. Viruses are colored on the basis of their hemagglutinin clade: 1.1.1 in light blue; 1.1.2 in green; 2.3.2.1a in red; 2.3.2.1b in purple; and 2.3.2.1c in dark blue. Scale bars indicate nucleotide substitutions per site.