

case we present suggests that local transmission of *L. loa* and prevalence of loiasis on the island may be higher than previously thought.

Acknowledgment

We thank Robert Cheke for his advice and assistance with this manuscript.

Dr. Priest is an infectious diseases clinician and Medical Director for Infection Prevention and Antimicrobial Stewardship for Novant Health, an integrated health care system. His interests include clinical care of patients with infectious diseases, antimicrobial stewardship, and infection prevention.

Dr. Nutman is deputy chief and head of the Helminth Immunology Section and Clinical Parasitology Section of the Laboratory of Parasitic Diseases at the National Institute of Allergy and Infectious Diseases, National Institutes of Health. His major research interest is the immune responses in parasitic helminth infections (primarily the filarial infections) and their regulation.

References

1. Metzger WG, Mordmüller B. *Loa loa*—does it deserve to be neglected? *Lancet Infect Dis*. 2014;14:353–7. [http://dx.doi.org/10.1016/S1473-3099\(13\)70263-9](http://dx.doi.org/10.1016/S1473-3099(13)70263-9)
2. Bockarie MJ, Kelly-Hope LA, Rebollo M, Molyneux DH. Preventive chemotherapy as a strategy for elimination of neglected tropical parasitic diseases: endgame challenges. *Philos Trans R Soc Lond B Biol Sci*. 2013;368:20120144. <http://dx.doi.org/10.1098/rstb.2012.0144>
3. Boussinesq M, Gardon J, Gardon-Wendel N, Chippaux JP. Clinical picture, epidemiology and outcome of *Loa*-associated serious adverse events related to mass ivermectin treatment of onchocerciasis in Cameroon. *Filaria J*. 2003;2(Suppl 1):S4. <http://dx.doi.org/10.1186/1475-2883-2-S1-S4>
4. Fink DL, Kamgno J, Nutman TB. Rapid molecular assays for specific detection and quantitation of *Loa loa* microfilaremia. *PLoS Negl Trop Dis*. 2011;5:e1299. <http://dx.doi.org/10.1371/journal.pntd.0001299>
5. Lal RB, Ottesen EA. Enhanced diagnostic specificity in human filariasis by IgG4 antibody assessment. *J Infect Dis*. 1988;158:1034–7. <http://dx.doi.org/10.1093/infdis/158.5.1034>
6. Burbelo PD, Ramanathan R, Klion AD, Iadarola MJ, Nutman TB. Rapid, novel, specific, high-throughput assay for diagnosis of *Loa loa* infection. *J Clin Microbiol*. 2008;46:2298–304. <http://dx.doi.org/10.1128/JCM.00490-08>
7. Cheke RA, Mas J, Chainey JE. Potential vectors of loiasis and other tabanids on the island of Bioko, Equatorial Guinea. *Med Vet Entomol*. 2003;17:221–3. <http://dx.doi.org/10.1046/j.1365-2915.2003.00436.x>
8. Moya L, Herrador Z, Ta-Tang TH, Rubio JM, Perteguer MJ, Hernandez-González A, et al. Evidence for suppression of onchocerciasis transmission in Bioko Island, Equatorial Guinea. *PLoS Negl Trop Dis*. 2016;10:e0004829. <http://dx.doi.org/10.1371/journal.pntd.0004829>
9. Nutman TB, Miller KD, Mulligan M, Ottesen EA. *Loa loa* infection in temporary residents of endemic regions: recognition of a hyperresponsive syndrome with characteristic clinical manifestations. *J Infect Dis*. 1986;154:10–8. <http://dx.doi.org/10.1093/infdis/154.1.10>
10. Herrick JA, Metenou S, Makiya MA, Taylor-Williams CA, Law MA, Klion AD, et al. Eosinophil-associated processes underlie differences in clinical presentation of loiasis between temporary residents and those indigenous to *Loa*-endemic areas. *Clin Infect Dis*. 2015;60:55–63. <http://dx.doi.org/10.1093/cid/ciu723>

Address for correspondence: David H. Priest, 1381 Westgate Center Dr, Winston-Salem, NC 27103, USA; email: dhpriest@novanthealth.org

Invasive Infections with Multidrug-Resistant Yeast *Candida auris*, Colombia

Soraya E. Morales-López,
Claudia M. Parra-Giraldo,
Andrés Ceballos-Garzón, Heidys P. Martínez,
Gerson J. Rodríguez, Carlos A. Álvarez-Moreno,
José Y. Rodríguez

Author affiliations: Universidad Popular del Cesar, Valledupar, Colombia (S.E. Morales-López, H.P. Martínez); Pontificia Universidad Javeriana, Bogotá, Colombia (C.M. Parra-Giraldo, A. Ceballos-Garzón); Centro de Investigaciones Microbiológicas del Cesar (CIMCE), Valledupar, Colombia (G.J. Rodríguez, J.Y. Rodríguez); Clínica Laura Daniela, Valledupar (G.J. Rodríguez, J.Y. Rodríguez); Universidad Nacional de Colombia, Bogotá (C.A. Álvarez-Moreno); Clínica Universitaria Colombia, Colsanitas, Colombia (C.A. Álvarez-Moreno)

DOI: <http://dx.doi.org/10.3201/eid2301.161497>

Candida auris is an emerging multidrug-resistant fungus that causes a wide range of symptoms. We report finding 17 cases of *C. auris* infection that were originally misclassified but correctly identified 27.5 days later on average. Patients with a delayed diagnosis of *C. auris* had a 30-day mortality rate of 35.2%.

Candida auris is an emerging multidrug-resistant fungus that causes a wide range of infections that are sometimes associated with high mortality rates (1–4). *C. auris* was first isolated in Japan and described as a new species in 2009 (5). In 2011, it was described as a cause of fungemia in South Korea (4) and was later isolated from patients in India (2), South Africa (6), Kuwait (3), and Venezuela (1).

We report 17 clinical isolates of *C. auris* recovered from 17 patients hospitalized in 6 institutions in the northern region of Colombia from February through July 2016. We reviewed patient medical records; analyzed microbiological, demographic, and clinical variables; and evaluated the mortality rate 30 days after yeast isolation. The initial

pathogen identification was made with the method available at each institution: VITEK 2 system (bioMérieux, Marcy l'Étoile, France); Phoenix (Becton Dickinson, Franklin Lakes, New Jersey, USA); MicroScan AutoSCAN 4 and MicroScan Walkaway (Beckman Coulter, Brea, California, USA); and API Candida (bioMérieux) (Table). Given the unusual prevalence of *C. haemulonii* and discordance in the micromorphologic characteristics of some isolates, we cultured strains in CHROMagar Candida medium (CHROMagar Candida, Paris, France) and identified them by using matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry (Bruker Daltonik, Bremen, Germany). All the isolates showed pink colonies on CHROMagar Candida medium and were identified as *C. auris* by MALDI-TOF mass spectrometry (scores >2.0).

We tested yeast isolates for in vitro susceptibility by VITEK cards (caspofungin, micafungin, fluconazole, voriconazole, and amphotericin B). Additionally, we performed the agar diffusion method using Etest strips (bioMérieux, France) to amphotericin B according to the manufacturer's instructions. *C. parapsilosis* ATCC 22019 was used as the control. The range, mode, MIC₅₀, and MIC₉₀ were calculated.

Of the 17 patients, 9 were men (52.9%); age range was 0–77 years (median 36 years). Fifteen (88.2%) were hospitalized in intensive care units and 2 in medical wards; no patients were transferred between hospitals.

Blood samples from 13 (76.4%) patients showed fungemia; for the remaining 4, *C. auris* was isolated from peritoneal fluid, cerebrospinal fluid, bone, or urine (Table). Most patients had a central venous catheter (n = 16, 94.1%), a urinary catheter (n = 15, 88.2%), and mechanical ventilation (n = 10, 58.8%). Additionally, some had risk factors described previously for candidemia: erythrocyte transfusion (n = 12, 70.5%), parenteral nutrition (n = 8, 47%), abdominal surgery (n = 7, 41.1%), hemodialysis (n = 5, 29.4%), diabetes (n = 3, 17.6%), pancreatitis (n = 2, 11.7%), cancer (n = 2, 11.7%), and HIV infection (n = 1, 5.8%).

The average number of days from hospitalization to isolation of *C. auris* was 27.5 days (SD ± 19.9 days). Before the isolation of *C. auris*, 15 (88.2%) and 12 (70.6%) patients received broad-spectrum antimicrobial therapy and antifungal therapy, respectively. Of the latter, 8 received fluconazole, 2 caspofungin, 1 fluconazole and caspofungin, and 1 fluconazole, caspofungin, and anidulafungin (given at a different time periods) (Table). The time from isolation of *C. auris* to the initial application of effective antifungal therapy averaged 3.7 days. The 30-day mortality rate in all patients and in those who had fungemia was 35.2% and 38.4%, respectively.

C. auris is phylogenetically related to *C. haemulonii*, for which it is often mistaken with the methods currently used for identification (2–6). Our isolates were originally

Table. Identification and antifungal susceptibilities of *Candida auris* clinical isolates of six hospitals, northern region of Colombia, 2016*

Isolate ID	Hospital no.	Specimen origin	Biochemical identification (system)	Pre-AFT	VITEK cards					Etest/AMB	
					FLC	MCF	CAS	VRC	AMB	24 h	48 h
001	1	Blood	<i>C. haemulonii</i> (VITEK)	None	16	0.12	<0.25	0.25	8	0.75	1
002	1	CSF	<i>C. tropicalis</i> (MicroScan Walkaway)	CAS	16	0.12	<0.25	<0.12	8	0.75	1
003	1	Blood	<i>C. famata</i> (API Candida)	FLC	16	<0.06	<0.25	<0.12	8	1	1
004	5	Blood	<i>C. haemulonii</i> (Phoenix)	FLC	16	0.12	<0.25	0.25	>16	1	1.5
005	2	Blood	<i>C. haemulonii</i> (VITEK)	FLC	>64	0.25	0.5	2	>16	1	1.5
006	4	Blood	<i>C. haemulonii</i> (VITEK)	None	>64	0.12	<0.25	0.5	8	1	1
007	3	Peritoneal fluid	<i>C. albicans</i> (MicroScan autoSCAN)	FLC	16	0.12	<0.25	0.25	8	1	1.5
008	2	Blood	<i>C. haemulonii</i> (VITEK)	FLC	16	0.12	<0.25	0.25	8	0.38	0.75
009	1	Blood	<i>C. tropicalis</i> (MicroScan Walkaway)/ <i>C. famata</i> (API Candida)	FLC, CAS	32	0.12	<0.25	0.25	8	2	3
010	5	Bone	<i>C. haemulonii</i> (VITEK)	FLC, AFG, CAS	16	0.12	<0.25	0.25	8	0.75	1
011	6	Urine	<i>C. haemulonii</i> (Phoenix)	None	32	0.12	<0.25	0.25	8	1.5	1.5
012	3	Blood	<i>C. albicans</i> (MicroScan AutoSCAN)	FLC	32	0.12	<0.25	1	8	1	2
013	3	Blood	<i>C. haemulonii</i> (VITEK)	None	32	0.12	<0.25	1	8	0.75	2
014	3	Blood	<i>C. haemulonii</i> (VITEK)	FLC	>64	0.12	<0.25	2	8	1.5	1.5
015	2	Blood	<i>C. haemulonii</i> (VITEK)	None	32	0.12	<0.25	0.25	8	0.75	2
016	2	Blood	<i>C. haemulonii</i> (VITEK)	FLC	>64	0.12	<0.25	2	>16	1	2
017	4	Blood	<i>C. haemulonii</i> (VITEK)	CAS	>64	0.12	<0.25	2	>16	2	4
MIC (mg/L) range					16 to >64	<0.06 to 0.25	<0.25 to 0.5	<0.12 to 2	8 to >16	0.38 to 2	0.75 to 4
MIC ₅₀					32	0.12	<0.25	0.25	8	1	1.5
MIC ₉₀					>64	0.12	<0.25	2	>16	2	2

*AFG, anidulafungin; AMB, amphotericin B; CAS, caspofungin; CSF, cerebrospinal fluid; FLC, fluconazole; ID, identification; MCF, micafungin; MIC₅₀, minimal inhibitory concentration for 50% of yeast; MIC₉₀, minimal inhibitory concentration for 90% of yeast; Pre-AFT, use of antifungal therapy before the isolation of *C. auris*; VRC, voriconazole.

misidentified as *C. haemulonii*, *C. famata*, *C. albicans*, or *C. tropicalis*, depending on the method used in the hospital. The identification of isolates by MALDI-TOF mass spectrometry has also been described in the literature as an adequate and fast method for identifying *C. auris* (7).

Because the Clinical and Laboratory Standards Institute does not currently provide breakpoints for *C. auris*, no categorical interpretation of results is available; thus, only the MICs obtained for antifungal drugs tested in our study were indicated (Table). Although misleading, elevated MICs of amphotericin B by VITEK card have been previously described (7); this study also found discrepancies with Etest strips, which could lead to the selection of inappropriate therapy if only 1 method is used.

The presence of *C. auris* in these patients has clinical and epidemiologic implications, considering the associated mortality rate confirmed in this report and the absence of sufficient technology in clinical laboratories both to confirm their identification and to carry out testing for antifungal susceptibility. The lack of suitable diagnostics complicates patient treatment and changes on the empiric treatment of invasive *Candida* spp. infections are needed.

Our data contributes to the knowledge of the epidemiology of this species at a regional level. Although we had already reported *Candida* spp. in Colombia (8), no information regarding these species on the Caribbean coast is available. Given the association of *Candida* spp. with outbreaks in hospitals, according to the Centers for Disease Control and Prevention, it is necessary to further strengthen measures for fungal infection control to prevent possible spread.

Acknowledgments

We thank Laura Marcela Ramirez and Juan Sebastian Monroy for their technical support and Heidy Pinzón, Kelin Esquea, Liliana Gómez, Alba Mosquera, Yiceth Acosta, Yenixa Fuentes, Leandro Narváez, Luis Rodrigo Ramírez, and Sindy Jiménez for providing the isolates.

Ms. Morales-López is a bacteriologist and doctoral candidate in the Microbiology program at the Universidad de Buenos Aires, Argentina. She works as an associate teacher of medical microbiology at Universidad Popular del Cesar in Valledupar, Colombia. Her areas of interest are medical microbiology and antimicrobial drug resistance.

References

1. Calvo B, Melo ASA, Perozo-Mena A, Hernandez M, Francisco EC, Hagen F, et al. First report of *Candida auris* in America: clinical and microbiological aspects of 18 episodes of candidemia. *J Infect*. 2016;73:369–74. <http://dx.doi.org/10.1016/j.jinf.2016.07.008>
2. Chowdhary A, Sharma C, Duggal S, Agarwal K, Prakash A, Singh PK, et al. New clonal strain of *Candida auris*, Delhi, India. *Emerg Infect Dis*. 2013;19:1670–3. <http://dx.doi.org/10.3201/eid1910.130393>
3. Emara M, Ahmad S, Khan Z, Joseph L, Al-Obaid I, Purohit P, et al. *Candida auris* candidemia in Kuwait, 2014. *Emerg Infect Dis*. 2015;21:1091–2. <http://dx.doi.org/10.3201/eid2106.150270>
4. Lee WG, Shin JH, Uh Y, Kang MG, Kim SH, Park KH, et al. First three reported cases of nosocomial fungemia caused by *Candida auris*. *J Clin Microbiol*. 2011;49:3139–42. <http://dx.doi.org/10.1128/JCM.00319-11>
5. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiol Immunol*. 2009;53:41–4. <http://dx.doi.org/10.1111/j.1348-0421.2008.00083.x>
6. Magobo RE, Corcoran C, Seetharam S, Govender NP. *Candida auris*—associated candidemia, South Africa. *Emerg Infect Dis*. 2014;20:1250–1. <http://dx.doi.org/10.3201/eid2007.131765>
7. Kathuria S, Singh PK, Sharma C, Prakash A, Masih A, Kumar A, et al. Multidrug-resistant *Candida auris* misidentified as *Candida haemulonii*: characterization by matrix-assisted laser desorption/ionization-time of flight mass spectrometry and DNA sequencing and its antifungal susceptibility profile variability by Vitek 2, CLSI broth microdilution, and Etest method. *J Clin Microbiol*. 2015;53:1823–30. <http://dx.doi.org/10.1128/JCM.00367-15>
8. Parra C, LePape P, Ceballos A, Cortes G, Alvarez-Moreno C, Valderrama S, et al. Performance of MALDI-TOF MS for the identification of emerging yeast of hospital patients, species distribution, in a third level hospital Bogotá-Colombia. In: Abstracts of the 26th European Congress of Clinical Microbiology and Infectious Diseases; Amsterdam; 2016 Apr 9–12. Abstract 5556. Basel: European Congress of Clinical Microbiology and Infectious Diseases; 2016.

Address for correspondence: José Y. Rodríguez, Centro de Investigaciones, Microbiológicas del Cesar (CIMCE), Calle 16c no 19d-14 Barrio Dangond, Valledupar, Colombia; email: jyrodriquezq@gmail.com

Zika Virus Knowledge among Pregnant Women Who Were in Areas with Active Transmission

Kate Whittemore, Anna Tate, Alex Illescas, Alhaji Saffa, Austin Collins, Jay K. Varma, Neil M. Vora

Author affiliations: New York City Department of Health and Mental Hygiene, New York, New York, USA (K. Whittemore, A. Tate, A. Illescas, A. Saffa, A. Collins, J.K. Varma, N.M. Vora); Centers for Disease Control and Prevention, Atlanta, Georgia, USA (J.K. Varma, N.M. Vora)

DOI: <http://dx.doi.org/10.3201/eid2301.161614>

We surveyed women in New York, New York, USA, who were in areas with active Zika virus transmission while pregnant. Of 99 women who were US residents, 30 were