

Significant Decrease in Pertactin-Deficient *Bordetella pertussis* Isolates, Japan

Yukihiro Hiramatsu, Yusuke Miyaji,
Nao Otsuka, Yoshichika Arakawa,
Keigo Shibayama, Kazunari Kamachi

Prevalence of pertactin-lacking *Bordetella pertussis* isolates has been observed worldwide. In Japan, however, we found that the frequency of pertactin-deficient isolates in 2014–2016 (8%) was significantly lower than the frequency in 2005–2007 (41%), 2008–2010 (35%), and 2011–2013 (25%). This reduction was closely associated with changes in genotypes.

Bordetella pertussis, a highly communicable, gram-negative coccobacillus, is the etiologic agent of pertussis (whooping cough), an acute respiratory infection that leads to severe illness in children. Vaccination is the most effective method for preventing and controlling pertussis. In Japan, acellular pertussis vaccines (ACVs) were introduced in 1981. Pertussis toxin and filamentous hemagglutinin derived from *B. pertussis* are the major antigens in ACVs in Japan, and certain ACVs also contain pertactin and fimbriae (*f*). Pertactin is believed to play a role in adherence to human epithelial cells (2); however, *B. pertussis* isolates that lack pertactin production have been identified in several countries where ACVs have been introduced (3–7). In Japan, pertactin-deficient isolates have increased significantly since the early 2000s, resulting in a high prevalence of these isolates (5,8). Recent studies have demonstrated that pertactin-deficient strains could colonize the respiratory tract more effectively than pertactin-producing strains in ACV-vaccinated mice (9,10). Supporting these results, an epidemiologic study suggested that ACV-vaccinated persons are more susceptible to pertactin-deficient strains than to pertactin-producing strains (11). These reports imply that pertactin-deficient strains have increased fitness in humans who have been vaccinated with ACVs and that their expansion may reduce the effectiveness of ACVs. We assessed trends in the frequency of pertactin-deficient isolates in Japan and further investigated their genotypes using multilocus variable-number tandem-repeat analysis (MLVA).

Author affiliations: National Institute of Infectious Diseases, Tokyo, Japan (Y. Hiramatsu, Y. Miyaji, N. Otsuka, K. Shibayama, K. Kamachi); St. Marianna University School of Medicine, Kawasaki, Japan (Y. Miyaji); Nagoya University Graduate School of Medicine, Nagoya, Japan (Y. Arakawa)

DOI: <http://dx.doi.org/10.3201/eid2304.161575>

The Study

We studied 232 *B. pertussis* clinical isolates collected from January 2005 through June 2016 in Japan. All isolates were derived from epidemiologically unrelated cases of pertussis. Pertactin production and MLVA types (MTs) of 111 isolates collected during 2005–2012 were previously determined by immunoblotting and MLVA typing, respectively (5,8). For our study, we extended these analyses to additional isolates collected during 2008–2016 ($n = 121$).

We determined the temporal trend in the frequency of pertactin-deficient isolates by 3-year periods (Figure, panel A). Percentages were 41% in 2005–2007 ($n = 39$ isolates), 35% in 2008–2010 ($n = 43$), 25% in 2011–2013 ($n = 97$), and 8% in 2014–2016 ($n = 53$). A significant decrease in the frequency of pertactin-deficient isolates was observed from 2005–2007 to 2014–2016 ($p < 0.05$ by Fisher exact test).

Among the 232 *B. pertussis* isolates, 25 MTs were identified; MT27 and MT186 isolates were predominant, and other MT isolates were found at low frequencies (online Technical Appendix Table 1, <http://wwwnc.cdc.gov/EID/article/23/4/16-1575-Techapp1.pdf>). The frequency of MT27 isolates increased significantly over time (Figure 1, panel B): 28% in 2005–2007, 44% in 2008–2010, 70% in 2011–2013, and 77% in 2014–2016. In contrast, the frequency of MT186 isolates decreased: 31% in 2005–2007, 44% in 2008–2010, 21% in 2011–2013, and 6% in 2014–2016. We also observed a substitution of the major genotype in the *B. pertussis* population from MT186 to MT27.

Of 59 pertactin-deficient *B. pertussis* isolates collected during 2005–2016, 45 (76.3%) were MT186 isolates, whereas 2 (3.4%) represented MT27 and 12 (20.3%) other MT isolates (MT194, MT224–226, MT314–316) (Table). Notably, 45 (83.3%) of 54 MT186 isolates were pertactin-deficient, whereas only 2 (1.4%) of 139 MT27 isolates were pertactin-deficient. This finding indicates that pertactin-deficient isolates predominate among the MT186 strain but are rare among the MT27 strain.

We previously showed that pertactin-deficient isolates in Japan were generated by 2 different mutations: an 84-bp deletion of the *prn* gene signal sequence (Δ SS) and an IS481 insertion at nucleotide position 1598 in *prn* (1598–1599::IS481) (5). Thus, to confirm the molecular basis for the loss of pertactin production, pertactin-deficient isolates ($n = 26$) that were newly identified in this study underwent PCR screening with 2 primer sets (online Technical Appendix Table 2). We summarized the molecular mechanisms of loss of pertactin production in 59 pertactin-deficient isolates




Figure. Temporal trends in the frequency of pertactin-deficient isolates and changes in multi-locus variable-number tandem repeat analysis types (MTs) in the *Bordetella pertussis* population in Japan. Pertactin production (A) and MTs (B) were analyzed for 232 *B. pertussis* isolates collected from January 2005 through June 2016. The frequencies of pertactin-deficient isolates and 2 major MTs (MT27 and MT186) are shown by time period. For convenience, minor MTs (MT194, MT224–226, and MT314–316) are included as “others.”

(Table); the Δ SS mutation was detected in 43 (72.9%) MT186 isolates and in 11 (18.6%) other MT isolates. In contrast, the 1598–1599::IS481 mutation was detected in 2 (3.4%) MT186 isolates and 1 other MT isolate (1.7%, MT226). Two MT27 isolates (BP394 and BP533) do not have either of these mutations. Instead, BP533 isolates (1.7%) have an IS481 insertion at nucleotide position 245 (245–246::IS481; GenBank accession no. KC445198), and BP394 (1.7%) isolates exhibit transcriptional down-regulation of *prn* gene expression (online Technical Appendix Figure 1).

Conclusions

The expansion of pertactin-deficient *B. pertussis* isolates has been reported worldwide (3–8). However, we observed a significant decrease in pertactin-deficient isolates within the *B. pertussis* population in Japan, caused by a genotypic replacement from the pertactin-deficient MT186 strain to the pertactin-producing MT27 strain.

The most likely explanation for the prevalence of pertactin-deficient strains is vaccine-driven strain evolution, because pertactin is a component of ACVs. No pertactin-deficient isolates have been detected in Denmark, where an ACV that does not contain pertactin is used, and pertactin-deficient strains exhibit a selective advantage against ACV-induced immunity (7,9–11). In Japan, 5 brands of the diphtheria–tetanus–acellular pertussis vaccine (DTaP) had been used to control pertussis for many years. These DTaPs had different formulations of components; only 3 contained the pertactin antigen (1). When the DTaP vaccine was replaced, however, 2 brands of combined DTaP–inactivated poliovirus (DTaP-IPV) vaccine that did not contain pertactin were introduced in November 2012. Therefore, most Japanese children ≤ 4 years of age do not have immunity to pertactin; consequently, the selective pressure for pertactin-deficient strains in the host environment has recently been reduced. This effect may be responsible for the recent decline in

pertactin-deficient isolates and further supports the hypothesis that pertactin-deficient strains are selected on the basis of host immunity to pertactin. Notably, a new brand of DTaP-IPV vaccine containing pertactin was also introduced in December 2015 in Japan. If the hypothesis of vaccine-driven evolution is correct, pertactin-deficient isolates should increase again in Japan in the near future. Thus, continued surveillance of pertactin-deficient isolates is of particular value.

We demonstrated that genotypic replacement from MT186 to MT27 has taken place among recent *B. pertussis* isolates in Japan: MT27 is a triple-locus variant of MT186. MT186 strains carry the pertussis-toxin promoter allele *ptxP1*, whereas MT27 strains carry the allele *ptxP3* (8). *B. pertussis* strains carrying *ptxP3* (i.e., MT27) produce more of several virulence factors than do *ptxP1* (i.e., MT186) strains (12,13). The population of MT27 strains carrying *ptxP3* has increased worldwide (14,15), although a low frequency of *ptxP1* isolates was observed in Japan (8), suggesting that MT27 strains are associated with the recent pertussis resurgence. It is possible, therefore, that the genotypic replacement in the *B. pertussis* population may have resulted from the expansion of the more virulent *ptxP3* (i.e., MT27) strains. In addition, given that pertactin-deficient MT27 isolates are rare, this genotypic replacement

Table. Molecular mechanisms of loss of pertactin production in 59 pertactin-deficient *Bordetella pertussis* isolates, Japan, 2005–2016*

Reason for loss of pertactin	No. (%) MTs		
	MT27	MT186	Others†
Δ SS	0	43 (72.9)	11 (18.6)
1598–1599::IS481	0	2 (3.4)	1 (1.7)
245–246::IS481	1 (1.7)	0	0
Transcriptional downregulation	1 (1.7)	0	0
Total	2 (3.4)	45 (76.3)	12 (20.3)

*MT, types determined by using multilocus variable-number tandem-repeat analysis.

†For convenience, minor MTs (MT194, MT224–226, and MT314–316) are included in this category.

may have contributed to the recent decrease in pertactin-deficient isolates in Japan.

In Japan, most pertactin-deficient isolates carry a deletion of the *prn* signal sequence (Δ SS), which has been found primarily in MT186 isolates carrying the *prn1* allele (online Technical Appendix Table 1). In other countries, a common *prn* mutation includes an insertion of IS481 into the *prn2* allele (4,6,7). In this study, we identified 2 pertactin-deficient MT27 isolates carrying the *prn2* allele, due to the IS481 insertion (245–246::IS481) and the transcriptional down-regulation of the *prn* gene. These pertactin-deficient isolates were previously identified in Europe and the United States (6,7). One possible explanation for the appearance of pertactin-deficient MT27 isolates is that they were imported from other countries.

This study was supported by the following funding sources: Research on Emerging and Re-emerging Infectious Diseases from the Ministry of Health, Labor, and Welfare of Japan; Research Program on Emerging and Re-emerging Infectious Diseases from the Japan Agency for Medical Research and Development.

Dr. Hiramatsu is a research scientist at the National Institute of Infectious Diseases in Tokyo, Japan. His research interests focus on the epidemiology, pathogenesis, and development of pertussis vaccines.

References

- Okada K, Komiya T, Yamamoto A, Takahashi M, Kamachi K, Nakano T, et al. Safe and effective booster immunization using DTaP in teenagers. *Vaccine*. 2010;28:7626–33. <http://dx.doi.org/10.1016/j.vaccine.2010.09.050>
- Leininger E, Roberts M, Kenimer JG, Charles IG, Fairweather N, Novotny P, et al. Pertactin, an Arg-Gly-Asp-containing *Bordetella pertussis* surface protein that promotes adherence of mammalian cells. *Proc Natl Acad Sci U S A*. 1991;88:345–9. <http://dx.doi.org/10.1073/pnas.88.2.345>
- Bouchez V, Brun D, Cantinelli T, Dore G, Njamkepo E, Guiso N. First report and detailed characterization of *B. pertussis* isolates not expressing pertussis toxin or pertactin. *Vaccine*. 2009;27:6034–41. <http://dx.doi.org/10.1016/j.vaccine.2009.07.074>
- Lam C, Octavia S, Ricafort L, Sintchenko V, Gilbert GL, Wood N, et al. Rapid increase in pertactin-deficient *Bordetella pertussis* isolates, Australia. *Emerg Infect Dis*. 2014;20:626–33. <http://dx.doi.org/10.3201/eid2004.131478>
- Otsuka N, Han HJ, Toyozumi-Ajisaka H, Nakamura Y, Arakawa Y, Shibayama K, et al. Prevalence and genetic characterization of pertactin-deficient *Bordetella pertussis* in Japan. *PLoS One*. 2012;7:e31985. <http://dx.doi.org/10.1371/journal.pone.0031985>
- Pawloski LC, Queenan AM, Cassiday PK, Lynch AS, Harrison MJ, Shang W, et al. Prevalence and molecular characterization of pertactin-deficient *Bordetella pertussis* in the United States. *Clin Vaccine Immunol*. 2014;21:119–25. <http://dx.doi.org/10.1128/CVI.00717-13>
- Zeddeman A, van Gent M, Heuvelman CJ, van der Heide HG, Bart MJ, Advani A, et al. Investigations into the emergence of pertactin-deficient *Bordetella pertussis* isolates in six European countries, 1996 to 2012. *Euro Surveill*. 2014;19:20881. <http://dx.doi.org/10.2807/1560-7917.ES2014.19.33.20881>
- Miyaji Y, Otsuka N, Toyozumi-Ajisaka H, Shibayama K, Kamachi K. Genetic analysis of *Bordetella pertussis* isolates from the 2008–2010 pertussis epidemic in Japan. *PLoS One*. 2013;8:e77165. <http://dx.doi.org/10.1371/journal.pone.0077165>
- Hegerle N, Dore G, Guiso N. Pertactin deficient *Bordetella pertussis* present a better fitness in mice immunized with an acellular pertussis vaccine. *Vaccine*. 2014;32:6597–600. <http://dx.doi.org/10.1016/j.vaccine.2014.09.068>
- Safarchi A, Octavia S, Luu LD, Tay CY, Sintchenko V, Wood N, et al. Pertactin negative *Bordetella pertussis* demonstrates higher fitness under vaccine selection pressure in a mixed infection model. *Vaccine*. 2015;33:6277–81. <http://dx.doi.org/10.1016/j.vaccine.2015.09.064>
- Martin SW, Pawloski L, Williams M, Weening K, DeBolt C, Qin X, et al. Pertactin-negative *Bordetella pertussis* strains: evidence for a possible selective advantage. *Clin Infect Dis*. 2015;60:223–7. <http://dx.doi.org/10.1093/cid/ciu788>
- King AJ, van der Lee S, Mohangoo A, van Gent M, van der Ark A, van de Waterbeemd B. Genome-wide gene expression analysis of *Bordetella pertussis* isolates associated with a resurgence in pertussis: elucidation of factors involved in the increased fitness of epidemic strains. *PLoS One*. 2013;8:e66150. <http://dx.doi.org/10.1371/journal.pone.0066150>
- Moore FR, van Loo IH, van Gent M, He Q, Bart MJ, Heuvelman KJ, et al. *Bordetella pertussis* strains with increased toxin production associated with pertussis resurgence. *Emerg Infect Dis*. 2009;15:1206–13. <http://dx.doi.org/10.3201/eid1508.081511>
- Schmidtke AJ, Boney KO, Martin SW, Skoff TH, Tondella ML, Tatti KM. Population diversity among *Bordetella pertussis* isolates, United States, 1935–2009. *Emerg Infect Dis*. 2012;18:1248–55. <http://dx.doi.org/10.3201/eid1808.120082>
- van Gent M, Heuvelman CJ, van der Heide HG, Hallander HO, Advani A, Guiso N, et al. Analysis of *Bordetella pertussis* clinical isolates circulating in European countries during the period 1998–2012. *Eur J Clin Microbiol Infect Dis*. 2015;34:821–30. <http://dx.doi.org/10.1007/s10096-014-2297-2>

Address for correspondence: Yukihiko Hiramatsu, Department of Bacteriology II, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashimurayama, Tokyo 208-0011, Japan; email: yhira@nih.go.jp

Significant Decrease in Pertactin–Deficient *Bordetella pertussis* Isolates, Japan

Technical Appendix

Technical Appendix Table 1. Characteristics of *Bordetella pertussis* isolates, Japan, 2005–2016*

Isolate	Isolation year	Patient age	Vaccine status	Origin (district)	Pertactin production	<i>prn</i> gene	<i>prn</i> allele	MLVA type
BP285	2005	2 mo	0	Kanto	–	ΔSS	1	186
BP291	2005	1 y	3	Chubu	–	ΔSS	1	186
BP292	2005	1 mo	0	Chubu	–	ΔSS	1	226
BP293	2005	4 mo	1	Chubu	–	ΔSS	1	186
BP299	2005	5 mo	0	Kyusyu	–	ΔSS	1	186
BP302	2005	1 mo	0	Kinki	–	1598–	1	226
						1599::IS481		
BP307	2006	2 mo	0	Chubu	–	1598–	1	186
						1599::IS481		
BP310	2006	11 y	Unknown	Tohoku	–	ΔSS	1	194
BP313	2006	58 y	Unknown	Kanto	–	1598–	1	186
						1599::IS481		
BP322	2006	1 mo	0	Kanto	–	ΔSS	1	186
BP323	2006	1 y	Unknown	Kanto	–	ΔSS	1	186
BP332	2006	35 y	Unknown	Kinki	–	ΔSS	1	194
BP333	2006	30 y	Unknown	Kinki	–	ΔSS	1	194
BP334	2006	0 y	0	Kinki	–	ΔSS	1	194
BP314	2007	3 mo	0	Kanto	–	ΔSS	1	186
BP318	2007	1 y	0	Kanto	–	ΔSS	1	186
BP466	2008	33 y	Unknown	Kanto	–	ΔSS	1	186
BP469	2008	8 y	Unknown	Kanto	–	ΔSS	1	186
BP346	2009	2 mo	0	Chubu	–	ΔSS	1	186
BP351	2009	4 mo	Unknown	Shikoku	–	ΔSS	1	186
BP458	2009	70 y	Unknown	Kanto	–	ΔSS	1	186
BP459	2009	5 mo	Unknown	Kanto	–	ΔSS	1	186
BP460	2009	6 mo	Unknown	Kanto	–	ΔSS	1	186
BP357	2010	12 y	Unknown	Kanto	–	ΔSS	1	186
BP358	2010	10 y	Unknown	Kanto	–	ΔSS	1	186
BP359	2010	Unknown	Unknown	Kanto	–	ΔSS	1	186
BP360	2010	Unknown	Unknown	Kanto	–	ΔSS	1	186
BP361	2010	10 y	Unknown	Kanto	–	ΔSS	1	186
BP363	2010	Unknown	Unknown	Kanto	–	ΔSS	1	186
BP365	2010	Unknown	Unknown	Kanto	–	ΔSS	1	186
BP366	2010	6 y	Unknown	Kanto	–	ΔSS	1	186
BP378	2011	21 y	Unknown	Kyusyu	–	ΔSS	1	186
BP390	2011	4 y	Unknown	Kinki	–	ΔSS	1	316
BP394	2011	5 mo	0	Kyusyu	–	<i>prn</i>	2	27
						(transcriptional down–regulation)		
BP395	2011	6 mo	0	Kinki	–	ΔSS	1	186
BP397	2011	7 y	Unknown	Kinki	–	ΔSS	1	186
BP398	2011	4 mo	0	Kinki	–	ΔSS	1	225
BP400	2011	5 y	Unknown	Kinki	–	ΔSS	1	186
BP401	2011	9 y	4	Kinki	–	ΔSS	1	314
BP403	2011	2 mo	0	Kinki	–	ΔSS	1	315
BP417	2011	35 y	4	Kanto	–	ΔSS	1	186
BP437	2011	1 mo	0	Kinki	–	ΔSS	1	186
BP438	2011	2 mo	0	Kinki	–	ΔSS	1	186
BP451	2011	3 mo	0	Kinki	–	ΔSS	1	186
BP452	2011	7 mo	Unknown	Kinki	–	ΔSS	1	186
BP410	2012	1 mo	0	Tohoku	–	ΔSS	1	186
BP412	2012	26 y	Unknown	Shikoku	–	ΔSS	1	186

Isolate	Isolation year	Patient age	Vaccine status	Origin (district)	Pertactin production	<i>prn</i> gene	<i>prn</i> allele	MLVA type
BP416	2012	3 mo	0	Kanto	-	ΔSS	1	186
BP418	2012	5 y	4	Kanto	-	ΔSS	1	186
BP440	2012	10 y	4	Kinki	-	ΔSS	1	186
BP442	2012	2 y	0	Kinki	-	ΔSS	1	316
BP447	2012	28 y	Unknown	Kyusyu	-	ΔSS	1	186
BP478	2012	5 y	Unknown	Kinki	-	ΔSS	1	186
BP480	2013	2 mo	0	Kinki	-	ΔSS	1	224
BP481	2013	1 mo	0	Kinki	-	ΔSS	1	186
BP510	2015	1 mo	0	Kinki	-	ΔSS	1	186
BP533	2015	1 mo	0	Kinki	-	245-246::IS481	2	27
BP535	2015	3 mo	0	Kyusyu	-	ΔSS	1	186
BP550	2016	7 y	4	Kanto	-	ΔSS	1	186
BP283	2005	9 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP284	2005	1 mo	0	Chubu	+	<i>prn</i>	1	186
BP289	2005	6 mo	0	Kinki	+	<i>prn</i>	2	27
BP290	2005	1 y	0	Chugoku	+	<i>prn</i>	2	27
BP294	2005	10 mo	1	Kyusyu	+	<i>prn</i>	2	27
BP296	2005	6 y	0	Kyusyu	+	<i>prn</i>	2	27
BP297	2005	3 mo	0	Chugoku	+	<i>prn</i>	2	22
BP298	2005	3 mo	0	Kinki	+	<i>prn</i>	2	69
BP300	2005	2 mo	0	Kyusyu	+	<i>prn</i>	2	26
BP301	2005	11 mo	0	Kyusyu	+	<i>prn</i>	2	27
BP303	2005	1 mo	0	Chugoku	+	<i>prn</i>	1	224
BP306	2006	10 y	4	Chubu	+	<i>prn</i>	2	69
BP311	2006	1 y	Unknown	Tohoku	+	<i>prn</i>	1	187
BP335	2006	0 y	Unknown	Kinki	+	<i>prn</i>	1	186
BP312	2007	4 mo	0	Kanto	+	<i>prn</i>	2	27
BP316	2007	53 y	Unknown	Tohoku	+	<i>prn</i>	1	234
BP317	2007	11 y	Unknown	Tohoku	+	<i>prn</i>	1	234
BP324	2007	65 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP327	2007	10 y	Unknown	Shikoku	+	<i>prn</i>	1	229
BP330	2007	6 mo	0	Chubu	+	<i>prn</i>	2	27
BP336	2007	0 y	2	Kinki	+	<i>prn</i>	2	27
BP337	2007	43 y	Unknown	Kinki	+	<i>prn</i>	2	27
BP343	2007	0 y	Unknown	Kanto	+	<i>prn</i>	2	95
BP331	2008	1 y	0	Chubu	+	<i>prn</i>	2	27
BP338	2008	11 y	2	Kinki	+	<i>prn</i>	2	27
BP339	2008	8 y	Unknown	Kinki	+	<i>prn</i>	2	27
BP340	2008	29 y	Unknown	Kinki	+	<i>prn</i>	2	27
BP341	2008	6 y	Unknown	Kinki	+	<i>prn</i>	1	187
BP342	2008	0 y	Unknown	Kinki	+	<i>prn</i>	2	27
BP344	2008	5 y	Unknown	Chubu	+	<i>prn</i>	2	27
BP345	2008	4 y	Unknown	Chubu	+	<i>prn</i>	2	27
BP464	2008	3 mo	0	Kanto	+	<i>prn</i>	2	27
BP465	2008	27 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP467	2008	8 mo	Unknown	Kanto	+	<i>prn</i>	2	27
BP347	2009	3 y	0	Shikoku	+	<i>prn</i>	9	27
BP348	2009	41 y	Unknown	Chubu	+	<i>prn</i>	2	27
BP349	2009	2 mo	0	Kyusyu	+	<i>prn</i>	1	186
BP350	2009	13 y	4	Chubu	+	<i>prn</i>	1	186
BP352	2009	1 mo	0	Shikoku	+	<i>prn</i>	1	34
BP353	2009	12 y	Unknown	Kyusyu	+	<i>prn</i>	2	22
BP354	2009	1 mo	0	Kyusyu	+	<i>prn</i>	1	311
BP355	2009	6 mo	Unknown	Kinki	+	<i>prn</i>	1	27
BP356	2010	2 mo	0	Kinki	+	<i>prn</i>	ND	221
BP362	2010	3 y	Unknown	Kyusyu	+	<i>prn</i>	2	27
BP364	2010	8 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP367	2010	9 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP368	2010	5 mo	Unknown	Kyusyu	+	<i>prn</i>	1	186
BP369	2010	15 y	Unknown	Kyusyu	+	<i>prn</i>	2	27
BP371	2010	1 mo	0	Kyusyu	+	<i>prn</i>	1	186
BP462	2010	6 mo	Unknown	Kanto	+	<i>prn</i>	2	27
BP470	2010	1 mo	0	Kanto	+	<i>prn</i>	2	27
BP376	2011	2 mo	0	Chubu	+	<i>prn</i>	2	27
BP377	2011	1 mo	0	Chugoku	+	<i>prn</i>	2	27
BP380	2011	11 y	Unknown	Kyusyu	+	<i>prn</i>	2	27
BP388	2011	3 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP389	2011	3 mo	1	Kanto	+	<i>prn</i>	2	27

Isolate	Isolation year	Patient age	Vaccine status	Origin (district)	Pertactin production	<i>prn</i> gene	<i>prn</i> allele	MLVA type
BP391	2011	5 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP392	2011	4 mo	Unknown	Kanto	+	<i>prn</i>	ND	27
BP393	2011	11 mo	Unknown	Kyusyu	+	<i>prn</i>	2	27
BP396	2011	1 mo	0	Kinki	+	<i>prn</i>	1	187
BP399	2011	2 y	Unknown	Kinki	+	<i>prn</i>	ND	27
BP402	2011	1 mo	0	Kinki	+	<i>prn</i>	ND	27
BP404	2011	10 y	Unknown	Kyusyu	+	<i>prn</i>	2	27
BP405	2011	14 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP406	2011	1 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP407	2011	6 mo	Unknown	Kyusyu	+	<i>prn</i>	2	27
BP430	2011	1 y	3	Kinki	+	<i>prn</i>	2	27
BP431	2011	2 mo	0	Kinki	+	<i>prn</i>	9	27
BP432	2011	8 y	0	Kinki	+	<i>prn</i>	2	27
BP433	2011	5 mo	0	Kinki	+	<i>prn</i>	2	27
BP436	2011	4 mo	0	Kinki	+	<i>prn</i>	ND	27
BP453	2011	6 mo	0	Kinki	+	<i>prn</i>	1	186
BP463	2011	3 mo	1	Kanto	+	<i>prn</i>	2	27
BP471	2011	2 mo	0	Kanto	+	<i>prn</i>	2	27
BP493	2011	Unknown	Unknown	Kanto	+	<i>prn</i>	ND	27
BP494	2011	0 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP501	2011	Unknown	Unknown	Kinki	+	<i>prn</i>	ND	27
BP408	2012	5 mo	Unknown	Kanto	+	<i>prn</i>	ND	27
BP409	2012	4 mo	Unknown	Tohoku	+	<i>prn</i>	2	27
BP411	2012	17 y	Unknown	Tohoku	+	<i>prn</i>	ND	27
BP413	2012	10 y	4	Kanto	+	<i>prn</i>	2	27
BP414	2012	1 mo	0	Chubu	+	<i>prn</i>	2	27
BP415	2012	8 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP420	2012	4 mo	0	Kanto	+	<i>prn</i>	2	27
BP421	2012	5 mo	Unknown	Kanto	+	<i>prn</i>	ND	27
BP422	2012	8 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP423	2012	8 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP424	2012	9 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP425	2012	7 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP426	2012	8 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP427	2012	4 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP428	2012	10 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP429	2012	12 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP434	2012	7 y	Unknown	Kanto	+	<i>prn</i>	2	27
BP435	2012	7 y	Unknown	Kanto	+	<i>prn</i>	ND	27
BP439	2012	3 mo	Unknown	Kinki	+	<i>prn</i>	2	27
BP441	2012	3 mo	Unknown	Kinki	+	<i>prn</i>	ND	186
BP443	2012	3 mo	Unknown	Kinki	+	<i>prn</i>	2	27
BP444	2012	1 mo	0	Tohoku	+	<i>prn</i>	3	27
BP445	2012	1 mo	0	Tohoku	+	<i>prn</i>	2	27
BP446	2012	2 mo	0	Tohoku	+	<i>prn</i>	2	27
BP448	2012	2 mo	0	Kinki	+	<i>prn</i>	2	27
BP449	2012	4 mo	2	Kinki	+	<i>prn</i>	2	27
BP450	2012	1 mo	0	Kinki	+	<i>prn</i>	2	27
BP454	2012	11 y	Unknown	Tohoku	+	<i>prn</i>	9	27
BP455	2012	1 mo	Unknown	Tohoku	+	<i>prn</i>	2	27
BP456	2012	0 mo	0	Tohoku	+	<i>prn</i>	2	27
BP472	2012	1 mo	0	Kanto	+	<i>prn</i>	2	27
BP477	2012	2 mo	0	Kinki	+	<i>prn</i>	ND	27
BP495	2012	0 y	Unknown	Tohoku	+	<i>prn</i>	ND	27
BP496	2012	Unknown	Unknown	Kanto	+	<i>prn</i>	ND	27
BP513	2012	3 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP514	2012	4 mo	≥1	Kyusyu	+	<i>prn</i>	ND	27
BP515	2012	5 mo	≥1	Kyusyu	+	<i>prn</i>	ND	26
BP538	2012	0 mo	0	Tohoku	+	<i>prn</i>	ND	27
BP473	2013	29 y	≥1	Shikoku	+	<i>prn</i>	ND	27
BP474	2013	31 y	Unknown	Kyusyu	+	<i>prn</i>	ND	186
BP475	2013	8 y	Unknown	Kyusyu	+	<i>prn</i>	ND	28
BP479	2013	10 y	0	Kinki	+	<i>prn</i>	ND	27
BP482	2013	4 mo	Unknown	Kanto	+	<i>prn</i>	ND	27
BP483	2013	52 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP497	2013	3 mo	Unknown	Kanto	+	<i>prn</i>	ND	27
BP498	2013	5 y	Unknown	Tohoku	+	<i>prn</i>	ND	27
BP516	2013	2 mo	0	Kyusyu	+	<i>prn</i>	ND	27


Isolate	Isolation year	Patient age	Vaccine status	Origin (district)	Pertactin production	<i>prn</i> gene	<i>prn</i> allele	MLVA type
BP484	2014	1 mo	0	Kyusyu	+	<i>prn</i>	ND	28
BP485	2014	13 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP486	2014	1 mo	0	Shikoku	+	<i>prn</i>	ND	27
BP487	2014	1 mo	0	Kanto	+	<i>prn</i>	ND	27
BP488	2014	1 mo	0	Chugoku	+	<i>prn</i>	ND	27
BP489	2014	5 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP490	2014	3 mo	0	Chugoku	+	<i>prn</i>	ND	27
BP491	2014	8 y	4	Kinki	+	<i>prn</i>	ND	27
BP499	2014	3 mo	Unknown	Kinki	+	<i>prn</i>	ND	27
BP500	2014	30 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP502	2014	1 mo	0	Kanto	+	<i>prn</i>	ND	27
BP503	2014	1 mo	0	Kanto	+	<i>prn</i>	ND	27
BP505	2014	36 y	Unknown	Kinki	+	<i>prn</i>	ND	27
BP506	2014	10 y	Unknown	Hokkaidou	+	<i>prn</i>	ND	27
BP517	2014	1 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP518	2014	1 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP519	2014	8 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP520	2014	13 y	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP521	2014	6 y	≥1	Kyusyu	+	<i>prn</i>	ND	27
BP522	2014	6 y	≥1	Kyusyu	+	<i>prn</i>	ND	27
BP523	2014	8 y	≥1	Kyusyu	+	<i>prn</i>	ND	27
BP524	2014	4 mo	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP525	2014	1 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP526	2014	2 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP527	2014	2 mo	0	Kyusyu	+	<i>prn</i>	ND	29
BP532	2014	4 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP507	2015	3 mo	Unknown	Kinki	+	<i>prn</i>	ND	78
BP508	2015	1 mo	0	Chubu	+	<i>prn</i>	ND	27
BP509	2015	32 y	Unknown	Kyusyu	+	<i>prn</i>	ND	29
BP511	2015	2 mo	0	Kyusyu	+	<i>prn</i>	ND	96
BP512	2015	8 y	4	Kinki	+	<i>prn</i>	ND	27
BP528	2015	1 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP529	2015	2 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP530	2015	3 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP531	2015	3 mo	Unknown	Kyusyu	+	<i>prn</i>	ND	27
BP534	2015	0 mo	0	Kyusyu	+	<i>prn</i>	ND	27
BP536	2015	3 mo	1	Shikoku	+	<i>prn</i>	ND	27
BP537	2015	2 mo	0	Kinki	+	<i>prn</i>	ND	31
BP539	2015	1 mo	0	Tohoku	+	<i>prn</i>	ND	32
BP540	2015	1 mo	0	Tohoku	+	<i>prn</i>	ND	27
BP541	2015	6 mo	0	Kinki	+	<i>prn</i>	ND	27
BP551	2015	12 y	3	Chubu	+	<i>prn</i>	2	27
BP542	2016	9 y	4	Hokkaidou	+	<i>prn</i>	2	27
BP543	2016	10 y	Unknown	Kinki	+	<i>prn</i>	2	27
BP544	2016	1 mo	0	Kyusyu	+	<i>prn</i>	1	34
BP545	2016	13 y	4	Chubu	+	<i>prn</i>	2	27
BP549	2016	9 y	4	Hokkaidou	+	<i>prn</i>	2	27
BP555	2016	6 mo	3	Chubu	+	<i>prn</i>	2	28
BP556	2016	8 y	4	Kanto	+	<i>prn</i>	2	27

*, negative; +, positive; ND, not determined; MLVA, multilocus variable number tandem repeat analysis.

Technical Appendix Table 2. Primers used in this study of *Bordetella pertussis* isolates, Japan, 2008–2016*

Designation	Primer name	Sequence (5' to 3')
Conventional PCR	SS-defect-F5	CTCTGTCACGCATTGACAAC
	SS-outerR	CTCGGCCGCGGGATTTTCTA
	IS481-combine-F	GCAGACGCCACTAGGTGTGA
	IS481-combine-R	AAAGGTCGCCGCGCTGCCTA
qRT PCR	qprn-F	ATCGTCAAGACCGGTGAGCG
	qprn-R	CTGACGGCCGCTTACCTTGA
	qrecA-F	CCAATGTGGTCGACAAGTCC
	qrecA-R	ATGGCCATTCCTTGTGCTC

*qRT-PCR, quantitative reverse transcription PCR.



Technical Appendix Figure. Lack of *pm* transcript expression in pertactin-deficient *Bordetella pertussis* isolate BP394. The pertactin-deficient isolate was cultured for 3 days on Bordet-Gengou agar. Total RNA was isolated by using the RNeasy Mini Kit (QIAGEN, Hilden, Germany) and treated with RNase-Free DNase (QIAGEN) to degrade contaminating DNA. Total RNA (0.1 μ g) was reverse-transcribed into cDNA by using the PrimeScript RT Master Mix with random hexamers (TaKaRa Bio, Inc., Shiga, Japan). Relative levels of *pm* and *recA* transcripts were determined using SYBR Premix Ex TaqII (TaKaRa Bio) with the ABI PRISM 7500 Fast Real-Time PCR System (Applied Biosystems, Waltham, MA, USA). Quantitative reverse transcription-PCR (qRT-PCR) conditions were 10 s at 95°C, followed by 40 cycles of 95°C for 3 s, and 60°C for 30 s. Primer sets qprn-F/qprn-R and qrecA-F/qrecA-R were used for *pm* and *recA* amplification, respectively (Technical Appendix Table 2). The relative *pm* transcript level was calculated using the $\Delta\Delta$ Ct method and was normalized to that of *recA*. The *recA* transcript was used as an internal control for each sample. Data are presented as -fold- changes in expression compared with

those observed in *B. pertussis* strain Tohama. The mean \pm SDs of results obtained from 3 separate experiments are shown.