

Appendix). The maximum-likelihood analysis using both P4b and DNA polymerase gene fragments showed that FPV-MOZ-608/2016 and FPV-MOZ-980/2016 clustered in clade E with the APV isolated in Hungary in 2011 (TKPV-HU1124/2011) (Figure, panels A and B) (6).

TKPV-HU1124/2011 was isolated from a flock of turkeys vaccinated with a commercial vaccine for FPV, and FPV-MOZ-608/2016 and FPV-MOZ-980/2016 were also obtained from vaccinated chickens. The laying pullets imported from South Africa had already been vaccinated for FPV on day 1 with the commercial fowlpox-vectored infectious laryngotracheitis vaccine and day 17 with the fowlpox-vectored infectious laryngotracheitis/avian encephalomyelitis vaccine. Our data suggest a possible failure of the vaccine to protect against clade E viruses, similar to what has been reported previously for TKPV-HU1124/2011 (6).

The identification of a clade E avipoxvirus in Mozambique requires further investigation to clarify how a virus that has only been reported once found its way to this country. Because the chickens in both infected flocks were purchased from the same pullet reseller who had (for both flocks) imported the birds from South Africa, it is likely that the source of infection was the same. However, the specific source has not been identified. FPVs are known to infect >230 species of wild birds, many of which are migratory (5); thus, introduction through migratory wild birds is a possibility.

Resolution of the full genome of these viruses might provide hints to their origin. The presence of fowlpox disease in birds vaccinated against FPV requires urgent re-evaluation of the vaccine formula and control strategies in Mozambique.

This work was supported through funding from the International Atomic Energy Agency Peaceful Uses Initiative (MOZ/14003) and the African Renaissance project, which is funded by the Republic of South Africa. Part of the field activities were supported by the Kyeema Foundation.

Dr. Mapaco is a researcher at the Central Veterinary Laboratory of the Agrarian Research Institute of Mozambique. His research interests are viral diseases of animals, including zoonoses.

References

1. Weli SC, Tryland M. Avipoxviruses: infection biology and their use as vaccine vectors. *Virology*. 2011;8:49. <http://dx.doi.org/10.1186/1743-422X-8-49>
2. Binns MM, Bourns ME, Tomley FM, Campbell J. Analysis of the fowlpoxvirus gene encoding the 4b core polypeptide and demonstration that it possesses efficient promoter sequences. *Virology*. 1989;170:288–91. [http://dx.doi.org/10.1016/0042-6822\(89\)90380-2](http://dx.doi.org/10.1016/0042-6822(89)90380-2)
3. Gyuranecz M, Foster JT, Dán Á, Ip HS, Egstad KF, Parker PG, et al. Worldwide phylogenetic relationship of avian poxviruses. *J Virol*. 2013;87:4938–51. <http://dx.doi.org/10.1128/JVI.03183-12>
4. Jarmin S, Manvell R, Gough RE, Laidlaw SM, Skinner MA. Avipoxvirus phylogenetics: identification of a PCR length polymorphism that discriminates between the two major clades.

J Gen Virol. 2006;87:2191–201. <http://dx.doi.org/10.1099/vir.0.81738-0>

5. Manarolla G, Pisoni G, Sironi G, Rampin T. Molecular biological characterization of avian poxvirus strains isolated from different avian species. *Vet Microbiol*. 2010;140:1–8. <http://dx.doi.org/10.1016/j.vetmic.2009.07.004>
6. Bányai K, Palya V, Dénes B, Glávits R, Ivanics É, Horváth B, et al. Unique genomic organization of a novel avipoxvirus detected in turkey (*Meleagris gallopavo*). *Infect Genet Evol*. 2015;35:221–9. <http://dx.doi.org/10.1016/j.meegid.2015.08.001>
7. Abdallah FM, Hassanin O. Detection and molecular characterization of avipoxviruses isolated from different avian species in Egypt. *Virus Genes*. 2013;46:63–70. <http://dx.doi.org/10.1007/s11262-012-0821-y>
8. Offerman K, Carulei O, Gous TA, Douglass N, Williamson AL. Phylogenetic and histological variation in avipoxviruses isolated in South Africa. *J Gen Virol*. 2013;94:2338–51. <http://dx.doi.org/10.1099/vir.0.054049-0>
9. Mzula A, Masola SN, Kasanga CJ, Wambura PN. Existence of variant strains fowlpox virus integrated with reticuloendotheliosis virus in its genome in field isolates in Tanzania. *Trop Anim Health Prod*. 2014;46:711–6. <http://dx.doi.org/10.1007/s11250-014-0552-2>

Address for correspondence: William G. Dundon, APHL Joint FAO/IAEA Division, International Atomic Energy Agency Laboratories, Seibersdorf A-2444, Austria; email: w.dundon@iaea.org

Indication of Cross-Species Transmission of Astrovirus Associated with Encephalitis in Sheep and Cattle

Céline L. Boujon, Michel C. Koch, Daniel Wüthrich, Simea Werder, Dennis Jakupovic, Rémy Bruggmann, Torsten Seuberlich

Author affiliation: University of Bern, Bern, Switzerland

DOI: <https://doi.org/10.3201/eid2309.170168>

We report the identification of a neurotropic astrovirus associated with encephalitis in a sheep. This virus is genetically almost identical to an astrovirus recently described in neurologically diseased cattle. The similarity indicates that astroviruses of the same genotype may cause encephalitis in different species.

Astroviruses are nonenveloped viruses with a single stranded, positive-sense RNA genome. They are best known to be associated with gastroenteritis, especially in humans. Recently, reports of these viruses in association

with encephalitis have increased dramatically, with reports of cases in humans (1), mink (2), and cattle (3–6).

The most common causes for viral encephalitis in sheep include maedi-visna virus, Borna disease virus, and rabies virus. In a high proportion of cases of nonsuppurative encephalitis cases (which is indicative of a viral infection) in sheep, however, the etiologic agent remains unknown (7). To investigate that matter, we subjected 3 ovine encephalitis samples from our archives to next-generation sequencing and a bioinformatics pipeline for virus discovery (online Technical Appendix, <https://wwwnc.cdc.gov/EID/article/23/9/17-0168-Techapp1.pdf>). In 1 animal (ID 41669), 1 of the contigs obtained had high similarity (>98%) to bovine astrovirus CH15 (BoAstV-CH15), a virus found recently in 2 cases of nonsuppurative encephalitis in cattle (5). The affected sheep was a 7-year-old Swiss white alpine ewe that was culled for reasons other than human consumption. No other information about the clinical history of the animal was available. Histological diagnosis consisted of severe nonsuppurative meningoencephalitis. Routine diagnostic tests for Borna, rabies, and maedi-visna viruses were all negative.

We used primers based on the BoAstV-CH15 genome sequence (GenBank accession no. KT956903) and the Sanger method to sequence the complete genome of the ovine strain (online Technical Appendix Figure 1), which we named ovine astrovirus CH16 (OvAstV-CH16). The sequence we obtained shared >98% identity with BoAstV-CH15 on the nucleotide and amino acid level (Genbank accession no. KY859988; online Technical Appendix Table 1). The virus reported here is genetically clearly distinct from intestinal OvAstV strains described previously (OvAstV-1 and OvAstV-2; online Technical Appendix Table 1).

A phylogenetic comparison confirmed the close relationship of OvAstV-CH16 with BoAstV-CH15 (5) and BoAstV-BH89/14, another astrovirus detected in association with encephalitis in a cow in Germany (6). Recently, 2 astroviruses were reported in association with encephalitis in sheep in Scotland (OvAstV UK/2013/ewe/lib01454 and UK/2014/lamb/lib01455) (8), and we included their genomic data in the study comparison. All these strains clustered in the same branch of the phylogenetic tree, with >95% amino acid sequence similarity in the viral capsid protein (online Technical Appendix Figure 2) and, therefore, should be considered 1 genotype species (9). When we compared all these viruses more closely on the amino acid level, we were not able to find any sequence variant that could be specifically associated with a tropism for sheep or cattle (online Technical Appendix Table 3).

We then analyzed brain samples of sheep with nonsuppurative encephalitis of unknown etiology (n = 47), which had been identified within the framework of active disease surveillance in Switzerland (7), by RT-PCR specific for

BoAstV-CH15 (online Technical Appendix). None reacted positively, implying a low incidence of OvAstV-CH16 infection associated with encephalitis in the sheep population in Switzerland.

To confirm the presence of OvAstV-CH16 in situ, we used polyclonal antisera targeted at the putative capsid protein of BoAstV-CH15 and tested formalin-fixed, paraffin-embedded brain tissues of sheep 41669 by immunohistochemistry (online Technical Appendix). We observed positive staining of neurons, assessed as such on the basis of morphological criteria, in all brain regions examined (e.g., medulla oblongata, cerebellum, thalamus, hippocampus, cortex, caudate nucleus), in particular in some areas (Figure; online Technical Appendix Figure 3). This finding supports a plausible biological association of OvAstV-CH16 infection and encephalitis in the sheep under investigation and underlines again the close relationship between OvAstV-CH16 and BoAstV-CH15. The identification of similar astroviruses in sheep and cattle with comparable diseases, by different methods and in distinct geographic areas, further strengthens these findings. We consider it unlikely that the ovine cases reported in Scotland and Switzerland are epidemiologically related and speculate that the respective viruses were already geographically widely spread but were undetected until recently, which also seems to be the case for neurotropic astroviruses in cattle.

The importance of the link between astroviruses and encephalitis is increasingly being brought to light, but the factors determining their tropism and neuroinvasion are still unknown. Deeper epidemiologic, genetic, and molecular investigations will help to clarify these aspects of astrovirus pathology. Astroviruses were traditionally considered to be host specific, but in recent years, several reports challenged this assumption; for instance, when human astroviruses were found in fecal samples of primates (10). In such cases, however, effective infection of atypical hosts was never shown. In this study, we demonstrated the presence of the virus in situ, a finding that strengthens the likelihood of such an infectious event. The fact that a virus of the same genotype was described in cattle with similar pathologic findings also challenges this concept of host specificity and suggests that astroviruses can cross the species barrier and, therefore, represent a zoonotic threat as not only a gastroenteric agent but also a potential cause of encephalitis.

Acknowledgments

The authors thank Muriel Fragnière and Camille Monney for excellent technical support, Anna Oevermann for neuropathology expertise, and the Next Generation Sequencing Platform of the University of Bern, which is supervised by Cord Drögemüller, for performing the high-throughput sequencing experiments. Thanks also to L. Boujon for proofreading the article.




Figure. Immunohistochemistry (IHC) for ovine astrovirus CH16 in brain tissues (hippocampus) of a sheep (ID 41669) with encephalitis using 2 polyclonal antisera targeted at the putative capsid protein of bovine astrovirus CH15. A) IHC using antiserum against the conserved region of the capsid protein showing positive staining (box at left; box at right shows area at higher magnification); B) negative control. C) IHC using antiserum against the variable regions of the capsid protein showing positive staining (box at left; box at right shows area at higher magnification); D) negative control. Scale bars indicate 50 μm .

This work was funded in part by the Federal Food Safety and Veterinary Office (grant MON-108), by the Swiss National Science Foundation (grant 31003A_163438), and by the Bangerter-Rhyner-Foundation.

Dr. Boujon is a veterinarian and PhD student at the Vetsuisse Faculty in Bern. Her primary research lies in the study of neurotropic astroviruses in cattle.

References

1. Vu DL, Cordey S, Brito F, Kaiser L. Novel human astroviruses: Novel human diseases? *J Clin Virol.* 2016;82:56–63. <http://dx.doi.org/10.1016/j.jcv.2016.07.004>
2. Blomström AL, Widén F, Hammer AS, Belák S, Berg M. Detection of a novel astrovirus in brain tissue of mink suffering from shaking mink syndrome by use of viral metagenomics. *J Clin Microbiol.* 2010;48:4392–6. <http://dx.doi.org/10.1128/JCM.01040-10>
3. Li L, Diab S, McGraw S, Barr B, Traslavina R, Higgins R, et al. Divergent astrovirus associated with neurologic disease in cattle. *Emerg Infect Dis.* 2013;19:1385–92. <http://dx.doi.org/10.3201/eid1909.130682>
4. Bouzalas IG, Wüthrich D, Walland J, Drögemüller C, Zurbriggen A, Vandevelde M, et al. Neurotropic astrovirus in cattle with nonsuppurative encephalitis in Europe. *J Clin Microbiol.* 2014;52:3318–24. <http://dx.doi.org/10.1128/JCM.01195-14>
5. Seuberlich T, Wüthrich D, Selimovic-Hamza S, Drögemüller C, Oevermann A, Bruggmann R, et al. Identification of a second encephalitis-associated astrovirus in cattle. *Emerg Microbes Infect.* 2016;5:e5. <http://dx.doi.org/10.1038/emi.2016.5>
6. Schlottau K, Schulze C, Bilk S, Hanke D, Höper D, Beer M, et al. Detection of a novel bovine astrovirus in a cow with encephalitis. *Transbound Emerg Dis.* 2016;63:253–9. <http://dx.doi.org/10.1111/tbed.12493>
7. Oevermann A, Botteron C, Seuberlich T, Nicolier A, Friess M, Doherr MG, et al. Neuropathological survey of fallen stock: active surveillance reveals high prevalence of encephalitic listeriosis in small ruminants. *Vet Microbiol.* 2008;130:320–9. <http://dx.doi.org/10.1016/j.vetmic.2008.01.015>
8. Pfaff F, Schlottau K, Scholes S, Courtenay A, Hoffmann B, Höper D, et al. A novel astrovirus associated with encephalitis and ganglionitis in domestic sheep. *Transbound Emerg Dis.* 2017;64:677–82. <http://dx.doi.org/10.1111/tbed.12623>
9. Bosch A, Pintó RM, Guix S. Human astroviruses. *Clin Microbiol Rev.* 2014;27:1048–74. <http://dx.doi.org/10.1128/CMR.00013-14>
10. Karlsson EA, Small CT, Freiden P, Feeroz MM, Matsen FA IV, San S, et al. Non-human primates harbor diverse mammalian and avian astroviruses including those associated with human infections. *PLoS Pathog.* 2015;11:e1005225. <http://dx.doi.org/10.1371/journal.ppat.1005225>

Address for correspondence: Torsten Seuberlich, NeuroCenter, Division of Neurological Sciences, Vetsuisse Faculty, University of Bern, Bremgartenstrasse 109 a, CH-3001 Bern, Switzerland; email: torsten.seuberlich@vetsuisse.unibe.ch

Indication of Cross-Species Transmission of Astrovirus Associated with Encephalitis in Sheep and Cattle

Technical Appendix

Tissue Samples

Sheep tissue samples were collected in the framework of a neuropathological survey in the small ruminant population in Switzerland in 2004 and 2005 (1). Whole brains of sheep and goats that died on farm for unknown reasons were collected according to a standardized procedure. Brains were cut sagittally into two equal halves. One half was stored at -20°C for the purpose of molecular testing. The other half was fixed in formalin, and representative brain regions were paraffin embedded and processed for histopathology. Brain sections were then HE stained and examined for histopathological lesions indicative of neurologic disease. Those samples with lesions consistent with a viral infection of the brain, i.e., a nonsuppurative inflammatory pattern, were further tested for common viral pathogens associated with encephalitis in sheep and goats (small ruminant lentiviruses, Borna virus, rabies virus). Tissues were archived in the Biobank of the Division of Neurologic Sciences, Vetsuisse Faculty, University of Bern, until further analysis in the current study.

Next-Generation Sequencing

DNA and RNA extracts from frozen brain tissues of 3 sheep diagnosed with severe nonsuppurative encephalitis, but with negative test results for the above-mentioned viruses, were processed to next-generation-sequencing libraries separately (2). Libraries were sequenced in an Illumina HiSeq using 1 lane of paired-end 150 bp reads, with yields of 177,146,907–195,143,918 reads for the DNA extracts and 113,160,073–251,062,995 reads for the RNA extracts. The reads of the DNA libraries were mapped to the *Ovis aries* reference genome (version 3.1) using Bowtie 2 (version 2.2.4) (3) and the reads of the RNA libraries were mapped to the *Ovis aries* reference

genome (version 3.1) using STAR (version 2.5.0b) (4). The reads that did not align to the reference genome were quality trimmed using Trimmomatic (version 0.33, option: SLIDINGWINDOW:4:14 MINLEN:127) (5). The trimmed reads were assembled using SPAdes (version 3.9.0, options:–meta -k 21,33,55,77,99,127) (6). The resulting assembly was subsequently analyzed using the virus discovery pipeline described by Wüthrich et al. (7).

Sanger Sequencing

The whole genome length of the virus was Sanger sequenced with primers that were designed on the sequence of BoAstV-CH15. Briefly, RNA was extracted from brain tissue of animal 41669 with TRI reagent (Sigma-Aldrich). Using primers BoAstV 15 2do, BoAstV 15 3do, BoAstV 15 q1do, BoAstV 15 q2do (Technical Appendix Table 3), and oligo(dT), viral RNA was reverse transcribed with ThermoScript Reverse Transcriptase (Life Technologies). Using different combinations of primers recognizing sequences scattered over the whole genome length of BoAstV-CH15, binding on both strands of the cDNA and spaced apart from one another with ~500 nt (the sequence of those primers are available on demand), PCR amplified fragments were directly sequenced in both directions, until the whole genome sequence of the new virus was covered.

Sequence Comparisons and Phylogenetic Analysis

The obtained full-length sequence was compared with full-length sequences of BoAstV-CH15 (8), BoAstV-BH89/14 (9), OvAstV-UK/2013/ewe/lib01454 and OvAstV-UK/2014/lamb/lib01455 (10), and 2 ovine astrovirus strains that were detected in feces samples of sheep (11, 12) (Technical Appendix Table 1 and Technical Appendix Figure 2), using the alignment function of the Geneious software Version 9.1.5 (Biomatters). The phylogenetic tree was constructed from amino acid sequences of the astrovirus capsid proteins by maximum-likelihood using the PHYML plug-in in Geneious with 100 bootstraps.

RT-PCR Screening

Forty-seven brain homogenates of sheep diagnosed with nonsuppurative encephalitis, which were available in our archives from another study (1), were tested by RT-PCR with

primers BoAstV CH15 3fo and BoAstV CH15 3do (8) (Technical Appendix Table 2) using the OneTaq One-Step RT-PCR Kit (New England Biolabs), with the alternative protocol described by the manufacturer. Cycling conditions were as follows: 48°C for 15 min, 94°C for 1 min, 40 cycles of 94°C for 15 sec, 52°C for 30 sec and 68°C for 45 sec, and 68°C for 5 min.

Antibody Production

The methods used to obtain antibodies were the same as those we described previously for another bovine astrovirus, BoAstV-CH13 (13). Briefly, rabbits were immunized with recombinant viral proteins expressed in bacteria and the resulting hyperimmune serum samples were collected and affinity purified. Only diverging elements will be mentioned here. First, because structural proteins are usually produced in greater amounts than nonstructural ones during viral replication, we chose only 2 regions on the corresponding putative open reading frame (ORF), ORF2, similar to ORF2-con and ORF2-var described in the previous study. CH15-ORF2-con corresponds to the conserved region, and N-terminal part of protein Vp90, CH15-ORF2-var to a more variable, more C-terminally located part of this protein (14,15) (Technical Appendix Figure 1). Reverse transcription was performed with oligo(dT). The primers subsequently used to amplify the regions of interest were the following: CH15O2c NheI F and CH15O2c BamHI R for the CH15-ORF2-con region and CH15O2v NheI F and CH15O2v BamHI R for CH15-ORF2-var (Technical Appendix Table 3). Lysis of transformed bacteria was performed using 1 mg/l Lysozyme from chicken egg white (Sigma Aldrich), and buffers used for protein purification consisted in PBS with 1 M urea, the concentration of imidazole used in the binding/washing buffer being 10 mM and 20 mM, respectively. One rabbit was injected subcutaneously with 50–200 µg of each antigen contained in SDS-PAGE gel pieces (LowDose Antiserum protocol).

Immunohistochemistry

For both antisera, a simple screening with 3 different antigen retrieval methods (microwave heating in citrate buffer at pH 6 and pH 9, proteinase K treatment) and dilutions (1:20, 1:50, 1:100) was sufficient to determine the best performing parameters, which were microwave heating in citrate buffer at pH 6 and 1:50 dilution of the primary antibody, the

protocol being otherwise the same as the one described in another of our studies (13). Each method was tested on case 41669 and a negative control sheep which was negative for OvAstV-CH16 by RT-PCR.

References

1. Oevermann A, Botteron C, Seuberlich T, Nicolier A, Friess M, Doherr MG, et al. Neuropathological survey of fallen stock: active surveillance reveals high prevalence of encephalitic listeriosis in small ruminants. *Vet Microbiol.* 2008;130:320–9. [PubMed](#)
<http://dx.doi.org/10.1016/j.vetmic.2008.01.015>
2. Bouzalas IG, Wüthrich D, Walland J, Drögemüller C, Zurbriggen A, Vandeveld M, et al. Neurotropic astrovirus in cattle with nonsuppurative encephalitis in Europe. *J Clin Microbiol.* 2014;52:3318–24. [PubMed](#) <http://dx.doi.org/10.1128/JCM.01195-14>
3. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012;9:357–9. [PubMed](#) <http://dx.doi.org/10.1038/nmeth.1923>
4. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics.* 2013;29:15–21. [PubMed](#)
<http://dx.doi.org/10.1093/bioinformatics/bts635>
5. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014;30:2114–20. [PubMed](#) <http://dx.doi.org/10.1093/bioinformatics/btu170>
6. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 2012;19:455–77. [PubMed](#) <http://dx.doi.org/10.1089/cmb.2012.0021>
7. Wüthrich D, Boujon CL, Truchet L, Selimovic-Hamza S, Oevermann A, Bouzalas IG, et al. Exploring the virome of cattle with non-suppurative encephalitis of unknown etiology by metagenomics. *Virology.* 2016;493:22–30. [PubMed](#) <http://dx.doi.org/10.1016/j.virol.2016.03.009>
8. Seuberlich T, Wüthrich D, Selimovic-Hamza S, Drögemüller C, Oevermann A, Bruggmann R, et al. Identification of a second encephalitis-associated astrovirus in cattle. *Emerg Microbes Infect.* 2016;5:e5. [PubMed](#) <http://dx.doi.org/10.1038/emi.2016.5>
9. Schlottau K, Schulze C, Bilk S, Hanke D, Höper D, Beer M, et al. Detection of a novel bovine astrovirus in a cow with encephalitis. *Transbound Emerg Dis.* 2016;63:253–9. [PubMed](#)
<http://dx.doi.org/10.1111/tbed.12493>

10. Pfaff F, Schlottau K, Scholes S, Courtenay A, Hoffmann B, Höper D, et al. A novel astrovirus associated with encephalitis and ganglionitis in domestic sheep. *Transbound Emerg Dis.* 2017;64:677–82. [PubMed http://dx.doi.org/10.1111/tbed.12623](http://dx.doi.org/10.1111/tbed.12623)
11. Jonassen CM, Jonassen TØ, Sveen TM, Grinde B. Complete genomic sequences of astroviruses from sheep and turkey: comparison with related viruses. *Virus Res.* 2003;91:195–201. [PubMed http://dx.doi.org/10.1016/S0168-1702\(02\)00269-1](http://dx.doi.org/10.1016/S0168-1702(02)00269-1)
12. Reuter G, Pankovics P, Delwart E, Boros Á. Identification of a novel astrovirus in domestic sheep in Hungary. *Arch Virol.* 2012;157:323–7. [PubMed http://dx.doi.org/10.1007/s00705-011-1151-4](http://dx.doi.org/10.1007/s00705-011-1151-4)
13. Boujon CL, Selimovic-Hamza S, Bouzalas I, Seuberlich T. Development and validation of an immunohistochemistry procedure for the detection of a neurotropic bovine astrovirus. *J Virol Methods.* 2017;239:26–33. [PubMed http://dx.doi.org/10.1016/j.jviromet.2016.10.013](http://dx.doi.org/10.1016/j.jviromet.2016.10.013)
14. Wang QH, Kakizawa J, Wen LY, Shimizu M, Nishio O, Fang ZY, et al. Genetic analysis of the capsid region of astroviruses. *J Med Virol.* 2001;64:245–55. [PubMed http://dx.doi.org/10.1002/jmv.1043](http://dx.doi.org/10.1002/jmv.1043)
15. Bass DM, Qiu S. Proteolytic processing of the astrovirus capsid. *J Virol.* 2000;74:1810–4. [PubMed http://dx.doi.org/10.1128/JVI.74.4.1810-1814.2000](http://dx.doi.org/10.1128/JVI.74.4.1810-1814.2000)

Technical Appendix Table 1. Pairwise sequence similarities of neurotropic astrovirus strains in ruminants.

ORF	Nucleotide / amino acid similarity, %						
	BoAstV-BH89/14	BoAstV-CH15	OvAstV-CH16*	OvAstV-1	OvAstV-2	UK/2013/ewe	UK/2014/lamb
<i>ORF 1a</i>							
BoAstV-BH89/14	-	98.6	98.5	72.7	n.a.	98.8	98.9
BoAstV-CH15	91.1	-	98.9	72.2	n.a.	98.6	98.5
OvAstV-CH16*	91.4	98.3	-	72.5	n.a.	98.2	98.3
OvAstV-1	68.0	67.6	67.8	-	n.a.	72.2	72.3
OvAstV-2	n.a.	n.a.	n.a.	n.a.	-	n.a.	n.a.
UK/2013/ewe	94.3	91.4	91.5	68.0	n.a.	-	99.9
UK/2014/lamb	94.4	91.3	91.6	68.0	n.a.	99.7	-
<i>ORF1ab</i>							
BoAstV-BH89/14	-	99.0	98.9	76.6	n.a.	99.0	76.6
BoAstV-CH15	92.4	-	99.2	76.3	n.a.	98.7	98.6
OvAstV-CH16	92.3	98.6	-	76.6	n.a.	98.5	98.5
OvAstV-1	70.0	70.4	70.2	-	n.a.	76.5	76.6
OvAstV-2	n.a.	n.a.	n.a.	n.a.	-	n.a.	n.a.
UK/2013/ewe	94.3	91.8	91.8	70.0	n.a.	-	99.9
UK/2014/lamb	94.3	92.1	92.1	70.0	n.a.	99.9	-
<i>ORF2</i>							
BoAstV-BH89/14	-	98.7	99.3	72.2	26.3	95.0	95.1
BoAstV-CH15	94.9	-	99.0	72.2	26.2	94.6	94.7
OvAstV-CH16	94.7	99.3	-	72.5	26.2	95.0	95.1
OvAstV-1	69.3	69.5	69.4	-	23.8	73.1	73.3
OvAstV-2	41.0	41.5	41.6	40.3	-	25.1	25.3
UK/2013/ewe	87.9	88.3	88.3	69.8	42.0	-	99.9
UK/2014/lamb	87.9	88.3	88.2	69.7	42.0	99.9	-

ORF: open reading frame. White cells: nucleotide similarity. Grey cells: amino acid similarity.

*Ovine astrovirus-CH16 (OvAstV CH16, in bold) shows high similarities with the neurotropic bovine astrovirus strains (BoAstV-BH89/14, BoAstV-CH15) and the neurotropic ovine strains (UK/2013/ewe, UK/2014/lamb), but not with the ovine feces strains (OvAstV-1, OvAstV-2).

Technical Appendix Table 2. Amino acid polymorphisms among neurotropic astrovirus strains in ruminants


<i>nsp1ab</i>	BoAstV-CH15	BoAstV-BH89/14	UK/2013/ewe	UK/2014/lamb	OvAstV-CH16
a.a.					
35	A	S	S	S	A
104	A	V	A	A	A
110	R	K	K	K	R
211	I	F	I	I	I
212	M	V	V	V	V
223	Q	R	Q	Q	Q
253	M	V	M	M	M
258	K	N	N	N	N
264	T	T	A	A	T
301	Y	Y	Y	Y	H
457	S	S	P	P	S
476	H	H	Y	Y	H
497	A	A	V	V	A
581	G	V	V	V	V
584	I	I	T	T	I
631	A	T	T	T	T
652	A	T	T	T	A
689	Y	Y	Y	Y	H
694	E	E	E	E	D
848	A	A	A	A	V
928	N	N	D	D	N
952	D	D	D	D	E
1081	V	V	I	I	V
1191	M	I	I	I	M
1200	N	N	N	N	S
1127	S	S	A	A	S
<i>capsid</i>	BoAstV-CH15	BoAstV-BH89/14	UK/2013/ewe	UK/2014/lamb	OvAstV-CH16
a.a.					
34	N	N	I	I	N
40	A	A	S	S	A
46	T	T	A	A	T
48	P	S	P	P	S
50	I	I	F	F	I
52	A	A	T	T	A
54	F	F	S	S	F
55	V	V	I	I	V
164	V	V	A	A	V
185	I	T	T	T	I
302	S	F	S	S	S
400	T	T	A	A	T
413	C	Y	Y	Y	Y
430	T	T	A	A	T
438	A	A	K	K	A
440	V	V	M	M	V
448	T	T	G	G	T
449	S	S	P	P	S
451	I	T	T	T	T
460	G	G	S	S	G
465	V	V	I	I	V
469	Y	Y	F	F	Y
516	A	A	S	S	A
518	T	T	S	S	T
524	V	V	I	I	V
531	V	V	I	I	V
554	N	N	D	N	N
555	R	R	Q	Q	R
563	E	E	D	D	E
564	T	T	P	P	T
565	S	S	A	A	S
575	M	M	I	I	M
579	Y	Y	I	I	Y
580	G	G	N	N	G
582	T	T	A	A	T
588	I	I	V	V	I
590	V	V	F	F	V
596	I	I	F	F	I
625	K	R	K	K	K

<i>nsp1ab</i>	BoAstV-CH15	BoAstV-BH89/14	UK/2013/ewe	UK/2014/lamb	OvAstV-CH16
635	T	T	S	S	T
664	S	N	N	N	N
732	I	I	I	I	V
754	P	S	S	S	S


nsp1ab: nonstructural precursor protein. *capsid*: capsid precursor protein. a.a.: amino acid position.

Technical Appendix Table 3. Primers used in this study


Primer name	Sequence (5'-3')	Orientation	Application
BoAstV 15 2do	GTC CCT TGA CCA TTG TTG C	Reverse	Reverse transcription
BoAstV 15 3fo	GTC TTG CGC GCT GAG C	Forward	Amplification in screening RT-PCR
BoAstV 15 3do	TGG GTA ATT CTC TAA GCT GTA CTT	Reverse	Reverse transcription and amplification in screening RT-PCR
BoAstV 15 q1do	AAG CTC GAA TTC GCT CCT TCT TAA CCT TAG AGT TAA C	Reverse	Reverse transcription
BoAstV 15 q2do	AGA CTC GAA TTC GGA GGT GTA GGG TAC TGC	Reverse	Reverse transcription
CH15O2c <i>NheI</i> F	ATC GAG CTA GCA AGG GAG GAC CAA AGT TTG AC	Forward	Amplification of ORF2-con fragment
CH15O2c <i>Bam</i> HI R	CGA CTG GAT CCA GCC AGC GAT GCA TAT ACA AG	Reverse	Amplification of ORF2-con fragment
CH15O2v <i>NheI</i> F	ATC GAG CTA GCG ACT TCT CAG CCA GCC CTT G	Forward	Amplification of ORF2-var fragment
CH15O2v <i>Bam</i> HI R	CGA CTG GAT CCT TCT GAG GAC GAC CCA GAC	Reverse	Amplification of ORF2-var fragment



Technical Appendix Figure 1. Sequencing information obtained by next-generation sequencing (NGS) investigation of the affected sheep (ID 41669), related to the genomic structure of bovine astrovirus CH15. nt: nucleotide position. ORF: open reading frame. Black rectangles: untranslated regions. Dark gray rectangle: contiguous sequence (contig) obtained by NGS. Black lines delimited by bars: genome regions used for the expression of recombinant antigens in antibodies production. Black lines with arrow heads: cDNA fragments obtained by reverse transcription and used for Sanger sequencing. Arrow heads: primer binding sites and direction of Sanger sequencing.



Technical Appendix Figure 2. Phylogenetic comparison of the putative capsid protein sequence of representative *Astroviridae* genotype species. The virus found in association with encephalitis in a sheep (OvAstV-CH16) is depicted in bold. Colors indicate viruses that belong to the same astrovirus species. Asterisks signal astrovirus strains that were found in association with encephalitis. HAstV: human astrovirus; PoAstV: porcine astrovirus; MiAstV: mink astrovirus; BoAstV: bovine astrovirus; MAstV: mamastrovirus; AAstV: avastrovirus.



Technical Appendix Figure 3. Correlation of histological lesions with immunohistochemistry (IHC) findings in the sheep positive for ovine astrovirus CH16 (ID 41669). A. Hippocampus, hematoxylin-eosin staining. The animal suffered from severe nonsuppurative encephalitis: marked perivascular cuffing (black arrow) and gliosis (white arrow) are characteristic lesions that can be seen here. B. Hippocampus, IHC with the antibodies against the variable region of the putative capsid protein of bovine astrovirus CH15 (BoAstV-CH15). Positive staining is recognizable by dark red coloring. C. Cerebellum, hematoxylin-eosin staining. Black arrows: perivascular cuffing. D. Cerebellum, IHC with the antibodies against the variable region of the putative capsid protein of BoAstV-CH15. Remarkable positive staining of Purkinje cells.