

# Veal Liver as Food Vehicle for Human *Campylobacter* Infections

Colette Gaulin, Danielle Ramsay, Réjean Dion,  
Marc Simard, Céline Gariépy, Éric Levac,  
Karon Hammond-Collins,  
Maude Michaud-Dumont, Mélanie Gignac,  
Marc Fiset

A matched case–control study in Quebec, Canada, evaluated consumption of veal liver as a risk factor for campylobacteriosis. *Campylobacter* was identified in 28 of 97 veal livers collected concurrently from slaughterhouses and retailers. Veal liver was associated with human *Campylobacter* infection, particularly when consumed undercooked.

Recent investigations conducted in Quebec, Canada, after an increased number of sporadic campylobacteriosis illnesses suggested that consumption of veal liver may be a risk factor for campylobacteriosis. Many of the persons infected reported eating veal liver, and many of those had eaten it pink or undercooked. The association between campylobacteriosis and the consumption of meat products, including chicken liver and offal from different animal species, has been previously described (1–5). We designed an epidemiologic study to examine the relationship between veal liver consumption and campylobacteriosis.

## The Study

We conducted a matched case–control study to examine a potential association between veal liver consumption and campylobacteriosis, using salmonellosis cases as controls. The study began in September 2016 and continued for 9 months. Salmonellosis and campylobacteriosis cases are reportable in Quebec; we selected all subjects from the provincial reportable disease registry. We used a systematic sampling method to select every fifth reported campylobacteriosis case-patient

≥45 years of age. We paired each campylobacteriosis case-patient with 1 salmonellosis case-patient by age group (45–64 and ≥65 y) and sex; both infections were confirmed by fecal culture. We matched case-patients if the salmonellosis sample was collected within a window of 7 days before to 60 days after the campylobacteriosis sample was collected. Inclusion criteria for cases and controls were infection that was sporadic and domestically acquired. Exclusion criteria were co-infection with another pathogen, being part of a recognized outbreak, or contact with a gastroenteritis case-patient ≤10 days before illness.

We administered a structured questionnaire by telephone to collect information on exposures in the 7 days before illness onset. Exposures were consumption of meat and unpasteurized milk products, contact with animals, drinking and recreational water exposures, and occupational exposures. In particular, we investigated consumption of a variety of livers and the degree to which they were cooked. We conducted matched univariate and multivariate analysis to estimate odds ratios (OR) for each exposure.

In addition, we collected samples of veal, chicken, pork, and beef livers from slaughterhouses and retail stores in Quebec between October 2014 and March 2017. We tested each liver specimen for the presence of *Campylobacter*, *Salmonella*, and *Escherichia coli* O157:H7 by using standardized methods (6,7).

We matched a total of 112 campylobacteriosis cases to salmonellosis cases. We found no significant statistical difference in the age or sex distribution of retained cases or controls and the excluded patients. The species of *Campylobacter* were *C. jejuni* (79.5%), *C. jejuni/coli* undifferentiated (3.6%), *C. coli* (0.9%), other (1.8%), and not identified (14.3%). Among campylobacteriosis case-patients, 42 (37.5%) consumed veal liver and 29 (69.0%) ate it undercooked.

Only the consumption of veal liver and having contact with farm animals were statistically significantly associated with campylobacteriosis (Table 1). After applying the Bonferroni correction to adjust for multiple comparisons (0.05 level of significance divided by 45 variables tested yields  $\alpha = 0.001$ ), only veal liver consumption remained as a statistically significant exposure (matched OR 9.50, 95% CI 3.39–26.62;  $p = 0.000001$ ).

Among veal liver consumers, adequate cooking (e.g., well-cooked vs. pink or rare, on the basis of the participant's subjective observation) was protective. Specifically, 13 (30.2%) of 43 case-patients versus 6 (85.7%) of 7 controls

Author affiliations: Ministère de la Santé et des Services Sociaux, Québec City, Québec, Canada (C. Gaulin, M. Fiset); Ministère de l'Agriculture des Pêcheries et de l'Alimentation du Québec, Québec City (D. Ramsay, M. Michaud-Dumont, M. Gignac); Laboratoire de Santé Publique du Québec, Sainte-Anne-de-Bellevue, Québec City (R. Dion); Institut National de Santé Publique du Québec, Québec City (M. Simard); Centre Intégré de Santé et de Services Sociaux de la Montérégie-Centre, Longueuil, Québec (C. Gariépy, É. Levac); Agence de la Santé Publique du Canada, Longueuil (K. Hammond-Collins)

DOI: <https://doi.org/10.3201/eid2406.171900>

**Table 1.** Results of univariate matched analysis for patients with campylobacteriosis or salmonellosis, Quebec, Canada, September 2016–May 2017\*

Exposures	No. pairs					Matched odds ratio†		p value§
	e	f	g	h	Total	Estimate	95% CI‡	
<b>Liver</b>								
Chicken liver	0	4	1	104	109	4.00	0.45–35.79	0.2
Pork liver	0	2	5	102	109	0.40	0.08–2.06	0.3
Beef liver	0	3	3	105	111	1.00	0.20–4.96	1.0
Veal liver	4	38	4	62	108	9.50	3.39–26.62	0.000001
Lamb liver	0	3	0	109	112	Undefined	Undefined	1.0
<b>Poultry</b>								
Breaded chicken	10	15	18	64	107	0.83	0.42–1.65	0.6
Ground chicken	0	3	10	95	108	0.30	0.08–1.09	0.07
Whole chicken	44	35	22	8	109	1.59	0.93–2.71	0.09
<b>Pork</b>								
Ham	27	26	26	29	108	1.00	0.58–1.72	1.0
Bacon	8	23	27	50	108	0.85	0.49–1.49	0.6
Ground pork	4	12	14	76	106	0.86	0.40–1.85	0.7
<b>Beef</b>								
Ground beef	47	25	24	9	105	1.04	0.60–1.82	0.9
Roast beef	3	22	15	66	106	1.47	0.76–2.83	0.3
Beef steak	20	21	25	39	105	0.84	0.47–1.50	0.6
<b>Veal</b>								
Ground veal	0	8	13	86	107	0.62	0.26–1.49	0.3
Veal escalope	0	8	2	98	108	4.00	0.85–18.84	0.08
<b>Unpasteurized milk products</b>								
Raw milk	0	2	1	107	110	2.00	0.18–22.06	0.6
Raw-milk cheese	0	10	4	93	107	2.50	0.78–7.97	0.1
<b>Water exposures</b>								
Drinking water from source other than aqueduct	3	12	12	81	108	1.00	0.45–2.23	1.0
<b>Animal exposures</b>								
Dog or cat	24	23	22	38	107	1.05	0.58–1.88	0.9
Farm animal	0	10	1	100	111	10.00	1.28–78.12	0.03
Work in contact with animals	0	8	2	102	112	4.00	0.80–38.67	0.07

\*Results by matched analysis: e, campylobacteriosis case exposed, salmonellosis case exposed; f, campylobacteriosis case exposed, salmonellosis case not exposed; g, campylobacteriosis case not exposed, salmonellosis case exposed; h, campylobacteriosis case not exposed, salmonellosis case not exposed.

†By McNemar method.

‡Lower and upper limits determined by McNemar or exact method.

§By Wald or Fisher exact test, bilateral.

ate their veal liver well-cooked (unmatched OR 0.07, 95% CI 0.002–0.72;  $p = 0.02$ ). Multivariate analysis using logistic regression confirmed that a statistically significant association between the consumption of veal liver and campylobacteriosis remained when all other exposures were included as covariates. Although we conducted this study among persons  $\geq 45$  years of age, it is reasonable to assume that eating veal liver, especially undercooked, would also carry risk for younger persons.

We sampled 339 veal, pork, chicken, and beef livers collected from 138 retailers and 16 slaughterhouses. When we evaluated all livers collected at these locations, we detected *Campylobacter* in 28.0% of veal livers, 22.2% of pork livers, 36.8% of chicken livers, and 19.1% of beef livers (Table 2). We detected *Salmonella* more frequently in chicken livers (22.1%) and pork livers (19.1%) than in veal livers (3.1%); we did not detect *Salmonella* in beef livers. We rarely identified *E. coli* O157:H7 in livers of any kind. The proportion of contaminated livers differed between animal species and also with respect to location of sampling. A higher proportion of veal livers (35.7%) collected from

retailers were contaminated by *Campylobacter*, compared with veal livers collected from slaughterhouses (16.2%). We observed the reverse for chicken and pork livers. The reason for these variations is unclear at this time, but this finding may be an artifact resulting from the relatively small number of samples taken at each location.

Cattle are a well-known reservoir for a variety of *Campylobacter* species, such as *C. jejuni*, *C. coli*, and *C. fetus* (8,9). *Campylobacter* species have been isolated from beef intestinal contents and also from beef bile, bile ducts, gallbladder, and liver (10–14). The gallbladder and bile contain substances that have a chemoattractant effect on *C. jejuni*, which explains the presence of *Campylobacter* within the biliary tract (10,15). Liver contamination varies between animal species (10–14). Chicken liver, for example, can be contaminated by *Campylobacter* and *Salmonella* and has been the source of several outbreaks (3,4,11,13). Because few case-patients consumed livers from other animal species during our study, we were not able to identify any substantial risks associated with those exposures.

**Table 2.** Animal livers collected from retailers and slaughterhouses and percentage positive for *Campylobacter*, *Salmonella*, and *Escherichia coli* O157:H7, Quebec, Canada, October 2014–March 2017

Source and pathogen	Veal livers		Pork livers		Chicken livers		Beef livers		Total
	No.	% Positive	No.	% Positive	No.	% Positive	No.	% Positive	
Retailer	59		27		19		41		146
<i>Campylobacter</i>		35.7		16.7		10.5		28.2	
<i>Salmonella</i>		5.1		44.4		32.0		0	
<i>E. coli</i> O157:H7		0		0		0		0	
Slaughterhouse	38		41		58		56		193
<i>Campylobacter</i>		16.2		25.6		45.6		12.7	
<i>Salmonella</i>		0		2.4		19.0		0	
<i>E. coli</i> O157:H7		0		2.4		0		0	
Total	97		68		77		97		339
<i>Campylobacter</i>		28.0		22.2		36.8		19.1	
<i>Salmonella</i>		3.1		19.1		22.1		0	
<i>E. coli</i> O157:H7		0		1.5		0		0	

Because livers may be collected from several animals and stored together, they may be contaminated during the evisceration process or by cross-contamination (11). Both the external and internal tissue of a liver may be contaminated with *Campylobacter*, and inadequate cooking may not fully inactivate *Campylobacter* and *Salmonella* (10,11), which is a cause for concern because  $\approx 70\%$  of the patients with campylobacteriosis who consumed veal liver in our study reported eating it undercooked. We did not examine possible cross-contamination of foods and surfaces and the host-related factors that may increase the risk for enteric diseases.

## Conclusions

Our study identified a strong and statistically significant association between the consumption of veal liver and sporadic, domestically acquired campylobacteriosis among persons  $\geq 45$  years of age in Quebec. We found that adequate cooking of veal liver mitigates the risk of infection. We detected *Campylobacter* in almost one third of veal livers we sampled from slaughterhouses and retail stores, which supports our finding that veal liver consumption is a risk factor for campylobacteriosis. In light of these results, we recommend the dissemination of safe food handling practices for veal liver and other offal for retailers, food establishments, slaughterhouses, and the general public.

## Acknowledgments

We thank the regional public health authorities for their participation in the study, as well as the food inspectors and laboratory staff who supported the collection and analysis of liver samples.

## About the Author

Dr. Gaulin is a physician epidemiologist who works in public health at the Protection Branch of the Ministry of Health in Quebec, Canada. She works in infectious diseases on enteric and nonenteric disease surveillance and also coordinates provincial outbreak investigations.

## References


- Little CL, Gormley FJ, Rawal N, Richardson JF. A recipe for disaster: outbreaks of campylobacteriosis associated with poultry liver pâté in England and Wales. *Epidemiol Infect.* 2010;138:1691–4. <http://dx.doi.org/10.1017/S0950268810001974>
- Oshin F. *Campylobacter* outbreak in North Ayrshire. Health Protection Scotland Weekly Report. 2005;39:232.
- Forbes KJ, Gormley FJ, Dallas JF, Labovitiadi O, MacRae M, Owen RJ, et al. *Campylobacter* immunity and coinfection following a large outbreak in a farming community. *J Clin Microbiol.* 2009;47:111–6. <http://dx.doi.org/10.1128/JCM.01731-08>
- O'Leary MC, Harding O, Fisher L, Cowden J. A continuous common-source outbreak of campylobacteriosis associated with changes to the preparation of chicken liver pâté. *Epidemiol Infect.* 2009;137:383–8. <http://dx.doi.org/10.1017/S0950268808001003>
- Stafford RJ, Schluter P, Kirk M, Wilson A, Unicomb L, Ashbolt R, et al.; OzFoodNet Working Group. A multi-centre prospective case-control study of *Campylobacter* infection in persons aged 5 years and older in Australia. *Epidemiol Infect.* 2007;135:978–88. <http://dx.doi.org/10.1017/S0950268806007576>
- Ministère de l'Agriculture, des Pêcheries et de l'Alimentation du Québec (MAPAQ). Laboratoire d'expertises et d'analyses alimentaires (LEAA). Isolation of *Campylobacter* spp. in food [in French]. LEAA-M-MIC-061. 2017 Apr 10.
- Ministère de l'Agriculture, des Pêcheries et de l'Alimentation du Québec (MAPAQ). Laboratoire d'expertises et d'analyses alimentaires (LEAA). Preparation of food specimens for microbiological analysis [in French]. LEAA-REF-MIC-550. 2015 Nov 26.
- Busato A, Hofer D, Lentze T, Gaillard C, Burnens A. Prevalence and infection risks of zoonotic enteropathogenic bacteria in Swiss cow-calf farms. *Vet Microbiol.* 1999;69:251–63. [http://dx.doi.org/10.1016/S0378-1135\(99\)00119-4](http://dx.doi.org/10.1016/S0378-1135(99)00119-4)
- Garcia MM, Lior H, Stewart RB, Ruckerbauer GM, Trudel JR, Skljarevski A. Isolation, characterization, and serotyping of *Campylobacter jejuni* and *Campylobacter coli* from slaughter cattle. *Appl Environ Microbiol.* 1985;49:667–72.
- Strachan NJ, MacRae M, Thomson A, Rotariu O, Ogden ID, Forbes KJ. Source attribution, prevalence, and enumeration of *Campylobacter* spp. from retail liver. *Int J Food Microbiol.* 2012;153:234–6. <http://dx.doi.org/10.1016/j.ijfoodmicro.2011.10.033>
- Noormohamed A, Fakhr MK. A higher prevalence rate of *Campylobacter* in retail beef livers compared to other beef and pork meat cuts. *Int J Environ Res Public Health.* 2013;10:2058–68. <http://dx.doi.org/10.3390/ijerph10052058>
- Enokimoto M, Kubo M, Bozono Y, Mieno Y, Misawa N. Enumeration and identification of *Campylobacter* species in the

- liver and bile of slaughtered cattle. *Int J Food Microbiol.* 2007;118:259–63. <http://dx.doi.org/10.1016/j.ijfoodmicro.2007.07.057>
13. Saito S, Yatsuyanagi J, Harata S, Ito Y, Shinagawa K, Suzuki N, et al. *Campylobacter jejuni* isolated from retail poultry meat, bovine feces and bile, and human diarrheal samples in Japan: comparison of serotypes and genotypes. *FEMS Immunol Med Microbiol.* 2005;45:311–9. <http://dx.doi.org/10.1016/j.femsim.2005.05.006>
  14. Kramer JM, Frost JA, Bolton FJ, Wareing DR. *Campylobacter* contamination of raw meat and poultry at retail sale: identification of multiple types and comparison with isolates from human infection. *J Food Prot.* 2000;63:1654–9. <http://dx.doi.org/10.4315/0362-028X-63.12.1654>
  15. Hugdahl MB, Beery JT, Doyle MP. Chemotactic behavior of *Campylobacter jejuni*. *Infect Immun.* 1988;56:1560–6.


Address for correspondence: Colette Gaulin, Ministère de la Santé et des Services Sociaux du Québec, 1075 chemin Ste-Foy, Québec City, QC G1S 2M1, Canada; email: [colette.gaulin@msss.gouv.qc.ca](mailto:colette.gaulin@msss.gouv.qc.ca)

## April 2016: Food Safety

- Determinants and Drivers of Infectious Disease Threat Events in Europe
- Shiga Toxin–Producing *Escherichia coli* O157, England and Wales, 1983–2012
- Nosocomial Co-Transmission of Avian Influenza A(H7N9) and A(H1N1)pdm09 Viruses between 2 Patients with Hematologic Disorders
- Quantifying Transmission of *Clostridium difficile* within and outside Healthcare Settings
- Microevolution of Monophasic *Salmonella Typhimurium* during Epidemic, United Kingdom, 2005–2010



- Molecular Typing and Epidemiology of Human Listeriosis Cases, Denmark, 2002–2012
- Limited Dissemination of Extended-Spectrum  $\beta$ -Lactamase– and Plasmid-Encoded AmpC–Producing *Escherichia coli* from Food and Farm Animals, Sweden
- Transmission of Middle East Respiratory Syndrome Coronavirus Infections in Healthcare Settings, Abu Dhabi
- Lassa Virus Seroprevalence in Sibirilia Commune, Bougouni District, Southern Mali
- *Neisseria meningitidis* Serogroup X in Sub-Saharan Africa Post-Ebola Syndrome, Sierra Leone
- Evaluation of Viremia Frequencies of a Novel Human Pegivirus by Using Bioinformatic Screening and PCR
- Shiga Toxin 1–Producing *Shigella sonnei* Infections, California, United States, 2014–2015
- Adenovirus Type 7 Pneumonia in Children Who Died from Measles-Associated Pneumonia, Hanoi, Vietnam, 2014



- Elevated *Toxoplasma gondii* Infection Rates for Retinas from Eye Banks, Southern Brazil
- Arenavirus Diversity and Phylogeography of *Mastomys natalensis* Rodents, Nigeria
- Cross-Neutralization between Human and African Bat Mumps Viruses
- Definitive Hosts of *Versteria* Species (Cestoda: Taeniidae) Causing Fatal Infection in North America
- Effectiveness of a Mobile Short-Message-Service–Based Disease Outbreak Alert System in Kenya
- Deletion Variants of Middle East Respiratory Syndrome Coronavirus from Humans, Jordan, 2015
- Nontyphoidal *Salmonella* Infection, Guangdong Province, China, 2012
- Nipah Virus Transmission from Bats to Humans Associated with Drinking Traditional Liquor Made from Date Palm Sap, Bangladesh, 2011–2014
- Low-Cost National Media-Based Surveillance System for Public Health Events, Bangladesh
- Exportations of Symptomatic Cases of MERS-CoV Infection to Countries Outside the Middle East
- Severe Infections with Human Adenovirus 7d in 2 Adults in Family, Illinois, USA, 2014



**EMERGING  
INFECTIOUS DISEASES**

<https://wwwnc.cdc.gov/eid/articles/issue/22/4/table-of-contents>