

clinical and microbiological cure of *M. conceptionense* pneumonitis by using azithromycin and doxycycline in a patient with HIV/AIDS in the United States.

About the Author

Dr. Michienzi is a clinical assistant professor and pharmacist at the University of Illinois at Chicago College of Pharmacy, Chicago, IL. Her research interests are HIV–hepatitis C virus co-infection, HIV in incarcerated and underserved populations, and pharmacist roles in care.

References

- Adékambi T, Stein A, Carvajal J, Raoult D, Drancourt M. Description of *Mycobacterium conceptionense* sp. nov., a *Mycobacterium fortuitum* group organism isolated from a posttraumatic osteitis inflammation. *J Clin Microbiol*. 2006; 44:1268–73. <https://doi.org/10.1128/JCM.44.4.1268-1273.2006>
- Oda G, Winters M, Pacheco SM, Sikka M, Bleasdale S, Dunn B, et al. Identical strain of *Mycobacterium conceptionense* isolated from patients at 2 veterans affairs medical centers within the same metropolitan area over a 4-year period. Abstract no. 648. In: Abstracts of ID Week 2017, San Diego, October 4–8, 2017. Arlington (VA): Infectious Diseases Society of America; 2017.
- Kim SY, Kim MS, Chang HE, Yim JJ, Lee JH, Song SH, et al. Pulmonary infection caused by *Mycobacterium conceptionense*. *Emerg Infect Dis*. 2012;18:174–6. <https://doi.org/10.3201/eid1801.110251>
- Shojaei H, Hashemi A, Heidarieh P, Ataei B, Naser AD. Pulmonary and extrapulmonary infection caused by *Mycobacterium conceptionense*: the first report from Iran. *JRSM Short Rep*. 2011;2:31. <https://doi.org/10.1258/shorts.2010.010103>
- Liao CH, Lai CC, Huang YT, Chou CH, Hsu HL, Hsueh PR. Subcutaneous abscess caused by *Mycobacterium conceptionense* in an immunocompetent patient. *J Infect*. 2009;58:308–9. <https://doi.org/10.1016/j.jinf.2009.02.012>
- Lee KH, Heo ST, Choi SW, Park DH, Kim YR, Yoo SJ. Three cases of postoperative septic arthritis caused by *Mycobacterium conceptionense* in the shoulder joints of immunocompetent patients. *J Clin Microbiol*. 2014;52:1013–5. <https://doi.org/10.1128/JCM.02652-13>
- Yang HJ, Yim HW, Lee MY, Ko KS, Yoon HJ. *Mycobacterium conceptionense* infection complicating face rejuvenation with fat grafting. *J Med Microbiol*. 2011;60:371–4. <https://doi.org/10.1099/jmm.0.024554-0>
- Zhang X, Liu W, Liu W, Jiang H, Zong W, Zhang G, et al. Cutaneous infections caused by rapidly growing mycobacteria: case reports and review of clinical and laboratory aspects. *Acta Derm Venereol*. 2015;95:985–9. <https://doi.org/10.2340/00015555-2105>
- Yaita K, Matsunaga M, Tashiro N, Sakai Y, Masunaga K, Miyoshi H, et al. *Mycobacterium conceptionense* bloodstream infection in a patient with advanced gastric carcinoma. *Jpn J Infect Dis*. 2017;70:92–5. <https://doi.org/10.7883/yoken.JJID.2015.626>
- Thibeaut S, Levy PY, Pelletier ML, Drancourt M. *Mycobacterium conceptionense* infection after breast implant surgery, France. *Emerg Infect Dis*. 2010;16:1180–1. <https://doi.org/10.3201/eid1607.090771>

Address for correspondence: Sarah M. Michienzi, University of Illinois at Chicago, College of Pharmacy, Pharmacy Practice (MC 886), 833 S Wood St, Ste 164, Chicago, IL 60612, USA; email: msarah@uic.edu

Emergence of Influenza A(H7N4) Virus, Cambodia

Dhanasekaran Vijaykrishna, Yi-Mo Deng, Miguel L. Grau, Matthew Kay, Annika Suttie, Paul F. Horwood, Wantanee Kalpravidh, Filip Claes, Kristina Osbjør, Phillipe Dussart, Ian G. Barr, Erik A. Karlsson

Author affiliations: Peter Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia (D. Vijaykrishna, Y.-M. Deng, M. Kay, I.G. Barr); Monash University, Melbourne (D. Vijaykrishna, M.L. Grau); Institut Pasteur du Cambodge, Phnom Penh, Cambodia (A. Suttie, P.F. Horwood, P. Dussart, E.A. Karlsson); James Cook University, Townsville, Queensland, Australia (P.F. Horwood); Food and Agriculture Organization of the United Nations, Bangkok, Thailand (W. Kalpravidh, F. Claes); Food and Agriculture Organization of the United Nations, Phnom Penh, Cambodia (K. Osbjør)

DOI: <https://doi.org/10.3201/eid2510.190506>

Active surveillance in high-risk sites in Cambodia has identified multiple low-pathogenicity influenza A(H7) viruses, mainly in ducks. None fall within the A/Anhui/1/2013(H7N9) lineage; however, some A(H7) viruses from 2018 show temporal and phylogenetic similarity to the H7N4 virus that caused a nonfatal infection in Jiangsu Province, China, in December 2017.

Avian influenza virus (AIV) subtype A(H7) is of concern because it has been a leading cause of zoonotic infections over the past 2 decades (1). The A/Anhui/1/2013-lineage A(H7N9) viruses, a leading cause of zoonotic infections in Asia since 2013, have not been detected in the Greater Mekong Subregion, but independent H7 lineages, including H7N3, H7N7, and H7Nx, have been detected occasionally in Cambodia since 2009 (2–4). H7N3 virus was detected from a duck mortality event in Kampong Thom during January 2017 (2), and H7N7 virus was detected in a live-bird market (LBM) in Takeo in September 2017 (4). Furthermore, highly pathogenic avian influenza (HPAI) A(H5N1) and low-pathogenicity avian influenza (LPAI) A(H9N2) are endemic in Cambodia (5); 59 poultry outbreaks of AIV and 56 human HPAI A(H5N1) cases have occurred since 2006. Although the exact ecologic links are unknown, serologic studies suggest that AIVs of multiple subtypes are frequently introduced into poultry in Cambodia, possibly through cross-border trade or through wild birds (2,6,7).

In December 2017, a 68-year-old woman in Jiangsu, China, who had underlying medical conditions was infected by an LPAI influenza A(H7N4) virus, which led to severe pneumonia and intensive care unit admission, but

she recovered and left the hospital after 21 days (8,9). Genetically similar H7N4 viruses were subsequently detected in contact chickens (9,10) and aquatic poultry in Jiangsu (GISAID, <https://www.gisaid.org>), substantiating that the infection was zoonotic and raising concerns of endemicity of H7N4 in the region. Because of the antigenic differences between the A/Jiangsu/1/2018-like A(H7N4) virus and other H7 lineages (10), including A/Anhui/1/2013(H7N9) lineage, this newly detected H7N4 virus has been proposed as a vaccine candidate for pandemic preparedness (10).

Beginning in February 2018, 2 months after the H7N4 case in China, this virus was detected in ducks in Cambodia; the frequency of detection increased in March and April (4). Therefore, because of the novelty of the strain and the association with human infection, we sought to understand the genomic diversity of H7 viruses in Cambodia.

We characterized the whole genomes (for sequencing methods, see Appendix, <http://wwwnc.cdc.gov/EID/article/25/10/19-0506-App1.pdf>) of 16 viruses collected during

2015–2018 subtyped by reverse transcription PCR (RT-PCR) as having an H7 hemagglutinin (HA) gene or an N4 neuraminidase (NA) gene; we also included viruses for which the HA or NA could not be typed but that were epidemiologically associated with A(H7) viruses (Appendix Table). We obtained samples from poultry swabs collected across multiple LBMs, slaughterhouses, and poultry collection centers in Cambodia; most H7 viruses originated from domestic ducks (4).

All AIV samples collected during February–April 2018 in Cambodia ($n = 9$) (Appendix Table 1, Figure 1) contained ≥ 1 segment with high similarity and common evolutionary origins to the Jiangsu H7N4 samples, whereas AIV collected before this period formed other independent lineages derived from wild birds. Seven H7-HA from viruses collected in 2018 in Cambodia (4 H7N4, 1 H7N5, 1 H7Nx, and 1 H7 with mixed N4 and N7 segments) were most closely related to the HA and NA genes of Jiangsu H7N4 isolates; all 6 N4 NA were most closely related to the NA genes of Jiangsu H7N4 isolates (Figure). We also




Figure. Maximum-likelihood phylogeny of the evolutionary origins of influenza A(H7N4) virus in Cambodia and comparison with reference isolates. H7 hemagglutinin (A) and N4 neuraminidase (B) genes were inferred using a general time-reversible nucleotide substitution model with a gamma distribution of among-site rate variation in RAXML version 8 (<https://cme.h-its.org/exelixis/web/software/raxml>) and visualized using Figtree version 1.4 (<http://tree.bio.ed.ac.uk/software/figtree/>). Branch support values were generated using 1,000 bootstrap replicates. Scale bars represent nucleotide substitutions per site. A color version of this figure is available online (<https://wwwnc.cdc.gov/EID/article/25/10/19-0506-F1.htm>).

observed close relationships between the Jiangsu and Cambodia isolates in the internal segments polymerase basic protein 2 (PB2), polymerase acidic protein (PA), and nucleoprotein (NP); most viruses carried a common PA gene (Appendix Figure 1). However, none of the H7N4 viruses from Cambodia shared all segments with Jiangsu isolates, indicating continual reassortment with AIV co-circulating in the region.

Phylogenetic analysis showed that the Cambodia–Jiangsu H7-HA genes emerged during late 2017 (mean time to most recent common ancestor November 2017; 95% CI August 2016–July 2017) and were derived from H7N7 and H7N2 viruses previously detected in aquatic birds in east Asia (Appendix Figure 2). In contrast, the N4-NA exhibited a greater diversity in Cambodia (mean time to most recent common ancestor January 2016; 95% CI January 2015–November 2016) and were derived from H10N4 and H8N4 viruses previously detected in Georgia, Russia, and Mongolia.

Our results show that H7N4 is a newly developing virus lineage that originated from divergent avian lineages within the Eurasian AIV gene pool. The dispersed genetic origins from locations in Europe and central Asia and the similarity of the Cambodia and Jiangsu H7N4 samples indicates that the H7N4 virus was generated in aquatic birds, likely just before their first detection. Detection of H7N4 in LBMs in Cambodia in such a short span of time at such a large spatial distance highlights the risk and potential for rapid spread of AIV lineages throughout the region. The ability to infect a human subject, the continual reassortment and antigenic evolution of this lineage, and the endemicity of numerous LPAI and HPAI viruses may further increase the risk for zoonotic infections and warrants vigilant, active surveillance in wild birds and poultry in the region.

Acknowledgments

The investigators thank everyone on the Influenza Team in the Virology Unit at Institut Pasteur du Cambodge (IPC) who contributed to this study, including Viseth Srey Horm, Songha Tok, Phalla Y, Ponnarath Keo, Sereyath Sun, Sonita Kol, Chiva Sum, Sarath Sin, Kim Lay Chea, and Sokhoun Yann. In addition, we thank Veasna Duong. We also thank all of the support teams at IPC, including the drivers and facilities personnel who make these studies possible, as well as all the local teams, epidemiologists, veterinary officers, and other staff from the National Health and Production Research Institute. We also thank everyone from the regional and country Food and Agriculture Organization of the United Nations (FAO) offices, especially Makara Hak and Aurelie Brioudes.

Work at Institut Pasteur in Cambodia was supported by the Office of the Assistant Secretary for Preparedness and Response,

US Department of Health and Human Services (grant no. IDSEP 140020-01-00; <http://www.asideproject.org>), and by the FAO, with funding from the United States Agency for International Development (USAID) under the Emerging Pandemics Threats 2 (EPT-2) project. The Melbourne WHO Collaborating Centre for Reference and Research on Influenza is supported by the Australian Government Department of Health. D.V. and M.L.G. are supported by contract no. HHSN27220140006C from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, US Department of Health and Human Services.

About the Author

Dr. Vijaykrishna is an evolutionary biologist based at the Biomedicine Discovery Institute in Melbourne, working closely with staff and students at IPC and World Health Organization Collaborating Centre, Melbourne. His primary research interests are in using disease surveillance and comparative genomics to track and solve problems in clinical and veterinary virology.

References

1. Abdelwhab EM, Veits J, Mettenleiter TC. Prevalence and control of H7 avian influenza viruses in birds and humans. *Epidemiol Infect.* 2014;142:896–920. <https://doi.org/10.1017/S0950268813003324>
2. Suttie A, Yann S, Y P, Tum S, Deng YM, Hul V, et al. Detection of low pathogenicity influenza A(H7N3) virus during duck mortality event, Cambodia, 2017. *Emerg Infect Dis.* 2018;24:1103–7. <https://doi.org/10.3201/eid2406.172099>
3. World Health Organization. Cumulative number of confirmed human cases for avian influenza A(H5N1) reported to WHO, 2003–2018. 2018 [cited 2018 Dec 13]. https://www.who.int/influenza/human_animal_interface/H5N1_cumulative_table_archives
4. Karlsson EA, Horm SV, Tok S, Tum S, Kalpravidh W, Claes F, et al. Avian influenza virus detection, temporality and co-infection in poultry in Cambodian border provinces, 2017–2018. *Emerg Microbes Infect.* 2019;8:637–9. <https://doi.org/10.1080/22221751.2019.1604085>
5. Suttie A, Karlsson EA, Deng YM, Horm SV, Yann S, Tok S, et al. Influenza A(H5N1) viruses with A(H9N2) single gene (matrix or PB1) reassortment isolated from Cambodian live bird markets. *Virology.* 2018;523:22–6. <https://doi.org/10.1016/j.virol.2018.07.028>
6. Horm SV, Tarantola A, Rith S, Ly S, Gambaretti J, Duong V, et al. Intense circulation of A/H5N1 and other avian influenza viruses in Cambodian live-bird markets with serological evidence of sub-clinical human infections. *Emerg Microbes Infect.* 2016;5:e70. <https://doi.org/10.1038/emi.2016.69>
7. Van Kerkhove MD, Vong S, Guitian J, Holl D, Mangtani P, San S, et al. Poultry movement networks in Cambodia: implications for surveillance and control of highly pathogenic avian influenza (HPAI/H5N1). *Vaccine.* 2009;27:6345–52. <https://doi.org/10.1016/j.vaccine.2009.05.004>
8. Gao P, Du H, Fan L, Chen L, Liao M, Xu C, et al. Human infection with an avian-origin influenza A (H7N4) virus in Jiangsu: a potential threat to China. *J Infect.* 2018;77:249–57. <https://doi.org/10.1016/j.jinf.2018.07.005>

9. Tong XC, Weng SS, Xue F, Wu X, Xu TM, Zhang WH. First human infection by a novel avian influenza A(H7N4) virus. *J Infect*. 2018;77:249–57. <https://doi.org/10.1016/j.jinf.2018.06.002>
10. World Health Organization. Human infection with avian influenza A(H7N4) virus—China. *Disease Outbreak News (DON)* 2018 [cited 2018 Dec 17]. <https://www.who.int/csr/don/22-february-2018-ah7n4-china>

Address for Correspondence: Erik A. Karlsson, Institut Pasteur du Cambodge Virology Unit, 5 Monivong Blvd, PO Box 983, Phnom Penh, Cambodia; email: ekarlsson@pasteur-kh.org

***Mycobacterium marseillense* Infection in Human Skin, China, 2018**

Bibo Xie,¹ Yanqing Chen,¹ Jian Wang, Wei Gao, Haiqing Jiang, Jiya Sun, Xindong Jin, Xudong Sang, Xiaobing Yu, Hongsheng Wang

Author affiliations: Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union Medical College, Nanjing, China (B. Xie, Y. Chen, W. Gao, H. Jiang, H. Wang); Zhejiang Institute of Dermatology, Deqing, China (B. Xie, J. Wang, X. Jin, X. Sang, X. Yu); Jiangsu Key Laboratory of Molecular Biology for Skin Diseases and STIs, Nanjing (Y. Chen, W. Gao, H. Jiang, H. Wang); Suzhou Institute of Systems Medicine, Chinese Academy of Medical Sciences, Suzhou, China (J. Sun); Centre for Global Health, Nanjing Medical University, Nanjing (H. Wang)

DOI: <https://doi.org/10.3201/eid2510.190695>

We describe a case of facial skin infection and sinusitis caused by *Mycobacterium marseillense* in an immunocompetent woman in China in 2018. The infection was cleared with clarithromycin, moxifloxacin, and amikacin. Antimicrobial drug treatments could not be predicted by genetic analyses; further genetic characterization would be required to do so.

Mycobacterium marseillense is a member of the *M. avium* complex (1) that has caused infections with lymphatic or pulmonary involvement sporadically in humans (2–4). We report *M. marseillense* infection involving facial skin in an immunocompetent woman in eastern China.

¹These authors contributed equally to this article.

In April 2018, a 59-year-old woman was referred to our institute (Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union Medical College, Nanjing, China) for a 4-year history of an erythematous plaque with ulceration located on the right cheek. The primary lesion was a small erythematous patch that gradually developed into an asymptomatic ulcerative plaque (i.e., the plaque had no heat, swelling, pain, or pruritus). She also reported occasional bloody, purulent nasal discharge over the course of 2 years. Two years before visiting our hospital, cutaneous tuberculosis was suspected, so she received treatment for tuberculosis (rifampin, isoniazid, ethambutol, pyrazinamide) for 10 months. No obvious improvement was observed with this treatment. Her medical history was otherwise unremarkable.

On physical examination, an infiltrated erythematous plaque with yellow scales and crusts on the right cheek was visible (Figure, panel A). Routine laboratory tests showed no remarkable findings. The results of autoantibody and HIV tests were negative, and immune subset cell counts were unremarkable. Histologic examination showed infiltration of a large number of lymphocytes, plasma cells, and neutrophils and some tissue cells in the dermis (Appendix Figure 1, <https://wwwnc.cdc.gov/EID/article/25/10/19-0695-App1.pdf>). Computed tomography scan of the paranasal sinuses showed bilateral maxillary, right ethmoid, and frontal sinusitis (Figure, panel C). Culture and PCR for mycobacteria in nasal discharge yielded negative findings.

After 3 weeks of skin tissue culture at 32°C in Löwenstein–Jensen medium, we observed smooth, yolk-yellow bacterial colonies (Appendix Figure 2). Ziehl–Neelsen staining confirmed the cultured organism was acid-fast bacilli. Sequence analysis indicated that the complete genetic sequence of 16S rRNA was 99.0%, *hsp65* 100%, and *rpoB* 99.8% homologous with *M. marseillense* strain FLAC0026. Phylogenetic analysis of the 16S rRNA sequence showed the isolate clustered with *M. chimaera* and *M. intracellulare* (Figure, panel D). Although the 16S rRNA gene sequence of the isolate was 100% similar to *M. intracellulare* subsp. *yongonense* 05-1390, the sequence similarities to *hsp65* and *rpoB* were relatively low. Sequence analyses suggested *M. marseillense* infection.

Referring to the guidelines for pulmonary *M. avium* complex disease, we treated the patient with the antimicrobial drugs clarithromycin, rifampin, and ethambutol (5). Afterward, in vitro drug susceptibility testing showed the isolate was sensitive to clarithromycin, azithromycin, and amikacin; moderately sensitive to moxifloxacin; and resistant to ethambutol and rifampin. Therefore, 3 months after initiating treatment, we changed the regimen to clarithromycin, moxifloxacin, and amikacin, which she received for 2 months. The patient’s skin lesions healed gradually, and nasal symptoms disappeared, but a scar and erythema

Emergence of Influenza A(H7N4) Virus, Cambodia

Appendix

Whole Genome Sequencing

Whole genome sequences were generated by next generation sequencing using Ion Torrent PGM with universal primers, and by Sanger sequencing using gene-specific primers, as described previously (1), followed by a customized de novo assembly pipeline of corrected reads using SPAdes v3 (2). Consensus sequences were then generated by BLAST against a local database built from all influenza records available in the National Center for Biotechnology Information database (<https://www.ncbi.nlm.nih.gov>) as of November 2018, followed by maximum likelihood phylogenetic analysis (3) using sequences from the GISAID repository (<https://www.gisaid.org/>; downloaded in January 2019).

Molecular Characterization of Cambodian Influenza A(H7N4) Viruses

None of the Cambodian and Jiangsu A(H7N4) viruses contained known amino acid mutations that confer adaptation of AIV to humans (e.g., PB2 627/701 or HA 186/226/228; H3-HA numbering) (4). However, despite showing independent origins, most of the Cambodian viruses contained amino acid mutations in the matrix (M) gene, namely N30D and T215A, shown to increase pathogenicity of A(H5N1) virus in rodents (5). All samples contained markers that indicated that these viruses would be sensitive to known antiviral drugs, including adamantanes, oseltamivir/zanamivir and baloxivir-marboxil (6).

References

1. Suttie A, Yann S, Y P, Tum S, Deng YM, Hul V, et al. Detection of low pathogenicity influenza A(H7N3) virus during duck mortality event, Cambodia, 2017. *Emerg Infect Dis.* 2018;24:1103–7. <https://doi.org/10.3201/eid2406.172099>

2. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 2012;19:455–77. <https://doi.org/10.1089/cmb.2012.0021>
3. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics.* 2014;30:1312–3. <https://doi.org/10.1093/bioinformatics/btu033>
4. Yamayoshi S, Fukuyama S, Yamada S, Zhao D, Murakami S, Uraki R, et al. Amino acids substitutions in the PB2 protein of H7N9 influenza A viruses are important for virulence in mammalian hosts. *Sci Rep.* 2015;5:8039. <https://doi.org/10.1038/srep08039>
5. Fan S, Deng G, Song J, Tian G, Suo Y, Jiang Y, et al. Two amino acid residues in the matrix protein M1 contribute to the virulence difference of H5N1 avian influenza viruses in mice. *Virology.* 2009;384:28–32. <https://doi.org/10.1016/j.virol.2008.11.044>
6. Shin WJ, Seong BL. Novel antiviral drug discovery strategies to tackle drug-resistant mutants of influenza virus strains. *Expert Opin Drug Discov.* 2019;14:153–68. <https://doi.org/10.1080/17460441.2019.1560261>
7. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evol.* 2018;4:vey016. <https://doi.org/10.1093/ve/vey016>

Appendix Table. Low pathogenic avian influenza viruses detected in poultry in Cambodia and their genomic similarity to A/Jiangsu-China/1/2018


Virus name	Subtype	Date	Location	HA†	NA	PB2	PB1	PA	NP	MP	NS
A/duck/Cambodia/12T-24-1-D3/2018	H7N4‡	2018 Mar 22	Takeo	99.4	99.4	98.1	96.1	99.4	96.7	98.0	97.2
A/duck/Cambodia/12T-24-1-D17/2018	H7N7/4	2018 Mar 22	Takeo	96.7	99.4	95.0	95.7	99.5	96.5	N/A	97.2
A/duck/Cambodia/C50W8M1/2018	H7N4	2018 Feb 24	Phnom Penh	99.4	99.5	93.7	96.3	99.4	96.5	98.3	97.2
A/duck/Cambodia/14T-24-1-D11/2018	HxN4	2018 Apr 10	Takeo	N/A	99.0	93.7	95.3	97.4	97.0	97.5	97.8
A/duck/Cambodia/12T-24-1-D1/2018	H7N4	2018 Mar 22	Takeo	99.0	99.3	93.7	95.7	95.9§	96.7	97.4	N/A
A/duck/Cambodia/12T-24-1-D7/2018	H7N4	2018 Mar 22	Takeo	99.3	99.4	N/A	N/A	N/A	N/A	N/A	N/A
A/host/Cambodia/C76W14M1/2018	H7Nx	2018 Apr 05	Phnom Penh	99.2	N/A	N/A	N/A	N/A	N/A	N/A	N/A
A/duck/Cambodia/C70W14M1/2018	H7N5	2018 Apr 05	Phnom Penh	99.4	N5	91.9	95.3	95.0	98.7	98.2	97.3
A/duck/Cambodia/C8W6M1/2018	H10N7	2018 Feb 06	Phnom Penh	H10	N7	93.5	95.6	94.5	96.5	97.1	97.0
A/duck/Cambodia/3T-24-1-D14/2017	H7N7	2017 Sep 17	Takeo	95.1	N7	91.0	94.4	91.0	97.0	97.4	97.3
A/duck/Cambodia/3T-24-1-D18/2017	H7N7	2017 Sep 17	Takeo	95.2	N7	92.8	94.6	91.7	97.0	97.4	88.8
A/duck/Cambodia/b0120501/2017	H7N3	2017 Jan 12	Outbreak	95.5	N3	86.7	94.9	92.4	88.1	94.7	71.5
A/duck/Cambodia/b0116502/2017	H7N3	2017 Jan 12	Outbreak	95.5	N3	86.7	95.0	92.5	88.2	94.6	71.8
A/chicken/Cambodia/Z437W28M3/2015	H7N7	2015 Jun 09	Takeo	96.7	N7	93.5	96.5	94.9	97.3	98.3	97.9
A/duck/Cambodia/Z535W33M3/2015	H7N3	2015 Aug 13	Takeo	95.9	N3	N/A	N/A	N/A	N/A	N/A	N/A
A/duck/Cambodia/Z683W41M1/2015	H7Nx	2015 Oct 10	Phnom Penh	96.0	N/A	N/A	N/A	N/A	N/A	N/A	N/A

*HA, hemagglutinin; MP, N/A, not available; NA, neuraminidase; NP, nucleoprotein; NS, nonstructural protein; PA, polymerase acidic protein; PB, polymerase basic protein.



†Percentage similarity of strains from Cambodia to corresponding segment of A/Jiangsu/1/2018, with bold values showing similar evolutionary origins. The evolutionary relationships of these segments are shown in the Figure in the main text and Appendix Figures 1 and 2.

‡Bold type represents similarity of surface genes of Cambodian viruses to A/Jiangsu/1/2018.

§Incomplete gene sequence.



Appendix Figure 1. Maximum likelihood trees of the PB2, PA, and NP gene segments showing the Jiangsu H7N4 and their most closely related strains. Phylogenetic trees were inferred using a general time reversible nucleotide substitution model with a gamma distribution of among-site rate variation in RAxML v8, and visualized using Figtree v1.4 (<http://tree.bio.ed.ac.uk/software/figtree>). Branch support values were generated using 1,000 bootstrap replicates. Scale bar represents nucleotide substitutions per site.

A**B**

Appendix Figure 2. Time scale of evolution of H7N4 virus. Dated phylogeny of the (A) H7-HA and (B) N4-NA genes inferred using an uncorrelated log-normal relaxed clock model with constant population demographic prior in BEAST 1.10 (7), and visualized using Figtree v1.4 (<http://tree.bio.ed.ac.uk/software/figtree>). Node bars represent 95% confidence intervals.