

Bombali Virus in *Mops condylurus* Bats, Guinea

Lyudmila S. Karan, Marat T. Makenov, Mikhail G. Korneev, Noumany Sacko, Sanaba Boumbaly, Sergey A. Yakovlev, Kerfalla Kourouma, Roman B. Bayandin, Anastasiya V. Gladysheva, Andrey V. Shipovalov, Irina A. Yurganova, Yana E. Grigorieva, Marina V. Fedorova, Svetlana A. Scherbakova, Vladimir V. Kutryev, Alexander P. Agafonov, Renat A. Maksyutov, German A. Shipulin, Viktor V. Maleev, Mamadou Boiro, Vasiliy G. Akimkin, Anna Y. Popova

Author affiliations: Central Research Institute of Epidemiology, Moscow, Russia (L.S. Karan, M.T. Makenov, Y.A. Grigorieva, M.V. Fedorova, V.V. Maleev, V.G. Akimkin); Russian Research Anti-Plague Institute, Saratov, Russia (M.G. Korneev, S.A. Yakovlev, S.A. Scherbakova, V.V. Kutryev); International Center for Research of Tropical Infections in Guinea, N'Zerekore, Guinea (N. Sacko, S. Boumbaly); Research Institute of Applied Biology of Guinea, Kindia, Guinea (N. Sacko, S. Boumbaly, K. Kourouma, M. Boiro); State Research Center of Virology and Biotechnology VECTOR, Kol'tsovo, Russia (R.B. Bayandin, A.V. Gladysheva, A.V. Shipovalov, I.A. Yurganova, A.P. Agafonov, R.A. Maksyutov); Center of Strategical Planning and Biomedical Health Risks Management, Moscow (G.A. Shipulin); Federal Service for Surveillance on Consumer Rights Protection and Human Wellbeing, Moscow (A.Y. Popova)

DOI: <https://doi.org/10.3201/eid2509.190581>

In 2018, a previously unknown Ebola virus, Bombali virus, was discovered in Sierra Leone. We describe detection of Bombali virus in Guinea. We found viral RNA in internal organs of 3 Angolan free-tailed bats (*Mops condylurus*) trapped in the city of N'Zerekore and in a near-by village.

In 2018, a new species of the genus *Ebolavirus* (family *Filoviridae*), Bombali virus (BOMV), was discovered in Sierra Leone (1). The virus was detected in oral and rectal swab specimens from 2 free-tailed bat species, *Chaerephon pumilus* (little free-tailed bat) and *Mops condylurus* (Angolan free-tailed bat). Both bat species are widespread in Africa, and their ranges include countries where human Ebola virus disease (EVD) outbreaks have occurred. Forbes et al. (2) detected BOMV RNA in mouth swabs and internal parenchymal organs, except kidneys, of *M. condylurus* bats in Kenya in May 2018.

Most known outbreaks of EVD among humans were Zaire Ebola virus, including the large epidemic in West Africa during 2013–2016 (3). The reservoir hosts of Ebola virus (EBOV) remain unclear, but bats commonly are suspected. Viral RNA and EBOV antibodies have been detected in a few species of fruit bats (4,5). The discovery of BOMV supports the hypothesis regarding the role of bats as hosts of EBOVs, but further study is required to determine the bat species involved in viral transmission, prevalence of the virus in bat populations, and geographic distribution of the virus.

We detected BOMV RNA in free-tailed bats in N'Zerekore Prefecture, Guinea. We trapped bats in Guinea and Liberia during 2018–2019 (Table; Appendix, <https://wwwnc.cdc.gov/EID/article/25/9/19-0581-App1.pdf>) and detected BOMV RNA by reverse transcription PCR in 2 pools of kidney and lung samples from 2 *M. condylurus* bats captured in Yalenzou village in May 2018 (cycle threshold [C_t] 17.4 and 19.6) and in a pool of liver and spleen tissues (C_t 28.2) of an *M. condylurus* bat from a school in the city of N'Zerekore in March 2019 (Table). Blood, intestine, and brain samples were negative for viral RNA. Sequencing of the 483-bp fragment of the large gene (GenBank accession no. MK543447) demonstrated 99.3% identity with BOMV RNA from Sierra Leone (accession no. NC039345) and 98.3% identity with BOMV RNA from Kenya (accession no. MK340750).

Marí Saéz et al. (5) suggested that the Angolan free-tailed bat was the most plausible zoonotic source of the EVD

Table. Locations where free-tailed bats were trapped and tested for Bombali virus, Guinea and Liberia*

Location	Date	Total trapped	Species, no. tested (no. positive)		
			<i>Mops condylurus</i>	<i>Chaerephon pumilus</i>	<i>Chaerephon cf. major</i>
Yalenzou	2018 May 4	26	26 (2)	0	0
Gbao	2018 May 2	1	0	1	0
Yalenzou	2019 Mar 2	30	30	0	0
Bololowee†	2019 Mar 3	11	11	0	0
N'Zerekore, school	2019 Mar 5	47	27 (1)	0	20
N'Zerekore, house	2019 Mar 6	23	1	0	22
N'Zerekore, gazebo	2019 Mar 7	5	0	0	5
Dar Salam‡	2019 Mar 17	22	14	8	0
Total		165	109 (3)	9	47

*All bats were collected in N'Zerekore Prefecture, Guinea, except as indicated, and were tested by reverse transcription PCR for Bombali virus.

†Liberia.

‡Madina Oula Prefecture, Guinea.

epidemic in West Africa. In addition, EBOV nucleotide sequences previously have been found in *Hypsignathus monstrosus*, *Epomops franqueti*, and *Myonycteris torquata* bats in Gabon (6). He et al. (7) detected filovirus RNA in brown fruit bats (*Rousettus leschenaultii*) in China, and another study showed that 3 distinct groups of unclassified filoviruses are circulating in *Eonycteris spelaea* and *Rousettus* spp. fruit bats in China (8). These studies demonstrate that bats are promising targets for identifying emerging filoviruses, and additional Chiroptera species, both insectivorous and fruit bats, should be examined for EBOVs.

EBOV IgG was detected in the human population of Sierra Leone in 2006, 8 years before the EVD outbreak began in that country (9). Seroprevalence to EBOVs was also found in the medical staff of hospitals that were not involved in treating EVD-positive patients and in community contacts that worked with villages where EVD was not detected (10). The highest seroprevalence to EBOVs was found in the inhabitants of villages with the lowest number of documented EVD cases during the 2013–2016 outbreak in Sierra Leone (10). Cross-reactivity or nonspecific binding could be responsible for artifacts of immunoassay. However, other plausible explanations for the presence of antibodies against EBOV among persons with no symptoms of EVD exist, including subclinical EBOV infection in humans and antibody reactions to previously undiscovered, nonpathogenic filoviruses. The newly discovered BOMV could be a causative agent of these types of asymptomatic infections that produce antibodies with cross-reactivity to other EBOVs. Other undiscovered filoviruses also could be circulating in the region. Further surveillance with family-level primers is needed for insectivorous bats, as well as fruit bats and patients with acute infections.

Although BOMV had been detected in the northern part of Sierra Leone (1) and in the Taita Hills area of Kenya (2), we isolated it from bats in Guinea, far from these sites. Our finding provides additional evidence that BOMV is more widely distributed than previously suspected. Consequently, we advise screening of free-tailed bats for BOMV across their range. The high concentration of BOMV RNA we found in the internal organs of *M. condylurus* bats provides additional confirmation that BOMV could amplify in these bats and that this species is a reservoir host of this virus.

About the Author

Mrs. Karan is the head of the research group of Vector-borne and Zoonotic Diseases at Central Research Institute of Epidemiology, Moscow, Russia. Her research interests include tickborne and mosquito-borne diseases and related molecular diagnostics.

References

1. Goldstein T, Anthony SJ, Gbakima A, Bird BH, Bangura J, Tremeau-Bravard A, et al. The discovery of Bombali virus adds further support for bats as hosts of ebolaviruses. *Nat Microbiol*. 2018;3:1084–9. <https://doi.org/10.1038/s41564-018-0227-2>
2. Forbes KM, Webala PW, Jääskeläinen AJ, Abdurahman S, Ogola J, Masika MM, et al. Bombali virus in *Mops condylurus* bat, Kenya. *Emerg Infect Dis*. 2019;25:955–7. <https://doi.org/10.3201/eid2505.181666>
3. Baize S, Pannetier D, Oestereich L, Rieger T, Koivogui L, Magassouba N, et al. Emergence of Zaire Ebola virus disease in Guinea. *N Engl J Med*. 2014;371:1418–25. <https://doi.org/10.1056/NEJMoa1404505>
4. Pourrut X, Délicat A, Rollin PE, Ksiazek TG, Gonzalez JP, Leroy EM. Spatial and temporal patterns of Zaire ebolavirus antibody prevalence in the possible reservoir bat species. *J Infect Dis*. 2007;196(Suppl 2):S176–83. <https://doi.org/10.1086/520541>
5. Marí Saéz A, Weiss S, Nowak K, Lapeyre V, Zimmermann F, Dux A, et al. Investigating the zoonotic origin of the West African Ebola epidemic. *EMBO Mol Med*. 2015;7:17–23. <https://doi.org/10.15252/emmm.201404792>
6. Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, et al. Fruit bats as reservoirs of Ebola virus. *Nature*. 2005;438:575–6. PubMed <https://doi.org/10.1038/438575a>
7. He B, Feng Y, Zhang H, Xu L, Yang W, Zhang Y, et al. Filovirus RNA in Fruit Bats, China. *Emerg Infect Dis*. 2015;21:1675–7. <https://doi.org/10.3201/eid2109.150260>
8. Yang XL, Zhang YZ, Jiang RD, Guo H, Zhang W, Li B, et al. Genetically diverse filoviruses in *Rousettus* and *Eonycteris* spp. bats, China, 2009 and 2015. *Emerg Infect Dis*. 2017;23:482–6. <https://doi.org/10.3201/eid2303.161119>
9. Richardson ET, Kelly JD, Barrie MB, Mesman AW, Karku S, Quiwa K, et al. Minimally symptomatic infection in an Ebola ‘hotspot’: A cross-sectional serosurvey. *PLoS Negl Trop Dis*. 2016;10:e0005087. <https://doi.org/10.1371/journal.pntd.0005087>
10. Mafopa NG, Russo G, Wadoum REG, Iwerima E, Batwala V, Giovanetti M, et al. Seroprevalence of Ebola virus infection in Bombali District, Sierra Leone. *J Public Health Africa*. 2017;8:732. <https://doi.org/10.4081/jphia.2017.732>

Address for correspondence: Marat T. Makenov, Research Group of Vector-Borne and Zoonotic Diseases, 3-A Novogireevskaya St. 415, Moscow 111123, Russia; email: mmakenov@gmail.com

Bombali Virus in *Mops condylurus* Bats, Guinea

Appendix

Methods

Animal Sampling

We trapped bats in the villages of Yalenzou (7°42'29.99" N, 8°41'20.60" W) and Gbao (7°32'19.42" N, 8°46'18.17" W) in N'Zerekore Prefecture, Guinea in May 2018. In both villages, bats had roosted under the corrugated tin roofs of houses and granaries. We captured 26 free-tailed bats in Yalenzou and 1 bat in Gbao.

In March 2019, we conducted a second trapping session in N'Zerekore Prefecture. We captured 30 free-tailed bats in Yalenzou and 76 free-tailed bats in 3 different sites of N'Zerekore city (a school, a house, and a gazebo in the yard of another house). In addition, we trapped 11 Angolan free-tailed bats in Bololowee village in Liberia (7°41'19.21" N, 8°40'32.47" W), and 22 bats in Dar Salam village (9°58'36.88" N, 12°20'21.08" W) in Madina Oula Prefecture, Guinea.

We weighed; morphologically described; and measured the length of the head and body, forearm, tail, tibia, hind foot, and ear of trapped bats. Then we genotyped them by sequencing the cytochrome c oxidase subunit I (COI) gene using the following primers: ST-COI-F2 5'-CTCYACYAAWCAAYAAAGACATTGGAAC-3' (1) and jgHCO2198 5'-TAIACYTCIGGRTGICCRARAAYCA-3' (2).

We used cardiac puncture to collect blood into sterile tubes with 0.5 M EDTA. We then euthanized the animals and obtained sections of the brain, liver, spleen, kidney, lung, and intestines through sterile necropsy. In 2018, we collected a blood sample, a pool of lung and kidney tissues, and a pool of liver and spleen tissues from each animal. In 2019, we expanded the sample panel and separately collected the following into sterile tubes: blood; an oral swab; a

rectal swab; and lung, kidney, and intestine samples. Because previous reports did not find viral RNA in the liver or spleen (3,4), we homogenized these tissues in a single pool for each animal.

We stored blood fractions from N’Zerekore Prefecture at 10°C for 5–7 days, and at –28°C for the next 10 days before studying. We collected tissue samples in RNAlater (ThermoFisher Scientific, <https://www.thermofisher.com>) and stored them at 6–10°C for 12–15 days before studying. We stored samples from Madina Oula Prefecture at –70°C. We followed standard methods for the safe handling and sampling of small mammals potentially infected with infectious pathogens (5).

PCR Array and Sequencing Analyses

We extracted total RNA from blood, brain, and internal organs (lung, liver, spleen, kidney, and intestines) by using RIBO-prep kit (Central Research Institute of Epidemiology, Moscow, Russia, <http://www.crie.ru>). Then we prepared cDNA by using REVERTA-L (Central Research Institute of Epidemiology, Moscow, Russia). We screened samples for Bombali virus RNA by quantitative reverse transcription PCR by using Filo_UCD_qFor and Filo_UCD_qRev primers, and the Filo_UCD_probe probe sequence (3). We amplified cDNA with the Filo-MOD-FWD and FiloL.conR primers, and sequenced it with Filo-MOD-RVS probe (3).

References

1. Makenov MT, Karan LS, Shashina NI, Akhmetshina MB, Zhurenkova OB, Kholodilov IS, et al. First detection of tick-borne encephalitis virus in Ixodes ricinus ticks and their rodent hosts in Moscow, Russia. *bioRxiv*. 2018;480475. <http://biorxiv.org/content/early/2018/12/20/480475>
2. Geller J, Meyer C, Parker M, Hawk H. Redesign of PCR primers for mitochondrial cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic surveys. *Mol Ecol Resour*. 2013;13:851–61.
3. Goldstein T, Anthony SJ, Gbakima A, Bird BH, Bangura J, Tremeau-Bravard A, et al. The discovery of Bombali virus adds further support for bats as hosts of ebolaviruses. *Nat Microbiol*. 2018;3:1084–9.
4. Forbes KM, Webala PW, Jääskeläinen AJ, Abdurahman S, Ogola J, Masika MM, et al. Bombali Ebola virus in *Mops condylurus* bat, Kenya. *Emerg Infect Dis*. 2019;25.

5. Mills JN, Childs JE, Ksiazek TG, Peters CJ, Velleca WM. Methods for trapping and sampling small mammals for virologic testing. Atlanta: U.S. Centers for Disease Control and Prevention. 1995.