

References

1. Telenti A, Marchesi F, Balz M, Bally F, Böttger EC, Bodmer T. Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *J Clin Microbiol*. 1993;31:175–8. <https://doi.org/10.1128/JCM.31.2.175-178.1993>
2. Klotz D, Barth SA, Baumgärtner W, Hewicker-Trautwein M. *Mycobacterium avium* subsp. *hominissuis* infection in a domestic rabbit, Germany. *Emerg Infect Dis*. 2018;24:596–8. <https://doi.org/10.3201/eid2403.171692>
3. Agdestein A, Olsen I, Jørgensen A, Dønne B, Johansen TB. Novel insights into transmission routes of *Mycobacterium avium* in pigs and possible implications for human health. *Vet Res (Faisalabad)*. 2014;45:46. <https://doi.org/10.1186/1297-9716-45-46>
4. Ignatov D, Kondratieva E, Azhikina T, Apt A. *Mycobacterium avium*-triggered diseases: pathogenomics. *Cell Microbiol*. 2012;14:808–18. <https://doi.org/10.1111/j.1462-5822.2012.01776.x>
5. Nishiuchi Y, Iwamoto T, Maruyama F. Infection sources of a common non-tuberculous mycobacterial pathogen, *Mycobacterium avium* complex. *Front Med (Lausanne)*. 2017;4:27. <https://doi.org/10.3389/fmed.2017.00027>
6. Komatsu T, Inaba N, Kondo K, Nagata R, Kawaji S, Shibahara T. Systemic mycobacteriosis caused by '*Mycobacterium avium* subspecies *hominissuis*' in a 14-month-old Japanese black beef steer. *J Vet Med Sci*. 2017;79:1384–8. <https://doi.org/10.1292/jvms.17-0204>
7. Yoshida S, Araki T, Asai T, Tsuyuguchi K, Arikawa K, Iwamoto T, et al. Phylogenetic uniqueness of *Mycobacterium avium* subspecies *hominissuis* isolated from an abnormal pulmonary bovine case. *Infect Genet Evol*. 2018;62:122–9. <https://doi.org/10.1016/j.meegid.2018.04.013>
8. Fitzgerald SD, Kaneene JB. Wildlife reservoirs of bovine tuberculosis worldwide: hosts, pathology, surveillance, and control. *Vet Pathol*. 2013;50:488–99. <https://doi.org/10.1177/0300985812467472>
9. Canada Food Inspection Agency, Government of Canada. Bovine tuberculosis. 2019 [cited 2020 Jan 13]. <https://inspection.gc.ca/animal-health/terrestrial-animals/diseases/reportable/bovine-tuberculosis/eng/1330205978967/1330206128556>
10. Wobeser G. Bovine tuberculosis in Canadian wildlife: an updated history. *Can Vet J*. 2009;50:1169–76.

Address for correspondence: Jamie L. Rothenburger, Faculty of Veterinary Medicine, University of Calgary, 3280 Hospital Dr NW, Calgary, Alberta T2N 4Z6, Canada; email: jamie.rothenburger@ucalgary.ca.

Public Mental Health Crisis during COVID-19 Pandemic, China

Lu Dong, Jennifer Bouey

Author affiliations: RAND Corporation, Santa Monica, California, USA (L. Dong); RAND Corporation, Arlington, Virginia, USA (J. Bouey); Georgetown University, Washington, DC, USA (J. Bouey)

DOI: <https://doi.org/10.3201/eid2607.200407>

The 2019 novel coronavirus disease emerged in China in late 2019–early 2020 and spread rapidly. China has been implementing emergency psychological crisis interventions to reduce the negative psychosocial impact on public mental health, but challenges exist. Public mental health interventions should be formally integrated into public health preparedness and emergency response plans.

China was the first country affected by the pandemic of 2019 novel coronavirus disease (COVID-19), caused by severe acute respiratory syndrome coronavirus 2. Several unique characteristics of China's COVID-19 epidemic patterns and its management policy prompted a heightened public mental health crisis. First, many Chinese residents still remember the 2003 outbreak of severe acute respiratory syndrome (SARS) and its effect on China's social life and economy (1). COVID-19 is more transmissible than SARS, and the case-fatality rate (2.3%) is substantially higher than that for seasonal influenza (2). The uncertain incubation period of the virus and its possible asymptomatic transmission cause additional fear and anxiety. Second, the government's initial downplaying of the epidemic's severity eroded public trust in the government's decision-making transparency and competency. Third, unprecedented large-scale quarantine measures in all major cities, which essentially confine residents to their homes, are likely to have a negative psychosocial effect on residents (3). Fourth, reports of shortages of medical protective supplies, medical staff, and hospital beds in Wuhan and the surrounding areas soon followed the citywide quarantine and caused enormous concern throughout the nation. Last, a unique "infodemic" – an overabundance of (mis)information on social media (4) and elsewhere – poses a major risk to public mental health during this health crisis.

As during the 2003 SARS and 2014 Ebola virus disease outbreaks, generalized fear and fear-induced overreactive behavior were common among the public; both can impede infection control (5,6). In addition, psychiatric disorders, such as depression,

anxiety, and posttraumatic stress disorder, developed in high-risk persons, especially survivors and front-line healthcare workers (7).

On the basis of these recent experiences, the National Health Commission of China released a notification on January 26, 2020, providing guiding principles of the emergency psychological crisis interventions to reduce the psychosocial effects of the COVID-19 outbreak (8). This notification specified that psychological crisis intervention should be part of the public health response to the COVID-19 outbreak, organized by the joint prevention and control mechanism at the city, municipal, and provincial levels, and that the interventions should be differentiated by group. The intervention workforce comprises psychological outreach teams led by psychiatrists and mental health professionals and psychological support hotline teams. An attachment to this notification further outlined the key intervention targets for 6 groups: confirmed patients, persons under investigation for COVID-19, healthcare workers, persons in immediate contact with patients, ill persons who refuse to seek care, and susceptible persons/the general public (Appendix, <https://wwwnc.cdc.gov/EID/article/26/7/20-0407-App1.pdf>).

The release of such policy guidance acknowledges China's recognition of public mental health needs during the outbreak. However, the notification does not specify how different resources should be mobilized and coordinated or, more important, who should deliver which type of interventions, for which group in need, and by which delivery mode(s). The policy guidance also does not indicate operationalization of how various groups should be screened or assessed to determine the type and level of interventions to provide to each. This level of detail is needed because China lacks a well-established mental healthcare system and has no existing national-level emergency response system and designated workforce to provide the psychological crisis interventions during a national emergency or disaster (X. Chen, X. Fu, unpub. data, <https://doi.org/10.16418/j.issn.1000-3045.20200213001>) (9). Other major challenges to successfully implementing the emergency psychological crisis interventions include China's severe shortage of mental healthcare providers (1.49 psychiatrists/100,000 population, and only half of these psychiatrists have attained a bachelor's degree in medicine), unevenly distributed healthcare resources, and the limitations posed by the mass quarantine (9). For example, hospitals, universities, and a variety of organizations have set up numerous hotlines staffed by volunteers with varying degrees of qualification and experience (8).

These well-meaning efforts can be uncoordinated and inadequately supervised and thus are likely to cause confusion to service consumers and inefficient use of resources.

The challenges reported in China indicate that, for many developing countries, telemedicine should be considered, given the widespread adoption of smartphones, to help remove barriers to accessing quality care for mental health. Task-shifting or -sharing (i.e., shifting service delivery of specific tasks from professionals to persons with fewer qualifications or creating a new cadre of providers with specific training) might help, especially in low-resource areas (10). Countries should also consider requesting support and guidance from global mental healthcare authorities and research communities through international collaborations.

Given lessons learned from past outbreaks in China and other parts of the world, public mental health interventions should be formally integrated into public health preparedness and emergency response plans to effectively curb all outbreaks. The World Health Organization's strategic preparedness and response plan for COVID-19, however, has not yet specified any strategies to address mental health needs of any kind (4). As the virus spreads globally, governments must address public mental health needs by developing and implementing well-coordinated strategic plans to meet these needs during the COVID-19 pandemic.

About the Authors

Dr. Dong is an associate behavioral scientist and a licensed clinical psychologist at RAND Corporation. Her primary research interests are development and improvement of evidence-based psychosocial interventions for youth and adults.

Dr. Bouey is a senior policy researcher and the Tang Chair in China Policy Studies at RAND Corporation and an associate professor of Global Health at Georgetown University. Her primary research interests include the social determinants of health among underserved populations.

References

1. Bouey J. From SARS to 2019-coronavirus (nCoV): U.S.-China collaborations on pandemic response: addendum. Santa Monica (CA): RAND Corporation; 2020 [cited 2020 Mar 23]. <https://www.rand.org/pubs/testimonies/CT523z2.html>
2. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19)—China, 2020. *China CDC Weekly*. 2020;2:113–22.
3. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the

- evidence. *Lancet*. 2020;395:912–20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
4. World Health Organization. 2019 Novel coronavirus (2019-nCoV): strategic preparedness and response plan Feb 3, 2020 [cited 2020 Feb 7]. <https://www.who.int/docs/default-source/coronaviruse/srp-04022020.pdf>
 5. Shultz JM, Cooper JL, Baingana F, Oquendo MA, Espinel Z, Althouse BM, et al. The role of fear-related behaviors in the 2013–2016 West Africa Ebola virus disease outbreak. *Curr Psychiatry Rep*. 2016;18:104. <https://doi.org/10.1007/s11920-016-0741-y>
 6. Person B, Sy F, Holton K, Govert B, Liang A, Garza B, et al; National Center for Infectious Diseases/SARS Community Outreach Team. Fear and stigma: the epidemic within the SARS outbreak. *Emerg Infect Dis*. 2004;10:358–63. <https://doi.org/10.3201/eid1002.030750>
 7. Mak IW, Chu CM, Pan PC, Yiu MG, Chan VL. Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry*. 2009;31:318–26. <https://doi.org/10.1016/j.genhosppsych.2009.03.001>
 8. National Health Commission of China. Principles of the emergency psychological crisis interventions for the new coronavirus pneumonia [in Chinese] [cited 2020 Feb 7]. <http://www.nhc.gov.cn/jkj/s3577/202001/6adc08b966594253b2b791be5c3b9467>
 9. Liang D, Mays VM, Hwang WC. Integrated mental health services in China: challenges and planning for the future. *Health Policy Plan*. 2018;33:107–22. <https://doi.org/10.1093/heapol/czx137>
 10. World Health Organization. Joint WHO/OGAC technical consultation on task shifting: key elements of a regulatory framework in support of in-country implementation of task shifting. Geneva: The Organization; 2007.

Address for correspondence: Lu Dong, RAND Corporation, 1776 Main St, Santa Monica, CA 90401, USA; email: ldong@rand.org

Rhabdomyolysis as Potential Late Complication Associated with COVID-19

Min Jin, Qiaoxia Tong

Author affiliations: Cancer Center, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China (M. Jin); Department of Infectious Diseases, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan (Q. Tong)

DOI: <https://doi.org/10.3201/eid2607.200445>

We describe a patient in Wuhan, China, with severe acute respiratory syndrome coronavirus 2 infection who had progressive pulmonary lesions and rhabdomyolysis with manifestations of lower limb pain and fatigue. Rapid clinical recognition of rhabdomyolysis symptoms in patients with severe acute respiratory syndrome coronavirus 2 infection can be lifesaving.

Recently, the outbreak of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in Wuhan, China, has attracted great attention worldwide (1). SARS-CoV-2, the cause of 2019 novel coronavirus disease (COVID-19), belongs to the β -coronavirus family, which also includes 2 other highly pathogenic human coronaviruses (2): severe acute respiratory syndrome coronavirus and Middle East respiratory syndrome coronavirus. Fever, cough, myalgia, and fatigue are the common symptoms of COVID-19, whereas expectoration, headache, hemoptysis, and diarrhea are relatively rare (3).

Rhabdomyolysis is a life-threatening disorder that manifests with myalgia, fatigue, and pigmenturia; it can also manifest as acute renal failure (4). The inducing factors of rhabdomyolysis include autoimmune myopathies, septicemia, electrolyte abnormalities, substance abuse, alcohol use, or infection (5). Viral infection, especially influenza virus infection, can lead to rhabdomyolysis (6). We report rhabdomyolysis related to COVID-19 in Wuhan, China.

A 60-year-old man in Wuhan sought care in February 2020 for a 6-day history of fever up to 38.3°C and cough. Chest computed tomography performed 3 days before in another hospital showed that the texture of both lungs was thickened and scattered with ground glass shadows (Appendix Figure, <https://wwwnc.cdc.gov/EID/article/26/7/20-0445-App1.pdf>). When the patient arrived, he was alert; heart rate was 89 bpm, blood pressure was 135/91 mm Hg, respiratory rate was 18 breaths/min, temperature was 38.5°C, and saturation of peripheral oxygen was 93%. Physical examination revealed a rough breath sound in the lungs. Laboratory findings included mild leukopenia (3.31×10^9 neutrophils/L [reference $3.5\text{--}9.5 \times 10^9$ neutrophils/L]), increased lactate dehydrogenase (280 U/L [reference 109–245 U/L]), and increased C-reactive protein (111 mg/L [reference 0–8 mg/L]) (Table). Results were in the normal range for creatine kinase (CK) and indicators of hepatic and kidney function. Screenings for common infectious diseases were negative. Real-time reverse-transcription PCR analysis of the patient's throat swab specimen indicated SARS-CoV-2 infection.

Public Mental Health Crisis during COVID-19 Pandemic, China

Appendix

The authors note that these documents below are not official translations provided by the Chinese government. The authors translated these documents for readers who may not read Chinese to understand the content of the notification referenced in the main text.

This technical appendix contains two parts:

1. Guiding Principles for Emergency Psychological Crisis Intervention for Outbreak of the Novel Coronavirus Outbreak
2. Attachment to the “Guiding Principles”: Key Points of Psychological Crisis Intervention for Different Groups

1. Guiding Principles for Emergency Psychological Crisis Intervention for Outbreak of Novel Coronavirus Outbreak

National Health Commission of the People's Republic of China

These guidelines should be implemented under the guidance of trained mental health professionals.

1. Organization and leadership:

The psychological crisis intervention work is led by the joint prevention and containment mechanism (leadership group, headquarters) in response to the COVID-19 epidemic at the provincial, regional, and municipal levels. This mechanism also provides the necessary coordination and financial support. The mental health professional associations and other related associations and institutions should assemble experts with experience in post-disaster psychological crisis intervention, to provide technical guidance and emergency

psychological crisis intervention under the coordination of the government's health authority.

2. Guiding principles

- 1) Integrate psychological crisis intervention into the overall deployment of epidemic prevention and control, with the premise of reducing potential psychological damage caused by the epidemic and promoting social stability and of adjusting the focus of psychological crisis intervention in a timely manner according to the progress of epidemic prevention and control.
- 2) Implement targeted interventions for different groups. Strictly protect the personal privacy of patients. Prevent secondary trauma for both providers and patients.

3. Formulate the intervention plan

1) Purpose

- a. Provide mental health services for the affected population
- b. Provide psychological crisis intervention for the population in need
- c. Actively prevent, mitigate and contain the psychosocial impact of the epidemic
- d. The management and treatment of severe mental disorders should not be interrupted

2) Tasks

- a. Understand the mental health status of various groups affected by the epidemic, and timely identify high-risk groups based on the information at hand to prevent extreme events such as suicide and other impulsive behaviors. Surveillance of early signs of group psychological crisis, timely reporting to the joint mechanism (lead group, command) of epidemic containment, and provide solutions.

- b. Comprehensively apply various psychological crisis intervention techniques and combine them with publicity and education to provide mental health services.
- c. Provide supports including training to social organizations to supply mental health services.
- d. Continue the management, treatment and community care of at-home patients with severe mental disorders.

3) Determine the target population and number.

Population affected by the COVID-19 outbreak is divided into four groups.

The focus of intervention should start with the first-tier population and gradually expand. General communication and education should reach all four tiers of the population.

1. First-tier population: patients confirmed with COVID-19 (hospitalized patients with severe illness and worse), front-line medical staff, front-line disease control staff, and front-line management staff
 2. Second-tier population: mild patients (close contacts, suspected patients) isolated at home, patients with fever who came to the hospital for treatment.
 3. Third-tier population: people related to the first-tier and second-tier population, such as family members, colleagues, friends, epidemic responders, including field commanders, organization managers, and volunteers.
 4. Forth-tier population: residents of geographic areas affected by the epidemic, susceptible population, and general public who are affected by the epidemic prevention and control measures.
- 4) Evaluation of target population and formulation of differentiated intervention plan. Assess the mental health of the target population, identify and distinguish between the high-risk population and the general population in a

timely manner; conduct psychological crisis interventions for the high-risk population and provide mental health education for the general population.

- 5) Develop work schedules. Based on the scope and size of target groups and the available number of psychological crisis intervention personnel, arrange work and develop work schedules.

4. Form a team

1. Psychological intervention medical team. Can be formed as standalone team or be part of the general medical team. The staff is mainly psychiatrists, with clinical psychologists and psychiatric nurses participating. Those with experience in psychological crisis intervention are preferred. When teaming alone, one team leader is assigned, and one liaison is assigned to take charge of the team's logistical support and contact with all parties.
2. Psychological assistance hotline team. Mainly mental health workers who have received psychological hotline training and volunteers who have experience in psychological crisis intervention in public emergencies. Before taking up the job, they should receive psychological assistance training on COVID-19 outbreak and organize experts to provide supervision to the hotline staff.

5. Working methods

1. Psychiatric and mental health experts conduct timely research and judgment based on the epidemic development and the psychological state of the population, and provide decision-making recommendations and consultation for the joint mechanism (lead group, headquarters), and provide training and supervision for personnel participating in the psychological crisis intervention, and provide mental health education for the public.
2. Give full play to the role of "Healthy China", "12320", provincial health platform, existing psychological crisis intervention hotline and multiple online communication methods, coordinate the rotation of psychological workers, and provide 24/7 online services. Provide real-time psychological support to the third- and fourth-tier populations in a timely manner, and provide

supplementary psychological assistance services to the first- and second-tier groups.

3. Extensive mobilization of the society to provide social support based on the needs and difficulties of various groups affected by the epidemic.

2. Attachment to the “Guiding Principles”: Key Points of Psychological Crisis Intervention for Different Groups

1. Confirmed patients

a. Early stage of isolation.

- i. Typical presentation: numbness, denial, anger, fear, anxiety, depression, disappointment, complaining, insomnia, aggression, etc.

- ii. Interventions:

1. Understand that the patients’ emotional responses are normal responses under stress. Be prepared in advance and not be irritated by patients’ aggressive or depressive behaviors and lose the stance of professional healthcare provider, such as arguing with the patient or being overly involved.
2. Under the premise of understanding the patient, psychological crisis intervention should be given in addition to medical treatments, such as timely assessment of suicide risk, self-injury risk, risk of aggressive behavior, positive psychological support, and avoid direct conflict with the patient. Seek a psychiatric consultation if necessary. Explain the importance and necessity of isolation and encourage patients to build confidence in recovery.
3. Emphasize that isolation is not only a way to better observe and treat patients, but also a way to protect loved ones and the society. Explain the treatment plans, and the effectiveness of the intervention.

- iii. Principles: Support and comfort. Treat patients with empathy, stabilize patients' emotions, and evaluate risks of suicide, self-injury, and aggressive behaviors early.
- b. Treatment stage of isolation.
 - i. Typical presentation: In addition to the above-mentioned possible presentation, there may also be loneliness, or lack of cooperation due to fear of the disease, abandonment of treatment, or excessive optimism and high expectations of treatment.
 - ii. Interventions:
 1. According to the patient's level of acceptance, objectively and truthfully explain the disease and the epidemic situation, so that the patient is informed;
 2. Assist the patient with communicating with loved ones and convey information if needed;
 3. Actively encourage patients to cooperate with all treatment activities;
 4. Make the environment suitable for the patients undergoing treatments;
 5. If necessary, please consult a psychiatric consultation.
 - iii. Principles: Actively communicate information and consult with psychiatrists if necessary.
- c. Patients with respiratory distress, extreme anxiety, and difficulty expressing themselves.
 - i. Typical presentation: near-death experiences, panic, despair, etc.
 - ii. Intervention measures: While calming and soothing, strengthen the treatment of the primary illness and reduce symptoms.
 - iii. Principles: soothing, sedation, pay attention to emotional communication, and enhance confidence in treatment.

d. Mild patients isolated at home, patients with fever who came to the hospital for treatment.

i. Typical presentation: panic, restlessness, loneliness, helplessness, depression, pessimism, anger, nervousness, stress from being alienated from others, grievance, shame, or disregard for illness.

ii. Interventions:

1. Assist patients to understand the situation with reliable information and knowledge, and trust scientific and authoritative medical sources;
2. Encourage active cooperation with treatment and isolation measures, healthy diet, balancing work and rest. Encourage soothing activities such as reading, listening to music, and communication using modern communication methods as well as other daily activities;
3. Accept isolation, mindful of their own reactions, and look for silver lining in adversity;
4. Seek social support to cope with stress: use modern communication methods to contact relatives, friends, colleagues, etc., and maintain social communication, for support and encouragement;
5. Encourage the use of psychological assistance hotline or online psychological intervention resources.

iii. Principles: Health education, encourage cooperation and adaptation to change.

2. Suspected Patients

a. Typical presentation: fluke-minded, avoiding treatment, fear of being discriminated against, or anxiety, seeking excessive treatment, frequent transfers among hospitals, etc.

b. Interventions:

- i. policy education, close observation, and early treatment;

- ii. adopt necessary protective measures;
 - iii. obey the overall public health strategy and report personal circumstances in accordance with regulations;
 - iv. adopt self-stress relieve methods and reduce stress
- c. Principles: Prompt education, adopt correct way of self-protection, obey the overall strategy, and reduce stress.

3. Health care and related personnel:

a. Typical presentation: Excessive fatigue and tension, exhaustion, anxiety, insomnia, depression, sadness, grievance, helplessness, down, frustration or self-blame in the face of patient death. Fear of being infected, concerned about family members, concerned about family members worried about themselves. Excessive excitement, refusal of reasonable rest, cannot well ensure their own health, etc.

b. Interventions:

- i. Conduct psychological crisis intervention training before participating in rescue, understand stress response, and learn how to respond to stress and regulate emotions. Conduct preventive interviews and openly discuss inner feelings; support and comfort; resource mobilization; help parties prepare psychologically for stress.
- ii. Eliminate the worries of frontline medical workers, arrange special personnel for logistical support, and staff in the quarantine area should rotate as much as possible every month.
- iii. Reasonable scheduling, arrange appropriate relaxation and rest, and ensure adequate sleep and diet. Try to arrange frontline staff at designated hospitals to stay near the hospital.

iv. Try to maintain contact and communication with family and the outside world when possible.

v. In case of insomnia, depression, or anxiety, you can seek professional psychological crisis intervention or mental health services. You can call the psychological assistance hotline or provide online psychological services, and face-to-face psychological crisis intervention can be performed in areas where conditions permit. It does not relieve for 2 weeks and affects the workers. It needs to be evaluated by a psychiatrist.

vi. If stress symptoms have occurred, you should promptly adjust your work position and seek professional help.

c. Principles: Rotate regularly, self-regulate, and ask for help if you have questions. 24.

4. Those who are in close contact with the patient (family members, colleagues, friends, etc.)

a. Typical presentations: avoidance, restlessness, anxiety during the waiting period; or blind bravery, refusal to protect and home observation.

b. Interventions:

i. Policy and education, encourage facing the reality, cooperate with home observation;

ii. Correct information dissemination and communication, release tension.

c. Principles: Mission, comfort and encourage communication through the Internet.

5. Patients who do not want to seek medical treatment

- a. Typical presentations: fear of being misdiagnosed and isolated, lack of awareness, avoidance, neglect, anxiety, etc.
- b. Interventions:
 - i. Education about the epidemic to eliminate fear;
 - ii. Advocate that seeking medical treatment early will benefit others;
 - iii. Eliminate public shaming and promote disease prevention based on scientific evidences;
- c. Principles: explain and persuade, not criticize. Support patients seeking medical treatments.

6. Susceptible people and the general public

- a. Typical presentations: panic, fear to go out, excessive disinfection, disappointment, fear, irritability, aggressive behavior, extreme optimism or pessimism, etc.
- b. Interventions:
 - i. Provide reliable information as well as information about further disease control methods and medical services;
 - ii. Ensure communication, provide guidance on how to adapt to changes;
 - iii. Do not discriminate against diagnosed or suspected patients;
 - iv. Advocate against unhealthy coping methods (such as drinking, smoking, etc.);
 - v. Health education so the general public can identify symptoms themselves.
- c. Principles: Health education, positive reinforcement, eliminate fear, prevention based on scientific evidences.