

References

1. Kinross P, Suetens C, Gomes Dias J, Alexakis L, Wijermans A, Colzani E, et al.; ECDC Public Health Emergency Team. Rapidly increasing cumulative incidence of coronavirus disease (COVID-19) in the European Union/European Economic Area and the United Kingdom, 1 January to 15 March 2020. *Euro Surveill.* 2020;25. <https://doi.org/10.2807/1560-7917.ES.2020.25.11.2000285>
2. CDC COVID-19 Response Team. Coronavirus disease 2019 in children – United States, February 12–April 2, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69:422–426 [Epub ahead of print]. <http://doi.org/10.15585/mmwr.mm6914e4>
3. Castagnoli R, Votto M, Licari A, Brambilla I, Bruno R, Perlino S, et al. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents: a systematic review. *JAMA Pediatr.* 2020 Apr 22 [Epub ahead of print]. <https://doi.org/10.1001/jamapediatrics.2020.1467>

Address for correspondence: Nicole E. Brown, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop H24-6, Atlanta, GA 30329-4027, USA; email: lvx6@cdc.gov

Severe Acute Respiratory Syndrome Coronavirus 2 among Asymptomatic Workers Screened for Work Resumption, China

Xiaoyu Han,¹ Xiong Wei,¹ Osamah Alwalid,¹ Yukun Cao, Yumin Li, Li Wang,² Heshui Shi²

Author affiliations: Union Hospital, Tongji Medical College, Wuhan, China (X. Han, O. Alwalid, Y. Cao, Y. Li, H. Shi); Hubei Province Key Laboratory of Molecular Imaging, Wuhan (X. Han, O. Alwalid, Y. Li, H. Shi); Wuhan Pingan Healthcare Diagnostic Center, Wuhan (X. Wei, L. Wang)

DOI: <https://doi.org/10.3201/eid2609.201848>

After the outbreak in Wuhan, China, we assessed 29,299 workers screened for severe acute respiratory syndrome coronavirus 2 by reverse transcription PCR. We noted 18 (0.061%) cases of asymptomatic infection; 13 turned negative within 8.0 days, and 41 close contacts tested negative. Among 6 contacts who had serologic tests, none were positive.

¹These first authors contributed equally to this article.

²These senior authors contributed equally to this article.

As the population of Wuhan, China, returns to work, asymptomatic cases of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) are being discovered among workers receiving health checkups for work resumption. Previous studies have shown that asymptomatic cases can be a public health threat and might lead to another outbreak (1,2). However, little is known about the clinical characteristics of asymptomatic infections. We report on cases of asymptomatic SARS-CoV-2 infection among persons during work resumption screening in Wuhan.

At Wuhan Pingan Healthcare Diagnostic Center, we reviewed 29,299 asymptomatic persons who were screened for SARS-CoV-2 by reverse transcription PCR (RT-PCR) and 22,633 asymptomatic persons tested for SARS-CoV-2 antibodies during March 13–April 25, 2020. Throat swab specimens were tested for SARS-CoV-2 by using Real-Time Fluorescent-PCR Kits (DAAN GENE Co., LTD, <https://www.en.daangene.com>; Appendix, <https://wwwnc.cdc.gov/EID/article/28/9/20-1848-App1.pdf>). We used colloidal gold-based immunochromatographic strip assay, Novel Coronavirus (SARS-CoV-2) IgM/IgG Antibody Detection Assay (Vazyme Biotech Co. Ltd., <http://vazyme.bioon.com.cn>) to perform antibody testing (Appendix). We recorded the demographic features, exposure history, RT-PCR and serology results, and imaging reports at the time of testing. We obtained follow-up data from persons screened by telephone.

Among 29,299 persons screened by RT-PCR, we confirmed 18 (0.061%) cases of SARS-CoV-2 infection. Of 22,633 persons tested for SARS-CoV-2 antibodies, 617 (2.7%) cases had positive IgG but negative IgM; 196 (0.87%) cases had positive IgG and IgM; and 40 (0.18%) cases had negative IgG but positive IgM.

The median age of 18 asymptomatic case-patients (10 male, 8 female) was 30.5 years (Table). Six (33.3%) cases had clear contact history with a confirmed case of SARS-CoV-2 infection. The median cycle threshold (C_t) values on the day of first positive RT-PCR were 38.2 (C_t range 37.2–39.3) for ORF1b gene and 38.1 (C_t range 36.81–38.5) for N gene (Table). All antibody tests were obtained on the day of first positive RT-PCR except in 1 case (obtained 6 days later). Half (7/14) the cases had negative IgM and IgG; the other half had positive IgG but negative IgM results (Table). Among 8 case-patients who had computed tomography imaging of the chest, none had remarkable findings. We closely observed the cases for 3–41 (median 16.5) days; 13 cases had negative RT-PCR within a median of 8 (range 3–14) days (Table), and none had symptoms. Among 41 close contacts, all had 2 consecutive negative RT-PCR tests ≥ 24 hours apart.

Among 6 contacts who had serologic tests, none had positive results.

According to Wuhan Municipal Health Commission (3), 275,400 RT-PCR tests were performed for universal screening during April 9–15, 2020. Among those, 182 (0.066%) asymptomatic persons were identified as SARS-CoV-2-positive, which is consistent with our study. Half the cases in our study showed negative IgM and IgG at the time of positive RT-PCR, suggesting recent infections (≤ 14 days). Seven (50%) cases in our study had positive IgG but negative IgM, indicating a late stage infection, 4 of which had a long interval of exposure (30–75 days). In addition, 13 cases had negative RT-PCR assays ≤ 8 (range 3–14) days, suggesting a favorable prognosis for persons with asymptomatic infections.

Epidemiologic, virologic, and modeling evidence support the possibility of SARS-CoV-2 transmission from persons who are presymptomatic (SARS-CoV-2 detected before symptoms onset) or asymptomatic (never develop symptoms) (4). None of the 18 asymptomatic persons in our study developed symptoms.

Recent reports showed that the viral load of SARS-CoV-2 infections in persons with no or mild symptoms was similar to the viral load of symptomatic patients (5,6), which could contribute to rapid transmissions (5). However, other studies demonstrated that asymptomatic patients had a lower viral load than symptomatic and presymptomatic patients (7,8,9), which might indicate less transmissibility from asymptomatic persons. The median cycle threshold values for the 18 cases were 38.2 for ORFa1b gene and 38.1 for the N gene, indicating a relatively low viral load. In addition, all 41 close contacts of the asymptomatic case-patients tested negative by RT-PCR. Possible explanations for this finding include that: the asymptomatic infected persons had relatively low viral load and were less infectious; that asymptomatic persons did not have clinical symptoms, such as sneezing or coughing, that could cause virus spread; and that, due to the strict isolation and preventive measures taken in Wuhan for ≥ 3 months, the population was generally protected from the spread of infection by mask-wearing and self-quarantine.

Table 1. Clinical characteristics of asymptomatic persons with detected severe acute respiratory syndrome coronavirus 2 infection, Wuhan, China*

| Age, y/sex | Occupation | Underlying condition | Interval, d† | Exposure source | RT-PCR | C _t ORFa1b | C _t N | Interval, d‡ | IgM | IgG | Chest CT | Interval, d§ | Follow-up, d |
|------------|----------------------|----------------------|--------------|-----------------|---------------|-----------------------|------------------|--------------|-----|-----|----------|--------------|--------------|
| 29/M | Office worker | N | 70 | Family | +, -, - | 37 | 38 | 5 | - | + | Normal | 0 | 41 |
| 32/M | Journalist | N | 54 | Bus passenger | +, +, - | 39 | 38 | 9 | - | + | Normal | 0 | 41 |
| 27/M | Journalist | N | NA | Unclear | +, - | 35 | 32 | 3 | - | - | Normal | 0 | 39 |
| 31/F | Researcher | N | 61 | Family | +, -, -, -, - | 36 | 38 | 7 | - | - | NA | NA | 23 |
| 27/M | Office worker | N | 75 | Family | +, -, -, - | 33 | 35 | 3 | - | + | Normal | 0 | 21 |
| 32/F | Housewife | N | NA | Unclear | +, -, - | 38 | 37 | 14 | - | - | NA | NA | 17 |
| 37/F | Office worker | N | 22 | Family | +, -, -, - | 40 | 39 | 8 | NA | NA | NA | NA | 16 |
| 57/M | Logistics dispatcher | Asthma | NA | Unclear | +, -, - | 39 | 37 | 8 | NA | NA | Normal | 10 | 16 |
| 25/M | Banking | N | NA | Unclear | +, -, - | 38 | 38 | 8 | NA | A | NA | NA | 16 |
| 26/M | Marketing | N | NA | Unclear | + | 39 | 38 | NA | - | - | NA | NA | 17 |
| 23/M | Marketing | N | NA | Unclear | +, - | 38 | 38 | 13 | - | - | Normal | 5 | 17 |
| 26/F | Office worker | N | NA | Unclear | +, - | 38 | 40 | 14 | NA | NA | NA | NA | 17 |
| 60/F | Cleaner | N | NA | Unclear | + | 37 | 37 | NA | - | + | NA | NA | 13 |
| 65/M | Cleaner | COPD | NA | Unclear | + | 39 | 39 | NA | - | - | Normal | 0 | 13 |
| 28/F | Office worker | N | NA | Unclear | + | 38 | 39 | NA | - | + | NA | NA | 8 |
| 33/F | Office worker | N | 30 | Family | +, - | 39 | 37 | 12 | - | + | Normal | 0 | 14 |
| 30/F | Office worker | N | NA | Unclear | +, - | 39 | 38 | 10 | - | + | NA | NA | 13 |
| 62/M | Worker | Hypertension | NA | Unclear | + | 39 | 39 | NA | - | - | NA | NA | 3 |

*COPD, chronic obstructive pulmonary disease; CT, computed tomography; C_t ORFa1b, RT-PCR cycle threshold for the ORFa1b gene; C_t N, RT-PCR cycle threshold for the N gene; N, no; NA, not available; +, positive; -, negative.

†Days between last exposure and initial positive RT-PCR.

‡Days between first positive and first negative RT-PCR results

§Days between first positive RT-PCR and CT scan.

Our report has limitations. Our sample size of asymptomatic cases is small, and follow-up was short. Recall bias of exposure history is another limitation; in the absence of clear symptom onset, asymptomatic persons might be less likely to accurately recall exposures than persons with symptoms. Finally, that the study took a place during the post-peak period of the epidemic in Wuhan, so contacts could have been seropositive already; those tested were seronegative, but most contacts did not have serologic testing.

In conclusion, as the population returns to the workplace, asymptomatic SARS-CoV-2-infected persons could be among workers. Although we did not detect transmission among 41 contacts of persons who were SARS-CoV-2-positive, such transmission cannot be excluded. Therefore, continued testing, self-quarantine, and mask-wearing should be encouraged to reduce the risk for additional outbreaks.

Acknowledgments

We thank all our colleagues for helping us during the current study.

This study was supported by Zhejiang University special scientific research fund for COVID-19 prevention and control, the Huazhong University of Science and Technology (HUST) COVID-19 Rapid Response Call (grant no. 2020kfyXGYJ019).

About the Author

Dr. Han is a clinician in the Department of Radiology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China. Her research interests including image diagnosis of pneumonia and lung cancer.

References

1. Qiu J. Covert coronavirus infections could be seeding new outbreaks. *Nature*. 2020 Mar 20 [Epub ahead of print]. <https://doi.org/10.1038/d41586-020-00822-x>
2. Pan A, Liu L, Wang C, Guo H, Hao X, Wang Q, et al. Association of public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan, China. *JAMA*. 2020;323:1-9. <https://doi.org/10.1001/jama.2020.6130>
3. Wuhan Municipal Health Commission. CN-HEALTHCARE: Hubei health, 2020 April 19 [in Chinese] [cited 2020 May 25]. <https://www.cn-healthcare.com/articlewm/20200419/content-1105403.html>
4. Furukawa NW, Brooks JT, Sobel J. Evidence supporting transmission of severe acute respiratory syndrome coronavirus 2 while presymptomatic or asymptomatic. *Emerg Infect Dis*. 2020 May 4 [Epub ahead of print]. <https://doi.org/10.3201/eid2607.201595>
5. Arons MM, Hatfield KM, Reddy SC, Kimball A, James A, Jacobs JR, et al. Presymptomatic SARS-CoV-2 infections and transmission in a skilled nursing facility. *N Engl J Med*. 2020;382:2081-90. <https://doi.org/10.1056/NEJMoa2008457>
6. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med*. 2020;382:1177-9. <https://doi.org/10.1056/NEJMc2001737>
7. Kim SE, Jeong HS, Yu Y, Shin SU, Kim S, Oh TH, et al. Viral kinetics of SARS-CoV-2 in asymptomatic carriers and presymptomatic patients. *Int J Infect Dis*. 2020;95:441-3. <https://doi.org/10.1016/j.ijid.2020.04.083>
8. He D, Zhao S, Lin Q, Zhuang Z, Cao P, Wang MH, et al. The relative transmissibility of asymptomatic COVID-19 infections among close contacts. *Int J Infect Dis*. 2020;94:145-7. <https://doi.org/10.1016/j.ijid.2020.04.034>
9. Zhou R, Li F, Chen F, Liu H, Zheng J, Lei C, et al. Viral dynamics in asymptomatic patients with COVID-19. *Int J Infect Dis*. 2020;96:288-290. <https://doi.org/10.1016/j.ijid.2020.05.03>

Address for correspondence: Heshui Shui, Department of Radiology, Union Hospital, Tongji Medical College, Wuhan 430030, China; email: heshuishui@hust.edu.cn

Effects of Proactive Social Distancing on COVID-19 Outbreaks in 58 Cities, China

Zhanwei Du, Xiaoke Xu, Lin Wang, Spencer J. Fox, Benjamin J. Cowling, Alison P. Galvani, and Lauren Ancel Meyers

Author affiliations: The University of Texas at Austin, Austin, Texas, USA (Z. Du, S.J. Fox, L.A. Meyers); Dalian Minzu University, Dalian, China (X. Xu); University of Cambridge, Cambridge, UK (L. Wang); The University of Hong Kong, Hong Kong, China (B.J. Cowling); Center for Infectious Disease Modeling and Analysis, Yale School of Public Health, New Haven, Connecticut, USA (A.P. Galvani); Santa Fe Institute, Santa Fe, New Mexico, USA (L.A. Meyers)

DOI: <https://doi.org/10.3201/eid2609.201932>

Cities across China implemented stringent social distancing measures in early 2020 to curb coronavirus disease outbreaks. We estimated the speed with which these measures contained transmission in cities. A 1-day delay in implementing social distancing resulted in a containment delay of 2.41 (95% CI 0.97–3.86) days.