

the low prevalence of SARS-CoV-2 infection. We do not have a clear explanation for how the 2 residents became infected after the introduction of these measures; we were unable to determine whether surveys were useful tools. It is possible that routine testing discouraged persons with symptoms from visiting. We observed a very low rate of positive tests in the LTCF staff; only 1 staff member tested positive. Potential explanations for this low rate could be that testing had an impact on behavior, symptom screening kept ill staff home, or the virus was less prevalent in the community surrounding the LTCF. Although symptom surveys were used and absentee rates were normal, staff did not report symptoms as a reason for missed work. Despite these limitations, this study suggests that a proper testing strategy coupled with other measures may result in protection of vulnerable populations.

About the Author

Dr. Shimotsu is an epidemiologist at United Health Group. His research areas include clinical care model development, chronic disease treatment, and social determinants of chronic disease.

References

1. Lai C-C, Wang J-H, Ko W-C, Yen M-Y, Lu M-C, Lee C-M, et al. COVID-19 in long-term care facilities: an upcoming threat that cannot be ignored. *J Microbiol Immunol Infect*. 2020;53:444–6. <https://doi.org/10.1016/j.jmii.2020.04.008>
2. Wortham JM, Lee JT, Althomsons S, Latash J, Davidson A, Guerra K, et al. Characteristics of persons who died with COVID-19—United States, February 12–May 18, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:923–9. <https://doi.org/10.15585/mmwr.mm6928e1>
3. Fisman DN, Bogoch I, Lapointe-Shaw L, McCreedy J, Tuite AR. Risk factors associated with mortality among residents with coronavirus disease 2019 (COVID-19) in long-term care facilities in Ontario, Canada. *JAMA Netw Open*. 2020;3:e2015957. <https://doi.org/10.1001/jamanetworkopen.2020.15957>
4. Rossen LM, Branum AM, Ahmad FB, Sutton P, Anderson RN. Excess deaths associated with COVID-19, by age and race and ethnicity—United States, January 26–October 3, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:1522–7. <https://doi.org/10.15585/mmwr.mm6942e2>
5. McMichael TM, Clark S, Pogosjans S, Kay M, Lewis J, Baer A, et al.; Public Health – Seattle & King County, EvergreenHealth, and CDC COVID-19 Investigation Team. COVID-19 in a long-term care facility—King County, Washington, February 27–March 9, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69:339–42. <https://doi.org/10.15585/mmwr.mm6912e1>

Address for correspondence: Daniel Griffin, Columbia University Irving Medical Center—Medicine, 701 W 168th St, HHSC 1310, New York, NY 10032-3784, USA; email: DGriffin@ProHEALTHcare.com

Severe Acute Respiratory Syndrome Coronavirus 2 Outbreak Related to a Nightclub, Germany, 2020

Nadine Muller,¹ Mareike Kunze,¹ Fabienne Steitz, Neil J. Saad, Barbara Mühlemann, Jörn I. Beheim-Schwarzbach, Julia Schneider, Christian Drosten, Lukas Murajda, Sandra Kochs, Claudia Ruscher, Jan Walter, Nadine Zeitlmann,² Victor M. Corman²

Author affiliations: European Centre for Disease Prevention and Control, Stockholm, Sweden (N. Muller, N.J. Saad); Robert Koch Institute, Berlin, Germany (N. Muller, N.J. Saad, J. Walter, N. Zeitlmann); Charité—Universitätsmedizin Berlin, Berlin (N. Muller, B. Mühlemann, J.I. Beheim-Schwarzbach, J. Schneider, C. Drosten, V.M. Corman); Local Health Authority Berlin-Mitte, Berlin (M. Kunze, F. Steitz, L. Murajda, S. Kochs); German Centre for Infection Research, Berlin (B. Mühlemann, C. Drosten, V.M. Corman); State Office for Health and Social Affairs, Berlin (C. Ruscher)

DOI: <https://doi.org/10.3201/eid2702.204443>

We report an outbreak of coronavirus disease with 74 cases related to a nightclub in Germany in March 2020. Staff members were particularly affected (attack rate 56%) and likely caused sustained viral transmission after an event at the club. This outbreak illustrates the potential for superspreader events and corroborates current club closures.

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) superspreading events are particularly linked to indoor settings, such as religious venues (1), restaurants (2), and bars or nightclubs (3–6). To provide further details on the extent and transmission dynamics in nightclubs, we describe a SARS-CoV-2 outbreak related to a Berlin, Germany, nightclub during the early phase of the coronavirus disease (COVID-19) pandemic, before infection prevention measures were applied.

On March 5, 2020, contact tracing activities in Berlin revealed several COVID-19 cases linked by visiting the same nightclub, club X, on February 29, 2020 (event 1). Estimates suggest ≈300 guests attended event 1. Club X then held other events: event 2 with ≈150 guests on March 2 and event 3 with ≈200 guests on March 5. On March 6, the local health

¹These first authors contributed equally to this article.

²These senior authors contributed equally to this article.

authority of Mitte district, Berlin, published announcements in local newspapers and on social media to identify other attendees of the events. Everyone attending ≥ 1 event was categorized as a high-risk contact person and ordered to self-quarantine for 14 days. If symptoms occurred, laboratory testing was recommended. Mandatory case notification occurred from the laboratory to the local health authority based on Germany's Protection against Infection Act (7). Due to the increasing spread of COVID-19, on March 16, 2020, government authorities in Germany prohibited social gatherings, including events in nightclubs, until further notice.

Confirmed cases in the outbreak were defined as persons with laboratory-confirmed SARS-CoV-2 (Appendix, <https://wwwnc.cdc.gov/EID/article/27/2/20-4443-App1.pdf>). We retrieved dates of symptom onset and sociodemographic data of 64 outbreak cases from the national infectious diseases notification database. We considered staff and persons who attended any event at club X to have first-generation cases and their contacts to have second-generation cases.

We interviewed 44 persons with first-generation cases whose contact information was available and with all 16 club X staff members who

worked any of the 3 events. For staff members who were not tested after the events or who tested negative despite reporting symptoms, we offered SARS-CoV-2 antibody testing 3 months after the outbreak to ascertain their infection status. We also mapped the space inside club X (Appendix Figure 1).

In total, 74 reported cases were linked to the outbreak. Median age was 30 (range 2–63) years; cases were equally distributed by sex, 37 female (50%) and 37 male (50%). Among 41 first-generation cases with known date of symptom onset and only 1 exposure, the median incubation period was 4 days (interquartile range 3–6 days). The calculated attack rates (ARs) show that guests attending event 1 were particularly affected. Staff pooled over all events had the highest risk for infection (AR 56%) (Table).

Among guests, 1 PCR-confirmed case had self-reported initial symptoms 1 day before attending event 1 and could be a potential source of the outbreak. The most probable source for continued viral transmission at event 3 was a PCR-confirmed case in a staff member working event 1 and event 3, with symptom onset 1 day before event 3. Overall, staff members reported symptom onset at a later stage of the outbreak than guests (Figure).

Table. Calculated attack rates for identified coronavirus disease outbreak cases among staff members and guests attending events in a nightclub, Berlin, Germany, March 2020*

Characteristics	Cases, no. (%)	No. attending		
		Event 1	Event 2	Event 3
Estimated guests†	–	300	150	200
Staff members, n = 16‡	–	11	6	11
Total cases	74 (100)			
Cases by generation§				
First-generation, n = 55	55 (74.3)			
Guests¶	46 (83.6)	39	0	3
Staff	9 (16.4)	–	–	–
Second-generation, n = 10	10 (13.5)	–	–	–
Generation unknown, n = 9	9 (12.2)	–	–	–
Cases by case definition#				
Confirmed cases, n = 72	72 (97.3)	–	–	–
PCR-confirmed	70 (97.2)	–	–	–
Antibody testing-confirmed	2 (2.8)	–	–	–
Probable cases	2 (2.7)	–	–	–
Attack rate, %**	Pooled over all events	Event 1	Event 2	Event 3
Guests	–	13	–	2
Staff	56	–	–	–

*Event-related case numbers are shown only for first-generation guest cases as all of them confirmed to only have attended 1 of the 3 events. –, value not calculated.

†The exact number of guests attending the events is unknown. For event 1, an estimate of attending guests was based on the maximum capacity of the club; staff and contacted guests confirmed that the club was running at full capacity. For events 2 and 3, the club owner provided estimates listed here.

‡Most staff members attended ≥ 2 of the events.

§First-generation cases were defined as cases exposed during event 1, 2, or 3. Second-generation cases were defined as cases without exposure at club X but with exposure to first-generation cases. Cases of unknown generation were confirmed cases of the outbreak but without contact information to reveal whether they were first- or second-generation cases.

¶All guests contacted confirmed they attended only 1 of the 3 events. Information on the event of exposure was available for 42 first-generation cases among guests. No guest case reported visiting club X for event 2.

#The outbreak case definition is described in the Appendix (<https://www.nc.cdc.gov/EID/article/27/2/20-4443-App1.pdf>). All cases confirmed by antibody testing were otherwise probable cases.

**Calculation of primary attack rates for guests was based on approximations for the denominator, the number of guests attending. Because most staff members were exposed repeatedly while working at ≥ 1 event we separately calculated attack rates for staff pooled over all events.




Figure. Date of symptom onset among 64 coronavirus disease cases linked to an outbreak in a nightclub, Berlin, Germany, March 2020. The asterisks indicate cases with symptom onset prior attending event 1 (symptom onset on February 28, 2020) and event 3 (symptom onset on March 4, 2020). No guests among cases reported attending event 2, but all attended either event 1 or event 3. No staff among cases attended only event 2; all attended event 1, event 3, or both events.

SARS-CoV-2 whole-genome sequencing was performed on 17 available patient samples to assess clustering of sequences. Sequencing revealed that 10 cases among event 1 guests, 2 second-generation cases, and 5 cases of unknown generation all grouped within clade G (GISAID, <https://www.gisaid.org>) and B.1 (Pangolin clade naming) (Appendix Figure 2). This clade also was observed in the SARS-CoV-2 outbreak in Italy and many later outbreaks in Europe (8). Sequences from 11 samples were identical. The other 6 samples were otherwise identical, but had slight differences; 1 sequence had 1 position with ambiguous nucleotides; 3 other sequences had 3 positions with ambiguous nucleotides; 1 sequence had a substitution in the 3' untranslated region; and sequences from 2 cases, in a couple who attended event 1, had an identical substitution in the N gene (Appendix Table 1). This substitution could hint to a second independent transmission cluster comprising these 2 cases, but all observed sequence variants also can be explained by sporadic mutation events. Thus, the sequence data do not provide evidence against a single person as the outbreak source (Appendix Figure 2).

The large number of cases from event 1, the relatively low median incubation period (4 days) for first-generation cases, and the close genetic relatedness of the sequenced viruses corroborate the theory of transmission from a single person and the potential for superspreading in a nightclub when no social distancing measures are applied. This outbreak further illustrates the potential role of nightclub staff members in transmission. AR among staff was particularly high (56%), showing they had a particularly high risk

for infection. Because 1 staff member appears to have been infected at event 1, then worked with symptoms at event 3, continued viral transmission could have been caused by staff. However, without sequencing data for all cases, staff contribution to viral transmission cannot be confirmed. Nonetheless, once ease of restrictions is considered, our study suggests that infection protection should be targeted particularly toward staff in nightclubs and bars.

Acknowledgments

We thank all our colleagues in the local health authority in Berlin and other federal states who actively collaborated in case finding activities, especially the team from the local health authority in the Berlin district Mitte for their unremitting commitment in managing cases and contact persons during the outbreak. We also thank the affected nightclub's owner and staff members for their trustful sharing of information. This study was only possible thanks to the collaboration of the affected individuals from the outbreak. We also gratefully acknowledge the authors, originating and submitting laboratories of the genetic sequence and metadata made available through GISAID on which the phylogenetic analysis presented in this paper is based.

N.M. is a fellow of Postgraduate Training for Applied Epidemiology, supported financially by the Robert Koch Institute. N.J.S. is a fellow of the European Centre for Disease Prevention and Control (ECDC) Fellowship Programme, supported financially by the ECDC. Contributions by V.M.C. and C.D. were funded by the German Ministry of Health (Konsiliarlabor für Coronaviren) and the German Center for Infection Research (DZIF).

About the Author

Dr. Muller is a medical doctor working as a fellow of the Postgraduate Training for Applied Epidemiology at the Robert Koch Institute, Berlin, and has a research affiliation with the Department of Infectious Diseases and Respiratory Medicine, Charité University Medicine Berlin. Her main research interests include epidemiological investigations on respiratory diseases.

References

1. Leclerc QJ, Fuller NM, Knight LE, CMMID COVID-19 Working Group, Funk S, Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters? *Wellcome Open Res.* 2020;5:83. <https://doi.org/10.12688/wellcomeopenres.15889.2>
2. Lu J, Gu J, Li K, Xu C, Su W, Lai Z, et al. COVID-19 outbreak associated with air conditioning in restaurant, Guangzhou, China, 2020. *Emerg Infect Dis.* 2020;26:1628–31. <https://doi.org/10.3201/eid2607.200764>
3. Abbott S, Hellewell J, Thompson RN, Sherratt K, Gibbs HP, Bosse NI, et al. Estimating the time-varying reproduction number of SARS-CoV-2 using national and subnational case counts. *Wellcome Open Res.* 2020;5:112. <https://doi.org/10.12688/wellcomeopenres.16006.1>
4. Correa-Martínez CL, Kampmeier S, Kumpers P, Schwierzeck V, Hennies M, Hafezi W, et al. A pandemic in times of global tourism: superspreading and exportation of COVID-19 cases from a ski area in Austria. *J Clin Microbiol.* 2020;58:3. <https://doi.org/10.1128/JCM.00588-20>
5. Kang CR, Lee JY, Park Y, Huh IS, Ham HJ, Han JK, et al.; Seoul Metropolitan Government COVID-19 Rapid Response Team (SCoRR Team). Coronavirus disease exposure and spread from nightclubs, South Korea. *Emerg Infect Dis.* 2020;26:2499–501. <https://doi.org/10.3201/eid2610.202573>
6. Chau NVV, Hong NTT, Ngoc NM, Thanh TT, Khanh PNQ, Nguyet LA, et al. Superspreading event of SARS-CoV-2 infection at a bar, Ho Chi Minh City, Vietnam. *Emerg Infect Dis.* 2020 Oct 16 [Epub ahead of print]. <https://doi.org/10.3201/eid2701.203480>
7. Federal Ministry for Justice and Consumer Protection. Law on the Prevention and Control of Infectious Diseases in Humans [in German] [cited 2020 Nov 27]. <https://www.gesetze-im-internet.de/ifsg/index.html>
8. Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, et al. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nat Microbiol.* 2020;5:1403–7. <https://doi.org/10.1038/s41564-020-0770-5>

Address for correspondence: Nadine Muller, Department of Infectious Disease Epidemiology, Robert Koch Institute, Seestrasse 10, Berlin 13353, Germany; email: mullern@rki.de

Evidence of SARS-CoV-2 RNA in an Oropharyngeal Swab Specimen, Milan, Italy, Early December 2019

Antonella Amendola,¹ Silvia Bianchi,¹ Maria Gori, Daniela Colzani, Marta Canuti, Elisa Borghi, Mario C. Raviglione, Gian Vincenzo Zuccotti, Elisabetta Tanzi

Author affiliations: University of Milan, Milan, Italy (A. Amendola, S. Bianchi, M. Gori, D. Colzani, E. Borghi, M.C. Raviglione, G.V. Zuccotti, E. Tanzi); Memorial University of Newfoundland, St. John's, Newfoundland, Canada (M. Canuti)

DOI: <https://doi.org/10.3201/eid2702.204632>

We identified severe acute respiratory syndrome coronavirus 2 RNA in an oropharyngeal swab specimen collected from a child with suspected measles in early December 2019, ≈3 months before the first identified coronavirus disease case in Italy. This finding expands our knowledge on timing and mapping of novel coronavirus transmission pathways.

Coronavirus disease (COVID-19) symptoms can encompass a Kawasaki disease-like multisystem inflammatory syndrome and skin manifestations that accompany common viral infections such as chickenpox and measles (1,2). Some of the earliest reports of COVID-19 cutaneous manifestations came from dermatologists in Italy. In fact, Italy was the first Western country severely hit by the COVID-19 epidemic. The first known COVID-19 case in Italy was reported in the town of Codogno in the Lombardy region on February 21, 2020. However, some evidence suggests that severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) had been circulating unnoticed for several weeks in Lombardy before the first official detection (3). Phylogenetic studies highlighted an early circulation of SARS-CoV-2 in Italy and suggest multiple introductions of the virus from China and Germany, followed by an autochthonous transmission (4,5). Furthermore, environmental surveillance has unequivocally demonstrated the presence of the virus, at concentrations comparable to those obtained from samples collected at later stages of the pandemic, in the untreated wastewater of the Milan area as early as mid-December 2019 (6).

¹These authors contributed equally to this article.

Severe Acute Respiratory Syndrome Coronavirus 2 Outbreak in a Nightclub, Germany, 2020

Appendix

Outbreak Case Definition

A confirmed case in the outbreak was defined as any person with laboratory-confirmed severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection who attended club X between February 29 and March 5, 2020; had an epidemiologic link to a case that attended club X between February 29 and March 5, 2020; or both. A probable case was defined as any person with clinical symptoms of coronavirus disease (COVID-19) (*I*) who attended club X between February 29 and March 5, 2020; had an epidemiologic link to a case that attended club X between February 29 and March 5, 2020; or both.

Laboratory confirmation of SARS-CoV-2 was defined by the detection of SARS-CoV-2 nucleic acid via PCR in a clinical specimen or by detection of SARS-CoV-2–specific IgG antibodies. Because club X was closed on March 6, 2020 until further notice, no cases linked to the outbreak were reported after March 2020.

Cases were assigned to the outbreak if they fulfilled the case definition and confirmation of positive case status was given by the person or by the local health authority of their place of residence.

Epidemiologic Outbreak Investigation

Data on the day of symptom onset was retrieved from the national infectious diseases notification database, which was collected and notified by public health officials from local public health authorities. Among all cases linked to the outbreak, dates of symptom onset were available for 64 cases.

We conducted semistructured telephone interviews with first-generation cases to gather information related to their exposure in the club, prior travel history, and characteristics of clinical symptoms. Among all first-generation cases linked to the outbreak, contact information was available for 44 cases and the study team interviewed them. We performed analysis by time, generation, symptoms, sex, and age. For analysis by time, we stratified cases by guests, staff members, and generation. For continuous variables, if not normally distributed, we calculated medians and interquartile ranges (IQR). In addition, members of the outbreak investigation team performed a site visit of club X to gain insight into the outbreak setting (Appendix Figure 1).

Virological Outbreak Investigation

SARS-CoV-2 Antibody Screening of Staff Members

For laboratory-confirmation of cases, qualitative real-time RT-PCR for SARS-CoV-2 was performed on purified RNA from swabs as described (2), or the Cobas SARS-CoV-2 test (Roche, <https://www.roche.com>), both of which target the SARS-CoV-2 E gene.

For nightclub staff members who had negative PCR tests for SARS-CoV-2 or who were not tested after the exposure, we performed SARS-CoV-2 antibody screening during June 2–24, 2020, approximately 3 months after the outbreak, by using a 2-step approach. First, we screened samples by using Anti-SARS-CoV-2 S1 IgG and IgA ELISAs (Euroimmun, <https://www.euroimmun.com>) according to the manufacturer's protocol. Second, we performed a plaque reduction neutralization test (PRNT), as previously described (3,4). In the PRNT, we tested all dilutions in duplicate. Only serum samples showing an optical density ratio >0.8 in the IgA or IgG ELISA were considered reactive and tested in the PRNT.

Whole-Genome Sequencing

To investigate the sequence diversity of the outbreak, we performed whole-genome sequencing (WGS) on available samples from initial diagnostic testing that had sufficient sample material. For WGS we followed two approaches. First, we performed direct sequencing of native samples with a high viral load (cycle threshold [C_t] value <25); then, for samples with lower SARS-CoV-2 concentration, we used a PCR amplicon-based sequencing approach.

For sequencing of native samples with a high viral load (C_t value <25), we used ≤ 100 ng in 5 μ L of extracted RNA for library preparation by using the KAPA RNA Hyper Prep Kit

(Roche Molecular Diagnostics, <https://diagnostics.roche.com>) according to manufacturer's instructions. The RNA was fragmented for 6 min at 85°C. Indexed libraries were then amplified for 8–13 PCR cycles. All DNA libraries were measured by Qubit dsDNA HS Assay kit (Thermo Fisher Scientific, <https://www.thermofisher.com>), pooled together at equimolar ratios, and normalized.

For amplicon-based complete genome sequencing of samples with a lower viral load (C_t value ≥ 25) we followed 2 approaches. First, we used 108 SARS-CoV-2 whole genomes, available in early February 2020 to design 48 overlapping heminested PCR fragment primers. Fragment size ranged between 507 bp and 950 bp for first-round products and 414–877 bp for second-round products. Primer names including “i” were modified versions (Appendix Table 2). For the first-round PCR, a 25 μ L reaction was performed by using the SuperScript III One-Step RT-PCR System with Platinum Taq DNA (Invitrogen, <https://www.thermofisher.com>) with 5 μ L of RNA, 12.5 μ L of 2 \times reaction buffer (provided with the kit), 1 μ L of enzyme mixture from the kit, additional 0.4 μ L of a 50 mmol magnesium sulfate solution, 400 nmol concentrations of each first-round primer, and 1 μ g of bovine serum albumin (Roche). For the second-round, 50 μ L reactions were carried out by using the Platinum Taq Polymerase Kit (Invitrogen), with 1 μ L of the first-round PCR product, 5 μ L of 10 \times reaction buffer provided with the kit, 2.5 mmol MgCl₂, 200 μ M of each dNTP, 0.2 μ L of Platinum Taq, and 400 nmol of each second-round primer. First-round RT-PCRs were carried out by using a thermocycling protocol with reverse transcription at 55°C for 20 min and subsequent PCR at 95°C for 3 min, followed by 45 cycles of 95°C for 15 s, 56°C for 15 s, and 72°C for 55 s, followed by a final 2-min extension step at 72°C. Second-round reactions used the same cycling protocol but without the RT step. Second, for amplicon-based WGS we used random hexamers and the SuperScript III Reverse transcription kit (Invitrogen) according to manufacturer's instructions, then amplified the SARS-CoV-2 genome by using the primer sets (V1) published by the Artic Network ([dx.doi.org/10.17504/protocols.io.bdbfi2jn](https://doi.org/10.17504/protocols.io.bdbfi2jn)). A 25 μ L PCR master mix was set up by using the Q5 High-Fidelity DNA Polymerase kit (New England Biolabs, <https://www.neb.com>) with 5 μ L 5 \times Q5 Reaction Buffer (New England Biolabs), 13.15 μ L RNase-free water, 0.5 10 mmol dNTPs, 3.6 μ L of either 10 μ mol primer pool 1 or 2, 2.5 μ L cDNA and 0.25 μ L Q5 High-Fidelity DNA Polymerase (New England Biolabs). PCR was carried out by using a thermocycling protocol with initial denaturation at 98°C for 30 sec, followed by 35 cycles of

98°C for 15 s, 65°C for 2 min 30 sec, followed by a final 2-min extension step at 72°C. PCR products were pooled and purified by using KAPA Pure Beads (Roche Molecular Diagnostics) according to manufacturer's instructions.

For DNA library preparation of the purified PCR amplicons, we used ≤ 5 ng DNA and the KAPA Hyper Prep Kit (Roche Molecular Diagnostics). All pooled PCR amplicons and DNA libraries were measured by Qubit dsDNA HS Assay kit (Thermo Fisher Scientific).

Sequencing was performed by using the 600-cycle MiSeq reagent v3 cartridge (Illumina, <https://www.illumina.com>), the 150/300-cycle NextSeq, and 100-cycle NovaSeq (Illumina) according to manufacturer's instructions (Appendix Table 3).

Bioinformatic Sequence Analysis

For each sample, data from the individual sequencing runs were mapped to a reference sequence (GISAID accession no. EPI_ISL_402125, GenBank accession no. NC_045512.2) by using bowtie2 version 2.3.5.1 and the sensitive-local option (5). Duplicates were removed using GATK MarkDuplicatesSpark version 4.1.4.1 (6), and the consensus reads were called at positions with coverage ≥ 3 reads by using bcftools version 1.10.2–31-gffa7016 and bcftools call–ploidy 1-mv-Oz-o (<https://samtools.github.io/bcftools/bcftools.html>). We included the following sequences in the phylogenetic tree: 1 sequence from each clade assigned by Pangolin; all sequences from Germany sampled before April 16, 2020 and available in GISAID on July 22, 2020; and representative sequences from GISAID clade G sampled by April 15, 2020 and available in GISAID on July 22, 2020. Sequences from each country were clustered by using CD-HIT version 4.8.1 by using a sequence identity threshold of 0.99 (7) and we picked 1 sequence from each cluster. Then we included 4 sequences from the U.S. and 1 from Canada that have the same additional SNP as sequences ChVir-W1248–16 and ChVir-D715-D799–17 from this outbreak. The phylogenetic tree was inferred by using RAxML-ng version 0.7.0 BETA (8) and an HKY substitution model, with gamma distribution rate heterogeneity among sites and invariant sites. We performed 100 bootstrap replicates and created a phylogenetic tree by using baltic (9) (Appendix Figure 2).

Ethics Approval

The outbreak investigation was conducted within the framework of the German Infection Protection Act (10) as part of an outbreak response and public health practice. Mandatory

regulations were respected, and thus review by an ethics committee was not required. Support by the Robert Koch Institute was provided after official request. Participation in the questionnaire and blood specimen collection for antibody testing was voluntary, for which verbal consent was obtained. For antibody testing, additional written informed consent was obtained.

Description of the Outbreak Setting

Club X is located in a basement. The area accessible to guests is $\approx 150 \text{ m}^2$ with a height of $\approx 3 \text{ m}$ (Appendix Figure 1). Ventilation of the space is ensured by a mechanical air exhaust and supply system and maintenance was performed according to the manufacturer's instructions. To avoid noise pollution in the surrounding neighborhood, windows are usually closed during events.

Clinical Symptoms of Cases

Among a total of 74 cases linked to the outbreak, dates of symptom onset were available for 64 cases. Of those, 44 cases could be interviewed on clinical symptoms during their COVID-19 infection. All 44 cases reported having ≥ 1 symptom. The most common symptoms experienced were dysgeusia (65%), cough (61%), headache (58%), and dysosmia (58%) (Appendix Table 4).

References

1. European Centers for Disease Control. Case definition for coronavirus disease 2019 (COVID-19), as of 29 May 2020 [cited 2020 Oct 21]. <https://www.ecdc.europa.eu/en/covid-19/surveillance/case-definition>
2. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill.* 2020;25. [PubMed](https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045)
<https://doi.org/10.2807/1560-7917.ES.2020.25.3.2000045>
3. Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature.* 2020;581:465–9. [PubMed](https://doi.org/10.1038/s41586-020-2196-x)
<https://doi.org/10.1038/s41586-020-2196-x>

4. Buchholz U, Müller MA, Nitsche A, Sanewski A, Wevering N, Bauer-Balci T, et al. Contact investigation of a case of human novel coronavirus infection treated in a German hospital, October-November 2012. *Euro Surveill.* 2013;18:20406. [PubMed](#)
5. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012;9:357–9. [PubMed](#) <https://doi.org/10.1038/nmeth.1923>
6. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernysky A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 2010;20:1297–303. [PubMed](#) <https://doi.org/10.1101/gr.107524.110>
7. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics.* 2006;22:1658–9. [PubMed](#) <https://doi.org/10.1093/bioinformatics/btl158>
8. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics.* 2019;35:4453–5. [PubMed](#) <https://doi.org/10.1093/bioinformatics/btz305>
9. baltic - Python library for parsing phylogenetic trees [cited 2020 Sep 8]. <https://github.com/evogytis/baltic>
10. Federal Ministry for Justice and Consumer Protection. Law on the Prevention and Control of Infectious Diseases in Humans [in German] [cited 2020 11 27]. <https://www.gesetze-im-internet.de/ifsg/index.html>

Appendix Table 1. Mapping statistics and single nucleotide polymorphisms of severe acute respiratory syndrome coronavirus 2 from an outbreak associated with a nightclub, Berlin, Germany, March 2020*

ID	Coverage depth†	Genome coverage, %	Ambiguous positions*	SNPs relative to the majority*	Sampling date, 2020
ChVir-D712-1	971.6 (3-6,636)	100	None	None	Mar 7
ChVir-D666-2	85.8 (4-307)	100	None	None	Mar 7
ChVir-D718-3	482.9 (3-6,898)	100	None	None	Mar 7
ChVir-D672-4	214 (3-487)	100	None	None	Mar 7
ChVir-D658-5	58.9 (3-206)	100	None	None	Mar 8
ChVir-D665-6	2,342.1 (3-5,966)	100	None	None	Mar 6
ChVir-D667-7	294.9 (3-773)	100	None	None	Mar 7
ChVir-D671-8	41.3 (2-107)	100	None	None	Mar 7
ChVir-D670-9	13.3 (0-51)	99.8	None	None	Mar 7
ChVir-D710-10	2,273.3 (34-4,992)	100	Position 20469 R, 2,144 reads have A, 1,573 reads have G. Confirmed by Sanger sequencing. 3rd codon position, synonymous.	None	Mar 7
ChVir-D711-11	2,924.1 (0-6,965)	99.8	Position 16570 K, 7,031 reads have G, 156 reads have T, but Sanger suggests T. 1st codon position, non-synonymous, C>G; Position 21515 R, 856 reads have A, 864 reads have G, 2nd codon position, non-synonymous, N > S; Position 25419 R; 4,657 reads have A, 6,708 reads have G, 3rd codon position, synonymous	Position 29780 A>G, 3' UTR	Mar 7
ChVir-D717-D761-12	3,645.9 (3-5,927)	100	Position 14801 R (reference has A), 93 reads have G, 122 reads have A. 2nd codon position, D>G change.	None	Mar 7
ChVir-W1191-13	96.6 (3-427)	100	None	None	Mar 8
ChVir-D679-14	52.6 (3-225)	100	None	None	Mar 4
ChVir-D929-15	3,791.3 (0-427)	98	Position 545 K, 4,591 reads have G, 7,074 have T confirmed by Sanger, 1st codon position, non-synonymous, G>C	None	Mar 5
ChVir-W1248-16	132.2 (0-422)	99	None	Position 29254 G>T, 3rd codon position, synonymous	Mar 7
ChVir-D715-D799-17	4,553.4 (3-6646)	100	None	Position 29254 G>T, 3rd codon position, synonymous	Mar 6

*Positions are given relative to the reference genome, EPI_ISL402125 (Wuhan-1).

†Coverage depth indicates the mean number of reads covering each position in the genome. Numbers in parentheses show the minimum and maximum number of reads.

Appendix Table 2. Oligonucleotide used for amplification and sequencing of SARS-CoV-2 genome fragments

Primer ID	Sequence (5'→3')	Use
SARS2_1_F	CAACTTTCGATCTCTTGTAGATCTG	1st round
SARS2_1_Fnest	GTCACCTCGGCTGCATGCTTAGTG	2nd round
SARS2_1_R	GTTATCGACATAGCGAGTGTATGC	1st round and 2nd round
SARS2_2_F	GGAGCTGGTGGCCATAGTTACG	1st round
SARS2_2_Fnest	CATTTGACTTAGGCCAGCAGC	2nd round
SARS2_2_R	TCCAAAGGCAATAGTGCAGC	1st round and 2nd round
SARS2_3_F	AAGTAGGACCTGAGCATAGTCTTG	1st round
SARS2_3_Fnest	CTTGCCGAATACCATAATGAATCTG	2nd round
SARS2_3_R	GTCTTAAGAACTCTACACCTTCT	1st round and 2nd round
SARS2_4_F	ATCTAGTTGTAATGGCCTACATTACAG	1st round
SARS2_4_iF	TTCAGTTGAYTTCGCAGTGCC	1st round
SARS2_4_Fnest	GGCTAACTAACATCTTTGGCACTG	2nd round
SARS2_4_R	CGAACTCATTTACTTCTGTACCGAG	1st round and 2nd round
SARS2_5_F	TGCCCTTGACCTAATATGATGG	1st round
SARS2_5_Fnest	ATAGAAGTGAAGTTCACAGAGTG	2nd round
SARS2_5_R	TGTTAGCAAGATTGTGTCGCT	1st round and 2nd round
SARS2_6_F	AGGAGGTGTTGCAGGAGCCT	1st round
SARS2_6_Fnest	ATAAGGCTACTAACAAATGCCATGC	2nd round
SARS2_6_R	CTTTGCCTCCTCTACAGTGTAAACC	1st round and 2nd round
SARS2_7_F	CAGATTCTGCCACTCTTGTTAGTG	1st round
SARS2_7_Fnest	CACTCTTGTTAGTGACATTGACATCAC	2nd round
SARS2_7_iFnest	TGGCACTACTGAAATGCTAGCGA	1st round
SARS2_7_R	AAGGTGATAACTTCACCATTAGGTG	1st round and 2nd round
SARS2_8_F	CACCTGATGCTGTTACAGCGT	1st round
SARS2_8_iF	GATCTCTCAAAGTGCCAGCTACAG	1st round
SARS2_8_Fnest	ACCATCTCACTTGCTGGTTCCCT	2nd round
SARS2_8_R	AAAGTGTGCCCATGTACATAACAGC	1st round and 2nd round
SARS2_9_F	CTGCTCTACAAGTCTTATTACAG	1st round
SARS2_9_Fnest	AAGACAGTAGGTGAGTTAGGTGATG	2nd round
SARS2_9_R	TCTCTTGAAGCAGGTTTCTTATAACC	1st round and 2nd round
SARS2_10_F	GTACAGAAATTGACCCTAAGTTGGACA	1st round
SARS2_10_Fnest	CCATATCCAACCGCAAGCTTCG	2nd round
SARS2_10_R	AACAGTATTCTTTGCTATAGTAGTCGG	1st round and 2nd round
SARS2_11_F	GCTACTCATGGTTAGCTGCTG	1st round
SARS2_11_Fnest	GCTGCTGTTAATAGTGTCCCTTG	2nd round
SARS2_11_R	TACATTCTAACCATAGCTGAAATCGG	1st round and 2nd round
SARS2_12_F	TGAACTCTACTAATGCTACTATTGCAAC	1st round
SARS2_12_Fnest	GCTTTTGGCTTAGTTGCAGAGT	2nd round
SARS2_12_R	TGCAACTTCCGCACTATCACC	1st round and 2nd round
SARS2_13_F	GAGCTAATAACACTAAAGKTTCAATTGC	1st round
SARS2_13_Fnest	AGCGTCTGTTTACTACAGTCAGC	2nd round
SARS2_13_R	GCGCACTACAGTCAATACAAGC	1st round and 2nd round
SARS2_14_F	GCAAGGGTTTGTGATTGATGATGTAG	1st round
SARS2_14_Fnest	ATCTGACATAGAAGTTACTGGCGATAG	2nd round
SARS2_14_R	CTGATGTTGCAAAGTCAAGTACTC	1st round and 2nd round
SARS2_15_F	TGCTGCAGTCATAACAAGAGAAG	1st round
SARS2_15_iF	AAGCTTGCCATTGATTGCTGC	1st round
SARS2_15_Fnest	GCCTGGCAGATATTACGCA	2nd round
SARS2_15_R	AAAGGTGTGAACATAACCATCCACTG	1st round and 2nd round
SARS2_16_F	GCTTTTGGTGAATACAGTYATGTAGTTG	1st round
SARS2_16_Fnest	CATTCACTGACTCTGTTAAACACCAG	2nd round
SARS2_16_R	CTTATACTTAGGTGTCTTAGGATTGGC	1st round and 2nd round
SARS2_17_F	CACCTCTGAAGACATGCTTAACC	1st round
SARS2_17_Fnest	ACAGGCTGGTAATGTTCAACTCAG	2nd round
SARS2_17_R	GAACAAAGACCATTGACTCTGGAC	1st round and 2nd round
SARS2_18_F	AGGACCTCTTCTGCTCAAACCTG	1st round
SARS2_18_iF	TTCTGCTCAAACCTGGAATTGCCG	1st round
SARS2_18_Fnest	TGCAAAATGGTATGAATGGACGTAC	2nd round
SARS2_18_iFnest	ACTGCAAAATGGTATGAATGGACGTAC	2nd round
SARS2_18_R	CCAAGAGTCAGTCTAAAGTAGCG	1st round and 2nd round
SARS2_19_F	GAGTATTGCCCTATTTTCTTCAACTG	1st round
SARS2_19_Fnest	TTCTTCAACTGGTAAATACACTTCAGTG	2nd round
SARS2_19_R	TCTAAGCATAGTGAAAAGCATTGTCTG	1st round and 2nd round
SARS2_20_F	TGAATGTGGCTAAATCTGAATTTGACC	1st round
SARS2_20_Fnest	CAGCCATGCAACGTAAGTTGG	2nd round
SARS2_20_R	CTTGTAGACGTAAGTGGCAGC	1st round and 2nd round
SARS2_21_F	GCACTGATGACAATGCGTTAGC	1st round
SARS2_21_Fnest	CTTGCACTGTTATCCGATTACAGG	2nd round

Primer ID	Sequence (5'→3')	Use
SARS2_21_R	AGACGGGCTGCACTTACACC	1st round and 2nd round
SARS2_22_F	CAAATACCTACAACCTTGTGCTAATGACC	1st round
SARS2_22_iF	TGCCGTTGCCACATAGATCATC	1st round
SARS2_22_Fnest	GGTTATGGCTGTAGTTGTGATCAAC	2nd round
SARS2_22_R	GCAGTTAAAGCCCTGGTCAAGGT	1st round and 2nd round
SARS2_22_iR	CCGAAATCATACCAGTTACCATTGAG	1st round and 2nd round
SARS2_23_F	TACGCCAACCTTAGTGAACGTG	1st round
SARS2_23_Fnest	TGATGCCATGCGAAATGCTG	2nd round
SARS2_23_R	CTGATAGCAGCATTACCATCCTG	1st round and 2nd round
SARS2_24_F	ACTAGATAAACGCACTACGTGCT	1st round
SARS2_24_iF	TAAGGAATTACTTGTGTGCTGCTG	1st round
SARS2_24_Fnest	TAGCTGCACTTACTAACAATGTTGC	2nd round
SARS2_24_iFnest	TTCTATGACTTTGCTGTGCTAAGG	2nd round
SARS2_24_R	GAGCAAGAACAAGTGAGGCCAT	1st round and 2nd round
SARS2_25_F	AATAGCCGCCACTAGAGGAG	1st round
SARS2_25_Fnest	GATTATCCTAAATGTGATAGCCATGC	2nd round
SARS2_25_R	CTATAGCTAAAGACACGAACCGTTC	1st round and 2nd round
SARS2_26_F	GCAAAATGTTGGACTGAGACTGACC	1st round
SARS2_26_Fnest	CTCAACATAACAATGCTAGTTAAACAGG	2nd round
SARS2_26_R	TGAGTCTTTTCAGTACAGGTGTTAGC	1st round and 2nd round
SARS2_27_F	TCAACTTTACTTAGGAGGTATGAGCT	1st round
SARS2_27_Fnest	CACCCATTAGTTTTCCATTGTGTGC	2nd round
SARS2_27_R	AAAGACATACTGTTCTAATGTTGAATTCAC	1st round and 2nd round
SARS2_28_F	AAGTATTCTACACTCAGGGACCAC	1st round
SARS2_28_iF	GAGCACTATGTTAGAATTACTGGCT	1st round
SARS2_28_Fnest	TACTACCCTTCTGCTCGCATAG	2nd round
SARS2_28_R	GAGCCCTGTGATGAATCAACAGT	1st round and 2nd round
SARS2_29_F	CAGGCCACAAATAGGCGTGG	1st round
SARS2_29_Fnest	CTTACACGTAACCCTGCTTGGAG	2nd round
SARS2_29_R	TCTCCAGGCGGTGGTTTAGC	1st round and 2nd round
SARS2_30_F	CCCGCGAAGAAGCTATAAGAC	1st round
SARS2_30_Fnest	CATGGATTGGCTTCGATGTCG	2nd round
SARS2_30_R	GGTTACCAATGTCGTGAAGAACTGG	1st round and 2nd round
SARS2_31_F	CCATGATCTGTATTGTCAAGTCCATG	1st round
SARS2_31_Fnest	TCTAGCTGTCCACGAGTGCT	2nd round
SARS2_31_R	CCACAAGCTAAAGCCAGCTGA	1st round and 2nd round
SARS2_32_F	GATTTGACACTAGAGTGCTATCTAACC	1st round
SARS2_32_Fnest	TAGAGTGCTATCTAACCTTAACTTGC	2nd round
SARS2_32_R	CAGTGAGTGGTGCACAAATCGT	1st round and 2nd round
SARS2_33_F	GCGCAACATTAACCAGTACCAG	1st round
SARS2_33_Fnest	ACATTGCTGCTAATACTGTGATCTG	2nd round
SARS2_33_R	CCTTAGAAACTACAGATAAATCTTGGGA	1st round and 2nd round
SARS2_34_F	TGGTTTACATCTACTGATTGGACTAGC	1st round
SARS2_34_Fnest	ATAACAGATGCGCAACAGGTTTC	2nd round
SARS2_34_R	TTATCTTTATAGCCACGGAACCTCC	1st round and 2nd round
SARS2_35_F	TTTGATTGGTGATTGTGCAACTGTAC	1st round
SARS2_35_Fnest	GGATCTCATTATTAGTGATATGTACGACC	2nd round
SARS2_35_R	TGGGTCTTCCAATCTAAAGTAGTACC	1st round and 2nd round
SARS2_36_F	CTCAGTTTTACATTTCAACTCAGGACT	1st round
SARS2_36_Fnest	TAACCCTGTCCCTACCATTTAATGATGG	2nd round
SARS2_36_R	GGTCAAGTGCACAGTCTACAGC	1st round and 2nd round
SARS2_37_F	AGTGCGTGATCTCCCTCAGG	1st round
SARS2_37_Fnest	AGGTTGGACAGCTGGTGCTG	2nd round
SARS2_37_R	AAGGTGTGCTACCGGCCTG	1st round and 2nd round
SARS2_37_iR	GTCCACAAACAGTTGCTGGTGC	1st round and 2nd round
SARS2_38_F	GAAGTCAGACAAATCGCTCCAG	1st round
SARS2_38_Fnest	CCAGATGATTTTACAGGCTGCG	2nd round
SARS2_38_R	ACTAGCGCATATRCCTGCACC	1st round and 2nd round
SARS2_39_F	CAGGAACAAATACTTCTAACCAGGTTG	1st round
SARS2_39_Fnest	GAAGTCCCTGTTGCTATTCATGC	2nd round
SARS2_39_R	TAACAGTGCAGAAAGTGTATTGAGC	1st round and 2nd round
SARS2_40_F	AGATCCATCAAACCAAGCAAGAG	1st round
SARS2_40_Fnest	GACACTTGCAGATGCTGGCT	2nd round
SARS2_40_R	CCATGAGGTGCTGACTGAGG	1st round and 2nd round
SARS2_41_F	AGGCTGAAGTGCAAATTTGATAGGT	1st round
SARS2_41_Fnest	TAGAGCTGCAGAAATCAGAGC	2nd round
SARS2_41_R	GACTCCTTTGAGCACTGGCT	1st round and 2nd round
SARS2_42_F	CTCATCGATCTCCAAGAACTTGG	1st round
SARS2_42_Fnest	GCTTGATTGCCATAGTAATGGTGAC	2nd round

Primer ID	Sequence (5'→3')	Use
SARS2_42_R	TGAGTACAGCTGGTAATAGTCTGAAG	1st round and 2nd round
SARS2_43_F	TTTGCTGGAAATGCCGTTCCA	1st round
SARS2_43_Fnest	CTTTGCTGGCATACTAATTGTTACG	2nd round
SARS2_43_R	TGTAGAAGACAATCCATGTAAGGAATAG	1st round and 2nd round
SARS2_44_F	CTTCTAGAGTTCCTGATCTTCTGG	1st round
SARS2_44_Fnest	CCATGGCAGATTCCAACGGTAC	2nd round
SARS2_44_R	GCTATAGTAACCTGAAAGTCAACGAG	1st round and 2nd round
SARS2_45_F	CAGTCGCTACAGGATTGGCA	1st round
SARS2_45_Fnest	CACAGACCATTCCAGTAGCAGTG	2nd round
SARS2_45_R	GACACGGGTCACTCAACTACATATGG	1st round and 2nd round
SARS2_46_F	TTAGGAATCATCACACTGTAGCTG	1st round
SARS2_46_Fnest	TAGCTGCATTTCCACCAAGAATGTAG	2nd round
SARS2_46_R	TGGTAGCTCTTCGGTAGTAGCC	1st round and 2nd round
SARS2_46_iR	GAAGTTGTAGCAGATTGCAGC	1st round and 2nd round
SARS2_47_F	TAACCAGAATGGAGAACGCAGTG	1st round
SARS2_47_Fnest	GGTTCACCGCTCTCACTCAAC	2nd round
SARS2_47_R	CGGCCAATGTTTGTAAATCAGTTCC	1st round and 2nd round
SARS2_48_F	CTGCTGAGGCTTCTAAGAAGC	1st round
SARS2_48_iF	GCTTGACAGATTGAACCAGCTTG	1st round
SARS2_48_Fnest	TGGCAGACGTGGTCCAGAAC	2nd round
SARS2_48_R	CTCCTRAGAAGCTATTAATCACATGG	1st round and 2nd round


Appendix Table 3. Coverage depth and number of reads mapped against the reference (GenBank accession no. NC_045512.2) for each sequencing run*

ID	Total reads mapped against SARS-CoV-2	Mean coverage depth (range)
ChVir-D712-1	102,968	971.6 (3–6,636)
ChVir-D666-2	21,192	85.8 (4–307)
ChVir-D718-3	482,720	482.9 (3–6,898)
ChVir-D672-4	52,215	214 (3–487)
ChVir-D658-5	14,485	58.9 (3–206)
ChVir-D665-6	569,265	2,342.1 (3–5,966)
ChVir-D667-7	71,902	294.9 (3–773)
ChVir-D671-8	10,051	41.3 (2–107)
ChVir-D670-9	2,561	13.3 (0–51)
ChVir-D710-10	212,223	2,273.3 (34–4,992)
ChVir-D711-11	301,653	2,924.1 (0–6,965)
ChVir-D717-D761-12	333,782	3,645.9 (3–5,927)
ChVir-W1191-13	9,892	96.6 (3–427)
ChVir-D679-14	15,850	52.6 (3–225)
ChVir-D929-15	447,129	3,791.3 (0–427)
ChVir-W1248-16	12,763	132.2 (0–422)
ChVir-D715-D799-17	444,374	4,553.4 (3–6,646)


*Coverage depth indicates the mean number of reads covering each base. Minimum and maximum number of reads are given in parentheses.

Appendix Table 4. Demographics and clinical symptoms of cases in a coronavirus disease outbreak in a nightclub, Berlin, Germany, March 2020

Demographics	No. (%)
No. total cases	74 (100)
Median age, y	30
Sex	
F	37 (50)
M	37 (50)
Clinical symptoms (no. queried)	
Dysgeusia (n = 43)	28 (65)
Cough (n = 44)	27 (61)
Headache (n = 43)	25 (58)
Dysosmia (n = 43)	25 (58)
Shivering, shaking (n = 43)	21 (49)
Myalgia (n = 43)	20 (47)
Rhinitis (n = 44)	20 (45)
Fever (n = 42)	18 (43)
Sore throat (n = 42)	12 (29)
Vertigo (n = 39)	5 (13)
Nausea (n = 42)	4 (9)



Appendix Figure 1. Illustration of the floorplan of nightclub involved in a coronavirus disease outbreak, Berlin, Germany, March 2020. Numerals represent the following: 1) entry area; 2) coat check area; 3) cashier; 4) bar counter 1; 5) lounge; 6) dance floor; 7) DJ booth; 8) bar counter 2; and 9) smoking lounge. This figure is not true to scale.



Appendix Figure 2. Maximum likelihood phylogenetic tree showing the positions of the sequences associated with a coronavirus disease outbreak in a nightclub, Berlin, Germany, March 2020. Orange circles indicate cases in the nightclub outbreak. Blue circles indicate available sequences from Germany sampled before April 15, 2020. Gray circles indicate a subset of sequences from additional countries. The x-axis shows substitutions per site. Asterisks indicate nodes with bootstrap support >70. Nodes with bootstrap support <5 are shown as polytomies. To view the sequences from club X in a wider context of currently unpublished sequences from Germany, see <https://civnb.info/sequences>.