

Survey of West Nile and Banzhi Viruses in Mosquitoes, South Africa, 2011–2018

Caitlin MacIntyre, Milehna Mara Guarido, Megan Amy Riddin, Todd Johnson, Leo Braack, Maarten Schrama, Erin Gorsich, Antonio Paulo Gouveia Almeida, Marietjie Venter

We collected >40,000 mosquitoes from 5 provinces in South Africa during 2011–2018 and screened for zoonotic flaviviruses. We detected West Nile virus in mosquitoes from conservation and periurban sites and potential new mosquito vectors; Banzhi virus was rare. Our results suggest flavivirus transmission risks are increasing in South Africa.

Flaviviruses have been major emerging zoonotic pathogens in Africa within the past decade (1). In South Africa, West Nile virus (WNV) is the main flavivirus detected in animals and humans (2,3). Several other lesser-known flaviviruses were first described in South Africa but are understudied and potentially underreported, including Wesselsbron, Usutu, and Banzhi (BANV) viruses (4). In South Africa, mosquito surveillance is not routinely performed and studies on flavivirus ecology are outdated (5). In this study, we aimed to update flavivirus vector epidemiology in northeastern provinces of South Africa through a large-scale ecologic survey.

The Study

We selected 15 sites (4 sentinel sites, 11 ad hoc sites) across 5 provinces in South Africa for mosquito collection according to recent cases of arboviral disease in humans and animals (2,3) (Figure 1). We established sentinel sites in Boschkop and Kyalami, both located in the Gauteng province (periurban sites), and Lapalala and Marakele, both located in the Lim-

popo province (conservation sites); collections were performed during 2011–2018 (Table 1). Opportunistic supplementary collections occurred at 10 ad hoc sites during 2015–2018 spanning 3 additional provinces that included urban, periurban, and conservation sites. We performed additional ad hoc collections during March–April 2017 in and around Kruger National Park (KNP) located within Limpopo and Mpumalanga provinces (periurban/conservation sites) (6). The methodologies for mosquito collection, identification, pooling, processing, flavivirus screening, and *COX1* gene sequencing have been previously described (7). To focus on months from midsummer to autumn in South Africa, when availability of mosquito breeding sites and vectorial capacity increases because of warmer and wetter weather conditions (8), we only screened mosquitoes collected during January–June. To submit sequences to GenBank, we generated *NS5* gene fragments >200 bp by using heminested PCR with 0.4 μmol/L each of forward primer (FU1, 5'-TACAACATGGGAAAGAGAGAA-3') and reverse primer (CFD2, 5'-GTGTCCCAGCCGGCGGTGTCATCAGC-3') and Platinum Taq DNA Polymerase (Thermo Fisher Scientific, <https://www.thermofisher.com>). For WNV positive pools, we performed reverse transcription PCR to amplify a 1,525 bp fragment of the WNV envelope protein gene for phylogenetic analysis (Appendix Table, <https://wwwnc.cdc.gov/EID/article/29/1/22-0036-App1.pdf>). We calculated the mosquito minimum infection rate (MIR) per site by using a standard formula: (number of positive pools/total number of individual mosquitoes tested) × 1000.

We collected a total of 66,299 mosquitoes belonging to 11 genera across 5 provinces in South Africa during 2011–2018 (Table 1; Appendix Figure 1). From those, we divided 40,731 female mosquitoes into 1,471 pools based on morphology, focusing on only 4 genera

Author affiliations: University of Pretoria, Pretoria, South Africa (C. MacIntyre, M.M. Guarido, M.A. Riddin, T. Johnson, L. Braack, M. Venter); Copperbelt University, Kitwe, Zambia (T. Johnson); Leiden University, Leiden, the Netherlands (M. Schrama); University of Warwick, Coventry, United Kingdom (E. Gorsich); NOVA University of Lisbon, Lisbon, Portugal (A.P.G. Almeida)

DOI: <https://doi.org/10.3201/eid2901.220036>




Figure 1. Sentinel and ad hoc mosquito collection sites across the northeastern region of South Africa in survey of West Nile and Banzi viruses in mosquitoes, South Africa, 2011–2018. Collection sites were selected according to recent cases of arboviral disease in humans and animals. Asterisks in the color-coded figure legend indicate sites where flaviviruses were identified in mosquitoes.

(*Culex*, *Mansonia*, *Anopheles*, and *Aedes*) and screened for flaviviruses. We detected WNV in 16 (1.09%) and BANV in 2 (0.14%) pools. We did not detect other zoonotic flaviviruses; however, insect-specific flaviviruses were detected and described elsewhere (7). WNV outbreaks can be expected once the MIR rises above 1 (9). We observed the highest MIRs in Kyalami (periurban site, MIR = 2.53), KNP (periurban/conservation site, MIR = 1.22), and Lapalala (conservation site, MIR = 1.01) (Table 2). Therefore, we identified those areas as higher risk sites for WNV outbreaks in humans and animals. These results correlated with areas where *Culex* spp. mosquitoes were the most abundant and where WNV cases were previously reported in humans and animals in South Africa (3,10).

Only 11 of 16 WNV-positive pools had partial *NS5* gene sequences of sufficient quality to perform maximum-likelihood analysis; we confirmed all 11 pools were WNV and also confirmed 2 BANV-positive pools (Figure 2; bootstrap value = 100 for both viruses). We observed high nucleotide similarity (94.62%–100.00%) between the identified WNV *NS5* gene sequences and those from previously identified, highly neuroinvasive strains from South Africa isolated from either equines or humans. We successfully amplified the 1,525 nt region of the envelope protein gene for 5 of 16 WNV-positive pools (Appendix Figure 2).

We performed *COX1* gene sequencing for 9 of 16 WNV-positive pools and confirmed morphologic identification of those mosquitoes as *Cx. univittatus*

except for 1 pool collected in KNP that was identified by sequencing as *Cx. perexiguus* (Table 2; Appendix Figure 3), a mosquito species not known to be present in South Africa (11). This 1 pool might have contained a mix of both species, but Sanger sequencing was unable to distinguish between the 2 species. In addition, the *COX1* reference sequences for *Cx. perexiguus* mosquitoes obtained from online databases may not be accurate because this species has not been identified in South Africa. Recently, *COX1* gene amplification using universal primers followed by next-generation sequencing was shown to distinguish between species in mixed pools (12) and might be useful in future studies to resolve this ambiguity. Further studies are necessary to clarify the status of the *Cx. perexiguus* mosquitoes in South Africa. Most of the WNV-positive pools consisted of *Cx. univittatus*, *Cx. pipiens s.l.*, and *Cx. theileri* mosquitoes, which we collected in high abundance. This result reiterates the importance of these species as WNV vectors in South Africa (5). We identified *Cx. simpsoni*, *Cx. bitaeniorhynchus*, *An. gambiae sensu lato*, and *Cx. poicilipes* mosquitoes, none of which have been previously associated with WNV in South Africa, as new potential vectors for WNV by using *COX1* gene sequencing (4). Globally, from this list of species, only *Cx. poicilipes* mosquitoes from Senegal were found to be infected with WNV (13). Experimental studies have shown that the *Cx. bitaeniorhynchus* mosquito is a likely vector

Table 1. Total number of mosquitoes collected across sentinel and ad hoc sites in survey of West Nile and Banzi viruses in mosquitoes in South Africa during 2011–2018

Site	Site type	Province	Region type	Coordinates	No. mosquitoes collected	No. trapping events	Mean no. mosquitoes/trapping event
Boschkop	Sentinel	Gauteng	Peri-urban	-25.82786, 28.42047	4,790	353	12
Kyalami	Sentinel	Gauteng	Peri-urban	-25.99183, 28.02947	8,736	338	26
Lapalala	Sentinel	Limpopo	Conservation	-23.88458, 28.26953	16,675	568	29
Marakele	Sentinel	Limpopo	Conservation	-24.29364, 27.50325	13,347	487	24
Total nos., sentinel sites					43,548	1,746	23
Benoni	Ad hoc	Gauteng	Peri-urban	-26.10611, 28.36689	1,191	31	38
Jozini	Ad hoc	KwaZulu-Natal	Rural	-27.41258, 32.20647	10,954	47	233
Kruger National Park	Ad hoc	Limpopo, Mpumalanga	Conservation, peri-urban	-25.35384, 31.79936	2,440	30	58
Matikwane	Ad hoc	Mpumalanga	Urban	-24.98545, 31.236342	615	9	68
Mnisi	Ad hoc	Mpumalanga	Rural	-24.48206, 31.38583	5,549	171	32
Pretoria north	Ad hoc	Gauteng	Urban	-25.68663, 28.15895	219	23	12
Roodeplaat	Ad hoc	Gauteng	Peri-urban	-25.62075, 28.37136	298	15	20
Shingwedzi	Ad hoc	Limpopo	Conservation	-23.10819, 31.43628	457	13	35
Skukuza	Ad hoc	Mpumalanga	Conservation	-24.99633, 31.59189	482	15	32
Southern African Wildlife College	Ad hoc	Mpumalanga	Conservation	-24.53886, 31.33369	56	9	15
Vulpro	Ad hoc	Northwest	Peri-urban	-25.7112, 27.95322	490	12	41
Total nos., ad hoc sites					22,751	375	53

for WNV because this species was able to successfully transmit WNV (14). We were unable to genetically characterize the remaining 7 of 16 pools

because of insufficient material for DNA extraction; we identified mosquitoes in those pools by morphologic characteristics.

Table 2. Detection of flaviviruses and their associated potential mosquito vectors in survey of West Nile and Banzi viruses in mosquitoes, South Africa, 2011–2018*

Site	Virus	Positive pool ID	No. mosquitoes†	Morphologic ID‡	Molecular ID§	MIR
Boschkop	WNV	GAU11MP26	2	<i>Culex pipiens sensu lato</i>	Not done	0.72
	WNV	GAU17MP72	33	<i>Cx. univittatus</i>	<i>Cx. univittatus</i>	
Kruger National Park	WNV	KNP17MP714	4	<i>Cx. simpsoni</i>	<i>Cx. simpsoni</i>	1.23
	WNV	KNP17MP718	36	<i>Cx. univittatus</i>	<i>Cx. perexiguus</i>	
	WNV	KNP17MP720	1	<i>Cx. bitaeniorhynchus</i>	<i>Cx. bitaeniorhynchus</i>	
Kyalami	WNV	KYA11MP11	14	<i>Cx. univittatus</i>	Not done	2.53
	WNV	KYA11MP13	10	<i>Cx. pipiens s.l.</i>	Not done	
	WNV	KYA14MP133	10	<i>Cx. univittatus</i>	<i>Cx. univittatus</i>	
	WNV	KYA14MP134	19	<i>Cx. univittatus</i>	<i>Cx. univittatus</i>	
	WNV	KYA14MP115	5	<i>Cx. theileri</i>	<i>Cx. theileri</i>	
Lapalala	WNV	LAP13LP71	44	<i>Anopheles</i> spp.	Not done	1.01
	WNV	LAP13LP28	50	<i>Anopheles</i> spp.	Not done	
	WNV	LAP13LP22	50	<i>Aedes</i> spp.	Not done	
	WNV	LAP14MP394	2	<i>Cx. univittatus</i>	<i>Cx. univittatus</i>	
Marakele	WNV	MAR13MP77	50	<i>Cx. poecilipes</i>	Not done	0.33
	WNV	MAR15MP18	1	<i>An. gambiae s.l.</i>	<i>An. gambiae s.l.</i>	
Lapalala	BANV	LAP13MP25	49	<i>Cx. spp.</i>	<i>Cx. rubinotus</i>	0.83
	BANV	LAP13MP26	50	<i>Cx. spp.</i>	<i>Cx. annulioris</i>	

*BANV, Banzi virus; ID, identification; MIR, minimum infection rate; WNV, West Nile virus.

†Number of mosquitoes in each positive pool.

‡Species of mosquito identified by morphologic characteristics.

§Species of mosquito identified by sequencing the COX1 gene.




Figure 2. Phylogenetic analysis of flaviviruses using *NS5* gene sequences in survey of West Nile and Banzi viruses in mosquitoes, South Africa, 2011–2018. Maximum likelihood analysis was used to identify flaviviruses found in mosquitoes after partial sequencing of the flavivirus *NS5* gene region (226 nt, Kimura 2-parameter model plus gamma distribution plus proportion of invariable sites). Sequence data were edited by using CLC Main Workbench version 8.0.1 (QIAGEN, <https://www.qiagen.com>). Reference genomes were downloaded from GenBank. Multiple sequence alignments were created by using MAFFT (<http://mafft.cbrc.jp/alignment/server/index.html>) with default parameters. Phylogenetic analysis was performed by using MEGA X software (MEGA, <https://www.megasoftware.net>) with bootstrap support for network groupings calculated from 1,000 replicates. Bootstrap values (>70%) are displayed on branches. GenBank accession numbers for newly sequenced virus strains: OL411950 (KYA11MP13 isolate), OL411951 (GAU11MP26 isolate), OL411952 (KYA14MP133 isolate), OL411953 (KYA14MP134 isolate), OL411954 (LAP14MP394 isolate), OL411955 (MAR15MP18 isolate), OL411956 (LAP13MP22 isolate), OL411957 (KNP17MP714 isolate), OL411958 (KNP17MP720 isolate), OL411959 (KNP17MP718 isolate), OL411960 (KYA11MP11 isolate), OL411961 (LAP13MP25 isolate), and OL411962 (LAP13MP26 isolate). Solid black triangles are new viral sequences that were detected in mosquitoes in this study. Scale bar indicates nucleotide substitutions per site.

Detection of BANV in mosquito pools in South Africa has not been described since the late 1970s (15), and a lack of surveillance raises the question regarding the true incidence of this virus. Only the *Cx. rubinotus* mosquito is recognized as a vector for BANV (15). Despite unclear morphologic identification, we identified *Cx. rubinotus* and *Cx. annulioris* mosquitoes in the 2 BANV-positive pools through *COX1* gene sequencing (Table 2; Appendix Figure 3), which should be investigated further to confirm vector status.

Conclusions

The first limitation of our study is that we did not separate voucher specimens for mosquito species from the pools before homogenization. A voucher specimen is a mosquito species that is preserved and serves as a reference used to document identity. Second, identifications of mosquito species not previously associated with WNV infection are preliminary findings, and further investigation of vector competency is required.

Mosquito surveillance is not routinely performed for arboviruses in South Africa, and most studies were performed 40 years ago (5). In this study, 5 provinces were targeted for mosquito surveillance over a 7-year period. These investigations revealed a wide range of new potential vectors that require further investigation. Both WNV and BANV were identified in mosquitoes in periurban and conservation areas at the animal/human interface in South Africa, suggesting increasing circulation potential for those viruses between humans, wildlife, domestic animals, and avian species that are common in those areas.

Acknowledgments

We thank Lapalala Wilderness, Marataba Conservation, and South African National Parks for logistical assistance and permission to collect mosquitoes; Basil Brooke and Danny Govender for their contribution to the collection of the mosquitoes; all previous staff and students in the Zoonotic Arbo- and Respiratory virus program for their technical support; and Clarence Yah for his assistance in preparing the manuscript. A.P.G. Almeida has been a recipient of the Visiting Professor Programme of the University of Pretoria and acknowledges the Global Health and Tropical Medicine unit.

The study was funded in part by a cooperative agreement (no. 5 NU2GGH001874-02-00) with the US Centers for Disease Control and Prevention. The study was solely the responsibility of the authors and does not necessarily represent the official views of the US Centers for Disease Control and Prevention or US Department of Health and Human Services.

About the Author

Ms. MacIntyre is an MSc candidate in the Zoonotic Arbo- and Respiratory virus research group, Department of Medical Virology, at the University of Pretoria. Her research focuses on One Health investigations of West Nile virus in South Africa.

References

- Hollidge BS, González-Scarano F, Soldan SS. Arboviral encephalitides: transmission, emergence, and pathogenesis. *J Neuroimmune Pharmacol.* 2010;5:428–42. <https://doi.org/10.1007/s11481-010-9234-7x>
- Burt FJ, Grobbelaar AA, Leman PA, Anthony FS, Gibson GV, Swanepoel R. Phylogenetic relationships of southern African West Nile virus isolates. *Emerg Infect Dis.* 2002;8:820–6. <https://doi.org/10.3201/eid0808.020027>
- Venter M, Pretorius M, Fuller JA, Botha E, Rakgotho M, Stivaktas V, et al. West Nile virus lineage 2 in horses and other animals with neurologic disease, South Africa, 2008–2015. *Emerg Infect Dis.* 2017;23:2060–4. <https://doi.org/10.3201/eid2312.162078>
- Venter M. Assessing the zoonotic potential of arboviruses of African origin. *Curr Opin Virol.* 2018;28:74–84. <https://doi.org/10.1016/j.coviro.2017.11.004>
- Jupp PG. The ecology of West Nile virus in South Africa and the occurrence of outbreaks in humans. *Ann N Y Acad Sci.* 2001;951:143–52. <https://doi.org/10.1111/j.1749-6632.2001.tb02692.x>
- Gorsich EE, Beechler BR, van Bodegom PM, Govender D, Guarido MM, Venter M, et al. A comparative assessment of adult mosquito trapping methods to estimate spatial patterns of abundance and community composition in southern Africa. *Parasit Vectors.* 2019;12:462. <https://doi.org/10.1186/s13071-019-3733-z>
- Guarido MM, Govender K, Riddin MA, Schrama M, Gorsich EE, Brooke BD, et al. Detection of insect-specific flaviviruses in mosquitoes (Diptera: *Culicidae*) in northeastern regions of South Africa. *Viruses.* 2021;13:2148. <https://doi.org/10.3390/v13112148>
- Cornel AJ, Lee Y, Almeida APG, Johnson T, Mouatcho J, Venter M, et al. Mosquito community composition in South Africa and some neighboring countries. *Parasit Vectors.* 2018;11:331. <https://doi.org/10.1186/s13071-018-2824-6>
- Bustamante DM, Lord CC. Sources of error in the estimation of mosquito infection rates used to assess risk of arbovirus transmission. *Am J Trop Med Hyg.* 2010;82:1172–84. <https://doi.org/10.4269/ajtmh.2010.09-0323>
- Zaayman D, Venter M. West Nile virus neurologic disease in humans, South Africa, September 2008–May 2009. *Emerg Infect Dis.* 2012;18:2051–4. <https://doi.org/10.3201/eid1812.111208>
- Mixão V, Bravo Barriga D, Parreira R, Novo MT, Sousa CA, Frontera E, et al. Comparative morphological and molecular analysis confirms the presence of the West Nile virus mosquito vector, *Culex univittatus*, in the Iberian Peninsula. *Parasit Vectors.* 2016;9:601. <https://doi.org/10.1186/s13071-016-1877-7>
- Alkan C, Erisoz Kasap O, Alten B, de Lamballerie X, Charrel RN. Sandfly-borne phlebovirus isolations from Turkey: new insight into the sandfly fever Sicilian and sandfly fever Naples species. *PLoS Negl Trop Dis.* 2016;10:e0004519. <https://doi.org/10.1371/journal.pntd.0004519>


13. Hubálek Z, Halouzka J. West Nile fever – a reemerging mosquito-borne viral disease in Europe. *Emerg Infect Dis.* 1999;5:643–50. <https://doi.org/10.3201/eid0505.990505>
14. Ilkal MA, Mavale MS, Prasanna Y, Jacob PG, Geevarghese G, Banerjee K. Experimental studies on the vector potential of certain *Culex* species to West Nile virus. *Indian J Med Res.* 1997;106:225–8.
15. Jupp PG, McIntosh BM, Anderson D. *Culex (Eumelanomyia) rubinotus* Theobald as vector of Banzi, Germiston and Witwatersrand viruses. IV. Observations on the biology of *C. rubinotus*. *J Med Entomol.* 1976;12:647–51. <https://doi.org/10.1093/jmedent/12.6.647>

Address for correspondence: Marietjie Venter, Zoonotic Arbo- and Respiratory Virus Program, Centre for Viral Zoonosis, Department of Medical Virology, Pathology Building, Prinshof Campus South, University of Pretoria, Private Bag X323, Gezina 0031, South Africa; email: marietjie.venter@up.ac.za

May 2022

Viral Infections

- Invasive Group A *Streptococcus* Outbreaks Associated with Home Healthcare, England, 2018–2019
- Genomic Epidemiology of Global Carbapenemase-Producing *Escherichia coli*, 2015–2017
- Risk for Asymptomatic Household Transmission of *Clostridioides difficile* Infection Associated with Recently Hospitalized Family Members
- Estimating Relative Abundance of 2 SARS-CoV-2 Variants through Wastewater Surveillance at 2 Large Metropolitan Sites, United States
- Effectiveness of BNT162b2 Vaccine Booster against SARS-CoV-2 Infection and Breakthrough Complications, Israel
- Effects of Tick-Control Interventions on Tick Abundance, Human Encounters with Ticks, and Incidence of Tickborne Diseases in Residential Neighborhoods, New York, USA
- Pertactin-Deficient *Bordetella pertussis* with Unusual Mechanism of Pertactin Disruption, Spain, 1986–2018
- Determining Existing Human Population Immunity as Part of Assessing Influenza Pandemic Risk
- Disparities in First Dose COVID-19 Vaccination Coverage among Children 5–11 Years of Age, United States
- Multisystem Inflammatory Syndrome in Children after SARS-CoV-2 Vaccination
- Severe Multisystem Inflammatory Symptoms in 2 Adults after Short Interval between COVID-19 and Subsequent Vaccination



- Evidence of Prolonged Crimean-Congo Hemorrhagic Fever Virus Endemicity by Retrospective Serosurvey, Eastern Spain
- Lack of Evidence for Crimean–Congo Hemorrhagic Fever Virus in Ticks Collected from Animals, Corsica, France
- Highly Pathogenic Avian Influenza A(H5N8) Clade 2.3.4.4b Viruses in Satellite-Tracked Wild Ducks, Ningxia, China, 2020
- Novel Hendra Virus Variant Circulating in Black Flying Foxes and Grey-Headed Flying Foxes, Australia
- Increased COVID-19 Severity among Pregnant Patients Infected with SARS-CoV-2 Delta Variant, France
- Mathematical Modeling for Removing Border Entry and Quarantine Requirements for COVID-19, Vanuatu
- SARS-CoV-2 Seroprevalence after Third Wave of Infections, South Africa
- *Angiostrongylus cantonensis* in a Red Ruffed Lemur at a Zoo, Louisiana, USA
- Breast Milk as Route of Tick-Borne Encephalitis Virus Transmission from Mother to Infant
- *atpE* Mutation in *Mycobacterium tuberculosis* Not Always Predictive of Bedaquiline Treatment Failure
- Emerging Novel Reassortant Influenza A(H5N6) Viruses in Poultry and Humans, China, 2021
- *Mycobacterium lepromatosis* as Cause of Leprosy, Colombia
- Rare Case of Rickettsiosis Caused by *Rickettsia monacensis*, Portugal, 2021
- Pathogens that Cause Illness Clinically Indistinguishable from Lassa Fever, Nigeria, 2018
- Duration of Infectious Virus Shedding by SARS-CoV-2 Omicron Variant–Infected Vaccinees
- Imported Monkeypox from International Traveler, Maryland, USA, 2021
- Intercontinental Movement of Highly Pathogenic Avian Influenza A(H5N1) Clade 2.3.4.4 Virus to the United States, 2021
- Rapid Replacement of SARS-CoV-2 Variants by Delta and Subsequent Arrival of Omicron, Uganda, 2021
- SARS-CoV-2 Antibody Prevalence and Population-Based Death Rates, Greater Omdurman, Sudan
- Cross-Variant Neutralizing Serum Activity after SARS-CoV-2 Breakthrough Infections

**EMERGING
INFECTIOUS DISEASES**

To revisit the May 2022 issue, go to:
<https://wwwnc.cdc.gov/eid/articles/issue/28/5/table-of-contents>