

differences in hematological and biochemical parameters between negative and positive animals indicate the asymptomatic features of infection and make clarifying the baboons' role as a potential reservoir more urgent. Measures to control tick circulation should be established to reduce risk for transmission of *R. conorii* subsp. *israelensis* to zoopark visitors and workers.

Acknowledgments

The authors are grateful to the staff of Fasano Zoopark involved in field activities.

R.I. and D.O. were partially supported by EU funding within the NextGeneration EU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (Project no. PE00000007, INF-ACT).

About the Author

Mr. Sgroi has a PhD in animal health and zoonosis from the University of Bari Aldo Moro. His main research activities focus on biology, epidemiology, and control of vector-borne pathogens of zoonotic concern.

References

- Nakayama J, Hayashida K, Nakao R, Ishii A, Ogawa H, Nakamura I, et al. Detection and characterization of zoonotic pathogens of free-ranging non-human primates from Zambia. *Parasit Vectors*. 2014;7:490. <https://doi.org/10.1186/s13071-014-0490-x>
- Sgroi G, Iatta R, Lia RP, D'Alessio N, Manoj RRS, Veneziano V, et al. Spotted fever group rickettsiae in *Dermacentor marginatus* from wild boars in Italy. *Transbound Emerg Dis*. 2021;68:2111–20. <https://doi.org/10.1111/tbed.13859>
- Roux V, Raoult D. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer-membrane protein rOmpB (ompB). *Int J Syst Evol Microbiol*. 2000;50:1449–55. <https://doi.org/10.1099/00207713-50-4-1449>
- Sgroi G, Iatta R, Veneziano V, Bezerra-Santos MA, Lesiczka P, Hrazdilová K, et al. Molecular survey on tick-borne pathogens and *Leishmania infantum* in red foxes (*Vulpes vulpes*) from southern Italy. *Ticks Tick Borne Dis*. 2021;12:101669. <https://doi.org/10.1016/j.ttbdis.2021.101669>
- Maia C, Cristóvão JM, Pereira A, Parreira R, Campino L. Detection of *Rickettsia conorii israelensis* DNA in the blood of a cat and a dog from southern Portugal. *Top Companion Anim Med*. 2019;36:12–5. <https://doi.org/10.1053/j.tcam.2019.06.001>
- Guccione C, Colomba C, Rubino R, Bonura C, Anastasia A, Agrenzano S, et al. A severe case of Israeli spotted fever with pleural effusion in Italy. *Infection*. 2022;50:269–72. <https://doi.org/10.1007/s15010-021-01693-8>
- Torina A, Naranjo V, Pennisi MG, Patania T, Vitale F, Laricchiuta P, et al. Serologic and molecular characterization of tickborne pathogens in lions (*Panthera leo*) from the Fasano Safari Park, Italy. *J Zoo Wildl Med*. 2007;38:591–3. <https://doi.org/10.1638/2007-0043R1.1>
- Rovero C, Brouqui P, Raoult D. Questions on Mediterranean spotted fever a century after its discovery. *Emerg Infect Dis*. 2008;14:1360–7. <https://doi.org/10.3201/eid1409.071133>
- Akinyi MY, Tung J, Jeneby M, Patel NB, Altmann J, Alberts SC. Role of grooming in reducing tick load in wild baboons (*Papio cynocephalus*). *Anim Behav*. 2013;85:559–68. <https://doi.org/10.1016/j.anbehav.2012.12.012>
- Otranto D, Dantas-Torres F, Giannelli A, Latrofa MS, Cascio A, Cazzin S, et al. Ticks infesting humans in Italy and associated pathogens. *Parasit Vectors*. 2014;7:328. <https://doi.org/10.1186/1756-3305-7-328>

Address for correspondence: Domenico Otranto, University of Bari Aldo Moro, strada provinciale per Casamassima km3, 70010 Valenzano, Italy; email: domenico.otranto@uniba.it

Prevention of *Thelazia callipaeda* Reinfection among Humans

Marija Trenkić, Suzana Tasić-Otašević, Marcos Antonio Bezerra-Santos, Marko Stalević, Aleksandar Petrović, Domenico Otranto

Author affiliations: Ophthalmology Clinic University Clinical Center, Niš, Serbia (M. Trenkić); University of Niš, Niš (M. Trenkić, S. Tasić-Otašević, M. Stalević, A. Petrović); Public Health Institute, Niš (S. Tasić-Otašević); University of Bari Aldo Moro, Bari, Italy (M.A. Bezerra-Santos, D. Otranto); Bu-Ali Sina University, Hamedan, Iran (D. Otranto)

DOI: <https://doi.org/10.3201/eid2904.221610>

Thelazia callipaeda is a zoonotic vector-borne nematode that infects and causes eye disease among a wide range of domestic and wild mammals, including humans. We describe an unusual case of reinfection by this nematode in Serbia and call for a focus on preventive measures in endemic areas.

The genus *Thelazia* (order Spirurida, family Thelaziidae) comprises several species of nematode that cause ocular infections in different host mammals, including humans (1). Over the past 20 years, the *T. callipaeda* eyeworm has gained interest among

Table. Morphometry of 11 specimens (6 female, 5 male) of *Thelazia callipaeda* from a naturally infected human in Serbia

Measurement	Male	Female
Body length, mm	9.58–12.7 (mean 11.05; SD ±1.16)	13.5–16.5 (mean 14.96; SD ±1.23)
Body width, μm	310–390 (mean 343; SD ±31.14)	360–490 (mean 434; SD ±56.83)
Buccal capsule length, μm	26.4	29.6
Buccal capsule width, μm	25.1	26.4
Nerve-ring from anterior extremity, μm	267.5	373.7
Esophagus length, μm	665.4	719.9
Vulva from anterior extremity, μm	Not applicable	663.2
Left spicule length, μm	1,648	NA
Right spicule length, μm	157.2	NA
Tail length, μm	62.8	72.4

the scientific community because several human cases have been reported in countries in Asia and Europe, making this parasite an agent of public health concern (1). Adult and larval forms of *T. callipaeda* eyeworms infect the ocular apparatus of a wide range of both domestic (e.g., dogs, cats) and wild (e.g., red foxes, wolves, jackals, bears, lagomorphs) animal species, including humans (1–3). In Europe, 2 species of drosophilid fruit flies were confirmed to act as vectors of *T. callipaeda* eyeworms: *Phortica variegata*, tested under laboratory and natural conditions (4), and *P. oldenbergi* only experimentally (5). Thelaziosis should be expected among humans in areas where *T. callipaeda* infection is endemic in animal reservoirs. Absence of preventive measures could be a factor influencing human reinfections in endemic areas. We describe a case of *T. callipaeda* reinfection in a human patient and call for focus on prevention in areas where the parasite and its vectors thrive.

A man, 41 years of age, living in a small village in the southern part of Serbia contacted an ophthalmologist at the University Clinical Center in Nis, Serbia,

because of ocular discomfort. The patient reported a history of clinical thelaziosis caused by *T. callipaeda* infection 5 years earlier. Ophthalmic manifestations were conjunctivitis with increased lacrimation, itching, and sensation of a foreign body in his left eye. During ophthalmologic examination, we removed 11 eyeworms (6 female, 5 male) from the eye and subsequently identified them morphologically as *T. callipaeda* according to an identification key (6). The nematodes had a filariform body type with a transversally striated cuticle and a buccal capsule of hexagonal shape. Among female worms, the vulva was located anterior to the esophagus-intestinal junction; male worms had a curved caudal end with 2 asymmetric spicules and precloacal and postcloacal papillae (Table; Figure).

To confirm morphologic identification, we extracted genomic DNA from individual worms using a QIAGEN DNeasy Blood & Tissue Kit (<https://www.qiagen.com>), and performed PCR analysis using the primers NTF (5'-TGATTGGTGGTTTTGGTAA-3') and NTR (5'-ATAAGTACGAGTATCAATATC-3'), which amplify a 689-bp portion of the mitochondrial




Figure. *Thelazia callipaeda* eyeworms collected from the left eye of a man in Serbia. A) Female worm; B) anterior end of adult female; C) posterior end of adult female; D) posterior end of adult male.

cytochrome *c* oxidase subunit 1 (*cox1*) gene. We purified amplified DNA products and sequenced them in both directions using ThermoFisher Big Dye Terminator version 3.1 chemistry in an Applied Biosystems 3130 genetic analyzer with an ABI-PRISM 377 automated sequencer (<https://www.thermofisher.com>). We analyzed sequences using MEGA version 7 software (<https://www.megasoftware.net>) and compared them with those available in GenBank using BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Nucleotide sequences had 100% identity with *T. callipaeda* (GenBank accession no. AM042549.1). In addition, phylogenetic analysis performed by using the maximum-likelihood method based on the Tamura-Nei model showed the representative sequence from our study clustered with other sequences of *T. callipaeda* belonging to haplotype 1 (Appendix, <https://wwwnc.cdc.gov/EID/article/29/4/22-1610-App1.pdf>), the only haplotype thus far described in Europe. We deposited the nucleotide sequence in GenBank (accession no. OP696980).

We treated the patient with topical antimicrobials and corticosteroids (rinsing with 3% boric acid 5×/d and topical tobramycin/deksametazon 5×/d). At clinical follow-up 7 and 14 days later, we found no signs or symptoms of eye infection.

T. callipaeda eyeworm prevalence in humans and animals has increased throughout Europe in recent decades (1). To date, human thelaziosis has been described in 12 patients from Europe, including a case-patient in Serbia (7) reinfected by *T. callipaeda* eyeworms 5 years after an initial case, as in the case we describe here. A related study called for implementing preventive measures, such as vector control and treatment of domestic reservoirs (e.g., dogs), to avoid zoonotic human infection (8). In addition, wild carnivore reservoirs, such as red foxes and wolves, should be considered as sources of infection for humans who frequent the same forest areas (9). The patient in our study reported that he spent long periods picking mushrooms in the forest, and he exhibited clinical manifestations of thelaziosis during the summer (July), when outdoor activities are most common and the *P. variegata* fruit fly, a *T. callipaeda* eyeworm vector, is most abundant.

Reinfection in this patient highlights that *T. callipaeda* eyeworms can cause recurrent infection in human hosts, which suggests the potential usefulness of implementing prevention and control strategies in the Balkan Peninsula, where this parasite and its vector and animal reservoirs are spreading (10). Moreover, it indicates that inspecting for *T. callipaeda* eyeworms should be part of routine periodic examinations in endemic areas, even among asymptomatic persons.

Acknowledgments

The authors thank Riccardo Paolo Lia for his support on morphologic identification and Giada Annoscia for her support on molecular analysis.

About the Author

Dr. Trenkić is an assistant professor in the medical faculty, Department of Ophthalmology, University of Niš. She performs ophthalmologic research.

References

- Otranto D, Mendoza-Roldan JA, Dantas-Torres F. *Thelazia callipaeda*. Trends Parasitol. 2021;37:263–4. <https://doi.org/10.1016/j.pt.2020.04.013>
- Papadopoulos E, Komnenou A, Karamanlidis AA, Bezerra-Santos MA, Otranto D. Zoonotic *Thelazia callipaeda* eyeworm in brown bears (*Ursus arctos*): a new host record in Europe. Transbound Emerg Dis. 2022;69:235–9. <https://doi.org/10.1111/tbed.14414>
- Bezerra-Santos MA, Moroni B, Mendoza-Roldan JA, Perrucci S, Cavicchio P, Cordon R, et al. Wild carnivores and *Thelazia callipaeda* zoonotic eyeworms: a focus on wolves. Int J Parasitol Parasites Wildl. 2022;17:239–43. <https://doi.org/10.1016/j.ijppaw.2022.03.005>
- Otranto D, Brianti E, Cantacessi C, Lia RP, Máca J. The zoophilic fruitfly *Phortica variegata*: morphology, ecology and biological niche. Med Vet Entomol. 2006;20:358–64. <https://doi.org/10.1111/j.1365-2915.2006.00643.x>
- Bezerra-Santos MA, Bernardini I, Lia RP, Mendoza-Roldan JA, Beugnet F, Pombi M, et al. *Phortica oldenbergi* (Diptera: Drosophilidae): a new potential vector of the zoonotic *Thelazia callipaeda* eyeworm. Acta Trop. 2022;233:106565. <https://doi.org/10.1016/j.actatropica.2022.106565>
- Otranto D, Lia RP, Traversa D, Giannetto S. *Thelazia callipaeda* (Spirurida, Thelaziidae) of carnivores and humans: morphological study by light and scanning electron microscopy. Parasitologia. 2003;45:125–33.
- Tasić-Otašević S, Gabrielli S, Trenkić-Božinović M, Petrović A, Gajić B, Colella V, et al. Eyeworm infections in dogs and in a human patient in Serbia: a One Health approach is needed. Comp Immunol Microbiol Infect Dis. 2016;45:20–2. <https://doi.org/10.1016/j.cimid.2016.01.003>
- Bezerra-Santos MA, Mendoza-Roldan JA, Sgroi G, Lia RP, Venegoni G, Solari Basano F, et al. Efficacy of a formulation of sarolaner/moxidectin/pyrantel (Simparica Trio®) for the prevention of *Thelazia callipaeda* canine eyeworm infection. Parasit Vectors. 2022;15:370. <https://doi.org/10.1186/s13071-022-05501-6>
- Čabanová V, Miterpáková M, Oravec M, Hurníková Z, Jerg S, Nemčíková G, et al. Nematode *Thelazia callipaeda* is spreading across Europe. The first survey of red foxes from Slovakia. Acta Parasitol. 2018;63:160–6. <https://doi.org/10.1515/ap-2018-0018>
- Hodžić A, Latrofa MS, Annoscia G, Alić A, Beck R, Lia RP, et al. The spread of zoonotic *Thelazia callipaeda* in the Balkan area. Parasit Vectors. 2014;7:352. <https://doi.org/10.1186/1756-3305-7-352>

Address for correspondence: Domenico Otranto, University of Bari Aldo Moro, Strada provinciale per Casamassima km3, 70010 Valenzano, Italy; email: domenico.otranto@uniba.it